

SMALL DATA SCATTERING FOR A SYSTEM OF NONLINEAR SCHRÖDINGER EQUATIONS

NAKAO HAYASHI, CHUNHUA LI AND TOHRU OZAWA

Abstract. We study the scattering theory for a system of nonlinear Schrödinger equations in space dimension $n \geq 3$. In the case $n \geq 4$, existence of the scattering operator is proved in small data setting in the Sobolev space $H^{n/2-2}$. In the case $n = 3$, a similar result is proved in the weighted L^2 space $\langle x \rangle^{-1/2} L^2 = \mathcal{F}(H^{-1/2})$ under the mass resonance condition.

Mathematics subject classification (2010): 35Q55.

Keywords and phrases: nonlinear Schrödinger equations, scattering theory.

REFERENCES

- [1] A. BACHELOT, *Problème de Cauchy global pour des systèmes de Dirac-Klein-Gordon*, Ann. Inst. Henri Poincaré, **48** (1988), 387–422.
- [2] J. BERGH AND J. LÖFSTRÖM, *Interpolation Spaces*, Springer, 1976.
- [3] T. CAZENAVE, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics vol. **10**, American Mathematical Society (2003).
- [4] T. CAZENAVE AND F. B. WEISSLER, The Cauchy problem for the critical nonlinear Schrödinger equation in H^s , *Nonlinear Analysis TMA*, **14** (1990), 807–836.
- [5] M. COLIN AND T. COLIN, *On a quasilinear Zakharov system describing laser-plasma interactions*, Diff. Integral Equations, **17** (2004), 297–330.
- [6] V. GEORGIEV, *Global solution of the system of wave and Klein-Gordon equations*, Math. Z., **203** (1990), 683–698.
- [7] N. HAYASHI, M. IKEDA AND P. I. NAUMKIN, Wave operator for the system of the Dirac-Klein-Gordon equations, to appear in *Math. Method. Appl. Sci.*
- [8] N. HAYASHI, C. LI AND P. I. NAUMKIN, *On a system of nonlinear Schrödinger equations in 2d*, preprint, *Differential Integral Equations*, **24** (2011), 417–434.
- [9] N. HAYASHI, C. LI AND P. I. NAUMKIN, *Modified wave operator for a system of nonlinear Schrödinger equations in 2d*, preprint.
- [10] N. HAYASHI AND T. OZAWA, *Scattering theory in the weighted $L^2(\mathbb{R}^n)$ spaces for some Schrödinger equations*, Ann. Inst. H. Poincaré Phys., **48** (1988), 17–37.
- [11] T. KATO, *On nonlinear Schrödinger equations II, H^s -solutions and nonconditional well-posedness*, J. Anal. Math., **67** (1995), 281–306.
- [12] M. KEEL AND T. TAO, *Endpoint Strichartz estimates*, Amer. J. Math., **120** (1998), 955–980.
- [13] S. MACHIHARA, K. NAKANISHI AND T. OZAWA, *Nonrelativistic limit in the energy space for nonlinear Klein-Gordon equations*, Math. Ann., **332** (2002), 603–621.
- [14] S. MACHIHARA, K. NAKANISHI AND T. OZAWA, *Small global solutions and the nonrelativistic limit for the nonlinear Dirac equation*, Rev. Math. Iberoamericana, **19** (2003), 179–194.
- [15] M. NAKAMURA AND T. OZAWA, *Low energy scattering for nonlinear Schrödinger equations in fractional order Sobolev spaces*, Rev. Math. Phys., **9** (1997), 397–410.
- [16] K. NAKANISHI, *Asymptotically-free solutions for the short-range nonlinear Schrödinger equation*, SIAM J. Math. Anal., **32** (2001), 1265–1271.
- [17] K. NAKANISHI AND T. OZAWA, *Remarks on scattering for nonlinear Schrödinger equations*, NoDEA Nonlinear Differential Equations Appl., **9** (2002), 45–68.

- [18] W. STRAUSS, *Nonlinear Wave Equations*, CBMS Regional Conference Series in Math., **73**, AMS, 1989.
- [19] H. TRIEBEL, *Theory of Function Spaces*, Monographs in Math., **78**, Birkhäuser, 1983.