

## PATTERNS IN A BALANCED BISTABLE EQUATION WITH HETEROGENEOUS ENVIRONMENTS ON SURFACES OF REVOLUTION

## MAICON SÔNEGO

Abstract. We use the variational concept of  $\Gamma$ -convergence to obtain sufficient conditions that guarantee existence, stability and the geometric structure of four families of stationary solutions to the singularly perturbed parabolic equation  $\partial_t u_\varepsilon = \varepsilon^2 \Delta u_\varepsilon + f(u_\varepsilon, x)$  on surfaces of revolution. We consider the bistable function f(u,x) = -(u-a(x))(u-b(x))(u-c(x)) and the conditions found relate the functions a,b,c to the geometry of the surface where such functions are defined.

*Mathematics subject classification* (2010): 35K57, 35B36, 35R01, 35B25, 35B35, 34K20, 58J32. *Keywords and phrases*: patterns, Γ-convergence, surface of revolution, stable transition layer.

## REFERENCES

- [1] C. BANDLE, F. PUNZO AND A. TESEI, Existence and nonexistence of patterns on Riemannian manifolds, J. Math. Anal. Appl. 387 (2012), 33–47.
- [2] A. L. BESSE, Manifolds all of whose Geodesics are Closed, Springer, 1978.
- [3] R. G. CASTEN AND C. J. HOLLAND, Instability results for reaction diffusion equations with Neumann boundary conditions, J. Diff. Eqns., 2 (1978), 266–273.
- [4] A. S. DO NASCIMENTO, Stable stationary solutions induced by spatial inhomogeneity via Γ-convergence, Bulletin of the Brazilian Mathematical Society, 29 No.1 (1998), 75–97.
- [5] A. S. DO NASCIMENTO, On the role of diffusivity in some stable equilibria of a diffusion equation, J. Diff. Eqns., 155 No. 2 (1999), 231–244.
- [6] A. S. DO NASCIMENTO AND M. SÔNEGO, Patterns on surfaces of revolution in a diffusion problem with variable diffusivity, Electr. J. Diff. Eqns., 2014, (2014) No. 238, 1–13.
- [7] A. S. DO NASCIMENTO AND M. SÔNEGO, The roles of diffusivity and curvature in patterns on surfaces of revolution, J. of Math. Anal. and Appl. Vol. 412 (2014), 1084–1096.
- [8] A. S. DO NASCIMENTO AND M. SÔNEGO, Stable equilibria to a singularly perturbed reactiondiffusion equation in a degenerated heterogeneous environment, J. of Math. Anal. and Appl. Vol. 433 (2016), 1743–1756.
- [9] A. S. DO NASCIMENTO AND M. SÔNEGO, Stable Transition Layers to Singularly Perturbed Spatially Inhomogeneous Allen-Cahn Equation, Advanced Nonlinear Studies 15 (2015), 363–376.
- [10] E. GIUSTI, Minimal Surfaces and Functions of Bounded Variation, Birkhauser, 1984.
- [11] D. HENRY, Geometric Theory of Semilinear Parabolic Equations, Springer Lecture Notes in Mathematics, 840, 1981.
- [12] R. V. KOHN AND P. STERNBERG, Local minimizers and singular pertubations, Proceedings of the R. Soc. of Edinburg, 111 (1989), 69–84.
- [13] K. KURATA, H. MATSUZAWA, Multiple stable patterns in a balanced bistable equation with heterogeneous environments, Applicable Analysis, 89 No. 7 (2010), 1023–1035.
- [14] F. LI, K. NAKASHIMA, Transition Layers for a spatially inhomogeneous Allen-Cahn equation in multi-dimensional domains, Discrete Contin. Dyn. Syst., 32 No. 4 (2012), 1391–1420.
- [15] F. MAHMOUDI, A. MALCHIODI AND J. WEI, Transition layer for the heterogeneous Allen-Cahn equation, Ann. I. H. Poincare A.N., 25 (2008), 609–631.
- [16] H. MATANO, Asymptotic behavior and stability of solutions of semilinear diffusion equations, Publ. RIMS, Kyoto Univ., 15 (1979), 401–454.



- [17] H. MATSUZAWA, Stable transition layers in a balanced bistable equation with degeneracy, Nonlinear Analysis, 58 (2004), 45–67.
- [18] K. NAKASHIMA, Stable transition layers in a balanced bistable equation, Differential Integral Equations, 13 No. 7-9 (2000), 1025–1038.
- [19] F. PUNZO, The existence of patterns on surfaces of revolution without boundary, Nonlinear Analysis, 77 (2012), 94–102.
- [20] J. SMOLLER, Shock Waves and Reaction Diffusion Equations, Springer New York, 258, 1994.
- [21] P. STERNBERG, The effect of a singular pertubation on nonconvex variational problems, Arch. Rat. Mech. Anal., 101 (1988), 209–260.