

WELL-POSEDNESS AND BLOW-UP FOR AN INHOMOGENEOUS SEMILINEAR PARABOLIC EQUATION

MOHAMED MAJDOUB

Abstract. We consider the large-time behavior of sign-changing solutions of the inhomogeneous equation $u_t - \Delta u = |x|^\alpha |u|^p + \zeta(t) \mathbf{w}(x)$ in $(0, \infty) \times \mathbb{R}^N$, where $N \geq 3$, $p > 1$, $\alpha > -2$, ζ, \mathbf{w} are continuous functions such that $\zeta(t) = t^\sigma$ or $\zeta(t) \sim t^\sigma$ as $t \rightarrow 0$, $\zeta(t) \sim t^m$ as $t \rightarrow \infty$. We obtain local existence for $\sigma > -1$. We also show the following:

- If $m \leq 0$, $p < \frac{N-2m+\alpha}{N-2m-2}$ and $\int_{\mathbb{R}^N} \mathbf{w}(x) dx > 0$, then all solutions blow up in finite time;
- If $m > 0$, $p > 1$ and $\int_{\mathbb{R}^N} \mathbf{w}(x) dx > 0$, then all solutions blow up in finite time;
- If $\zeta(t) = t^\sigma$ with $-1 < \sigma < 0$, then for $u_0 := u(t=0)$ and \mathbf{w} sufficiently small the solution exists globally.

We also discuss lower dimensions. The main novelty in this paper is that blow up depends on the behavior of ζ at infinity.

Mathematics subject classification (2010): 35K05, 35A01, 35B44.

Keywords and phrases: Inhomogeneous parabolic equation, global existence, finite time blow-up, differential inequalities, forcing term depending of time and space, critical Fujita exponent.

REFERENCES

- [1] J. AGUIRRE AND J. GIACOMONI, *The shape of blow-up for a degenerate parabolic equation*, Differ. Integral Equ., **14** (2001), 589–604.
- [2] B. BEN SLIMENE, S. TAYACHI AND F. B. WEISSLER, *Well-posedness, global existence and large time behavior for Hardy-Hénon parabolic equations*, Nonlinear Analysis, **152** (2017), 116–148.
- [3] C. BANDLE, H. A. LEVINE AND QI S. ZHANG, *Critical Exponents of Fujita Type for Inhomogeneous Parabolic Equations and Systems*, Journ. of Math. Anal. and App., **251** (2000), 624–648.
- [4] K. DENG AND H. A. LEVINE, *The role of critical exponents in blowup theorems, the sequel*, J. Math. Anal. Appl., **243** (2000), 85–126.
- [5] J. DIXON AND S. MCKEF, *Weakly singular discrete Gronwall inequalities*, Z. angew. Math. Mech., **64** (1986), 535–544.
- [6] M. J. ESTEBAN AND J. GIACOMONI, *Existence of global branches of positive solutions for semilinear elliptic degenerate problems*, J. Math. Pures Appl., **79** (2000), 715–740.
- [7] H. FUJITA, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo Sec. IA Math. **13** (1966), 109–124.
- [8] V. A. GALAKTIONOV AND J. L. VÁZQUEZ, *The problem of blow-up in nonlinear parabolic equations*, Discrete Contin. Dyn. Syst., **8** (2002), 399–433.
- [9] J. GIACOMONI, *Some results about blow-up and global existence to a semilinear degenerate heat equation*, Rev. Mat. Complut., **11** (1998), 325–351.
- [10] Y. HAN, *Blow-up phenomena for a reaction diffusion equation with special diffusion process*, Applicable Analysis (2020), doi:10.1080/00036811.2020.1792447.
- [11] K. HAYAKAWA, *On nonexistence of global solutions of some semilinear parabolic differential equations*, Proc. Japan Acad. **49** (1973), 503–505.
- [12] B. HU, *Blow Up Theories for Semilinear Parabolic Equations*, Springer, Berlin (2011).
- [13] M. JLELI, T. KAWAKAMI AND B. SAMET, *Critical behavior for a semilinear parabolic equation with forcing term depending of time and space*, J. Math. Anal. Appl. **486** (2020), 123931.

- [14] O. A. LADYZENSKAJA, V. A. SOLONNIKOV AND N. N. URALĆEVA, *Linear and quasilinear equations of parabolic type*, Amer. Math. Soc., Transl. Math. Monographs, Providence, R. I. (1968).
- [15] H. A. LEVINE, *The role of critical exponents in blowup theorems*, SIAM Rev., **32** (1990), 269–288.
- [16] H. A. LEVINE AND P. MEIER, *The value of the critical exponent for reaction-diffusion equations in cones*, Arch. Rational Mech. Anal., **109** (1989), 73–80.
- [17] A. V. MARTYNYENKO AND A. F. TEDEEV, *Cauchy problem for a quasilinear parabolic equation with a source term and an inhomogeneous density*, Comput. Math. Math. Phys., **47** (2007), 238–248.
- [18] A. V. MARTYNYENKO AND A. F. TEDEEV, *On the behavior of solutions to the Cauchy problem for a degenerate parabolic equation with inhomogeneous density and a source*, Comput. Math. Math. Phys., **48** (2008), 1145–1160.
- [19] E. MITIDIERI AND S. I. POHOZAEV, *A priori estimates and blow-up of solutions of nonlinear partial differential equations and inequalities*, Proc. Steklov Inst. Math., **234** (2001), 3–383.
- [20] C. A. STUART, *A critically degenerate elliptic Dirichlet problem, spectral theory and bifurcation*, Nonlinear Analysis, **190** (2020), Article ID 111620.
- [21] Z. TAN, *Reaction-diffusion equations with special diffusion processes*, Chin. J. Contemp. Math., **22** (2001), 371–382; translation from Chin. Ann. Math., Ser. A, **22** (2001), 597–607.
- [22] G. D. YOMGNE AND M. MAJDOUB, *Blow up and global existence of solutions to a Hardy-Hénon equation with a spatial-temporal forcing term*, submitted.
- [23] YUAN-WEI QI, *The critical exponents of parabolic equations and blow-up in \mathbb{R}^n* , Proceedings of the Royal Society of Edinburgh, **128A** (1998), 123–136.
- [24] P. QUITTNER AND P. SOUPLET, *Superlinear parabolic problems*, Birkhäuser Verlag, Basel (2007), xii+584.
- [25] S. TAYACHI, *Uniqueness and non-uniqueness of solutions for critical Hardy-Hénon parabolic equations*, J. Math. Anal. Appl., **488** (2020), 123976.
- [26] Q. S. ZHANG, *A new critical phenomenon for semilinear parabolic problem*, J. Math. Anal. Appl., **219** (1998), 123–139.
- [27] Q. S. ZHANG, *Blow up and global existence of solutions to an inhomogeneous parabolic system*, J. Differential Equations, **147** (1998), 155–183.
- [28] X. WANG, *On the Cauchy problem for reaction-diffusion equations*, Transactions of the American Mathematical Society, 337 (1993), 549–590.