THE SOLUTION OF FRACTIONAL NONLINEAR GINZBURG-LANDAU EQUATION WITH WEAK INITIAL DATA

JIE XIN, JIAQIAN HU AND HONG LU

Abstract. In this paper, we study the solution of the fractional nonlinear Ginzburg-Landau(FNGL) equation with weak initial data in the weighted Lebesgue spaces. The existence of a solution to this equation is proved by the contraction-mapping principle.

Mathematics subject classification (2010): 26A33, 35R11, 35Q56.

Keywords and phrases: Fractional nonlinear Ginzburg-Landau equation, global solution, contractionmapping principle, weighted Lebesgue spaces.

REFERENCES

- H. WEITZNER AND G. M. ZASLAVSKY, Some applications of fractional equations, Commun. Nonlinear Sci. Numer Simul 8 (2003), 273–281.
- [2] VASILY E. TARASOV AND GEORGE M. ZASLAVSKY, Fractional Ginzburg-Landau equation for fractal media, Physica A 354 (2005), 247–261.
- [3] A. V. MILOVANOV AND J. TUUL RASMUSSEN, Fractional generalization of the Ginzburg-Landau equation: an unconventional approach to critical phenomena in complex media, Physics Letters A 337 (2005), 75–80.
- [4] J.-H. WU, Quasi-geostrophic type equations with weak initial data, Electon. J. Differential Equations (1998), 1–10.
- [5] CARLOS E. KENIG, GUSTAVO PONCE AND LUIS VEGA, Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle, Comm. Pure Appl. Math. XLVI (1993), 527–620.
- [6] T. KATO AND G. PONCE, Commutator estimates and the Euler and Navier-Stokes equations, Comm. Pure Appl. Math. XXI (1998), 891–907.
- [7] R. R. COIFMAN AND Y. MEYER, Nonlinear harmonic analysis, operator theory and P.D.E., in "Beijing Lectures in Harmonic Analysis", Princeton University Press (1986), 3–45.
- [8] T. KATO AND G. PONCE, The Navier-Stokes equations with weak initial data, Int Math. Research Notices 10 (1994), 435–444.
- [9] Y. GIGA, T. MIYAKAWA AND H. OSADA, Two dimensional Navier-Stokes flow with measures as initial vorticity, Arch. Rotional Mech. Anal. 104 (1998), 223–250.

