

A NEW APPLICATION OF CONVEX SEQUENCES

HÜSEYİN BOR

Abstract. In this paper, an application of convex sequences dealing with $|C, \alpha|$ summability has been generalized to the $|C, \alpha, \beta; \delta|_k$ summability. Some new results have also been obtained.

Mathematics subject classification (2010): 40D15, 40F05, 40G05, 40G99.

Keywords and phrases: Cesàro mean, infinite series, summability factors, convex sequences.

REFERENCES

- [1] H. BOR, *On a new application of power increasing sequences*, Proc. Estonian Acad. Sci. Phys. Math. **57** (2008), 205–209.
- [2] H. BOR, *On the generalized absolute Cesàro summability*, Pac. J. Appl. Math. **2** (2009), 35–40.
- [3] D. BORWEIN, *Theorems on some methods of summability*, Quart. J. Math., Oxford Ser. **9** (1958), 310–316.
- [4] H. C. CHOW, *On the summability factors of Fourier Series*, J. London Math. Soc. **16** (1941), 215–220.
- [5] G. DAS, *A Tauberian theorem for absolute summability*, Proc. Camb. Phil. Soc. **67** (1970), 321–326.
- [6] T. M. FLETT, *On an extension of absolute summability and some theorems of Littlewood and Paley*, Proc. London Math. Soc. **7** (1957), 113–141.
- [7] T. M. FLETT, *Some more theorems concerning the absolute summability of Fourier series and power series*, Proc. London Math. Soc. **7** (1958), 357–387.
- [8] T. PATI, *The summability factors of infinite series*, Duke Math. J. **21** (1954), 271–284.