SOME PROPERTIES OF p–VALENT ANALYTIC FUNCTIONS INVOLVING CHO–KWON–SRIVASTAVA INTEGRAL OPERATOR

K. I. Noor, S. Z. H. Bukhari, M. Arif and M. Nazir

Abstract. In this paper, we introduce a subclass of p-valent functions involving Cho-Kwon-Srivastava integral operator. We study some interesting results including inclusion relations, convoltion with convex functions and integral preserving property for this class.

REFERENCES

[1] B. C. Carlson, B. D. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal., **15** (4) (1984), 737–745.

[2] N. E. Cho, O. S. Kwon, H. M. Srivastava, *Inclusion relationships and argument properties for certain subclasses of multivalent functions associated with a family of linear operators*, J. Math. Anal. Appl., **292** (2004), 470–483.

[3] J. H. Choi, M. Saigo, H. M. Srivastava, *Some inclusion properties of a certain family of integral operators*, J. Math. Anal. Appl. **276** (2002), 432–445.

[4] P. Eenigenburg, S. S. Miller, P. T. Mocanu, M. O. Reade, *On a Briot–Bouquet differential subordination*, Gen. Math. 3, Int. Series of Num. Math., Birkhäuser Verlag, Basel, 64 (1983), 339–348; see also Rev. Roumaine Math. Pures Appl., **29** (1984), 567–573.

[5] H. Irmak, R. K. Raina, *The starlikeness and convexity of multivalent functions involving certain inequalities*, Rev. Mat. Complut., **16** (2) (2003), 391–39.

[6] Y. C. Kim, J. H. Choi, T. Sugawa, *Coefficient bounds and convolution properties for certain classes of close-to-convex functions*, Proc. Japan Acad. Ser. A Math. Sci. **76** (2000), 95–98.

[7] J.-L. Liu, *The Noor integral and strongly starlike functions*, J. Math. Anal. Appl,. **261** (2001), 441–447.

[8] W. Ma, D. Minda, *An internal geometric characterization of strongly starlike functions*, Ann. Univ. Mariae Curie-Sklodowska Sect. A, **45** (1991), 89–97.

[9] S. S. Miller, P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Pure Appl. Math., Vol. 225, Dekker, New York, 2000.

[10] K. I. Noor, *On new classes of integral operators*, J. Natur. Geom., **16** (1999), 71–80.

[11] K. I. Noor, *On classes of analytic functions defined by convolution with incomplete beta functions*, J. Math. Anal. Appl., **307** (2005), 339–349.

[12] K. I. Noor, *On some applications of certain integral operators*, Appl. Math.Comput., **188** (2007), 814–823.

[13] K. I. Noor, S. Mustafa, B. Malik, *On some classes of p-valent functions involving certain Carlson Shaffer operator*, Appl. Math. Comput, **214** (2009), 336–341.

[14] K. I. Noor, M. A. Noor, *On integral operators*, J. Math. Anal. Appl., **238** (1999), 341–352.

[15] S. Owa, *On the distortion theorems I*, Kyungpook Math. J., **18** (1) (1978), 53–59.

[16] S. Owa, H. M. Srivastava, *Univalent and starlike generalized hypergeometric functions*, Canad. J. Math., **39** (5) (1987), 1057–1077.

[17] M. S. Robertson, *Multivalently starlike functions*, Duke Math. J., **20** (1953), 539–549.

[18] St. Ruscheweyh, *Convolutions in Geometric Function Theory*, Les Presses de l'Univ. de Montreal, 1982.

[19] H. Saitoh, *A linear operator and its applications of first order differential subordinations*, Math. Japon., **44** (1996), 31–38.

[20] J. SOKÓŁ, L. T. SPELINA, *Convolution properties for certain classes of multivalent functions*, J. Math. Anal. Appl. **337** (2008), 1190–1197.

[21] H. M. SRIVASTAVA, S. OWA, *An application of the fractional derivative*, Math. Japon., **29** (3) (1984), 383–389.

[22] T. YAGUCHI, *The radii of starlikeness and convexity for certain multivalent functions*, in: H. M. Srivastava, S. Owa (Eds.), Current Topics in Analytic Function Theory, World Scientific, Singapore, 1992, 375–386.

Journal of Classical Analysis
www.ele-math.com
jca@ele-math.com