SOME RESULTS ON POROUS SET RELATING TO RATIO SETS

D. K. GANGULY AND DHANANJOY HALDER

Abstract. An attempt has been made in this paper is to show that every Lebesgue measurable linear set with positive measure has a porous subset whose ratio set contains an interval. The category analogue of this result is also established.

Mathematics subject classification (2010): 28A05, 26A15. *Keywords and phrases*: Ratio set, property of Baire, porous set, difference set.

REFERENCES

- N. C. BOSE MAJUMDAR, On some properties of sets with positive measure, Annali Dell' Univ. di Ferrara (N. S.) Sez VII, Sci. Mat., X, 1 (1961), 1–12.
- [2] ZOLTAN BUCZOLICH, Every set of positive measure has a porous subset with difference set containing an interval, Real Analys Exchange 14 (1988–89).
- [3] H. I. MILLER AND LEJLA MILLER, A result about porous sets and difference sets, Real Analysis Exchange 13 (1987–88).
- [4] I. P. NATANSON, Theory functions of a real variable, Frederick Ungar, NY, 1955.
- [5] J. C. OXTOBY, *Measure and Category*, Second Edition, Springer-Verlag, NY, Heidelberg Berlin, 1980.
- [6] W. R. UTZ, The distance set of the Cantor discontinuum, Amer. Math. Monthly, June (1951), 407– 408.

