SHARING THE ZEROS OF POLYNOMIALS WITH REVERSE INDEX FOUR IN WEIGHTED WIDER SENSE

ABHIJIT BANERJEE* AND JHILIK BANERJEE

Abstract. In this paper, on the basis of zero set of reverse indexed polynomial, first introduced by us, we have investigated the uniqueness of meromorphic functions under weighted sharing in wider sense criteria [4], which in turn extend some earlier results in different directions. We have succeeded to identify a subclass of meromorphic functions for which uniqueness property exists for higher reverse indexed polynomial in literature. In the last section, we have presented the application of our results in case of derivatives of the concerned functions accompanied by series of examples.

Mathematics subject classification (2020): 30D35.

Keywords and phrases: Meromorphic functions, polynomials, weighted sharing, shared sets, uniqueness.

REFERENCES

- [1] A. BANERJEE, On a question of Gross, J. Math. Anal. Appl., 327, 2 (2007), 1273–1283.
- [2] A. BANERJEE, On the uniqueness of meromorphic functions that share two sets, Georgian Math. J., 15, 1 (2008), 21–38.
- [3] A. BANERJEE, Uniqueness of meromorphic functions sharing two sets with finite weight II, Tamkang J. Math., 41, 4 (2010), 379–392.
- [4] A. BANERJEE AND J. BANERJEE, On strong uniqueness polynomials in the wider sense and associated unique range sets of the same category, Adv. Stud: Euro-Tbilisi Math. J., 16, 4 (2023), 175–189.
- [5] A. BANERJEE AND S. MALLICK, On the characterisations of a new class of strong uniqueness polynomials generating unique range sets, Comput. Methods Funct. Theo., 17, 1 (2017), 19–45.
- [6] A. BANERJEE AND S. MALLICK, Uniqueness properties of meromorphic functions in the light of three shared sets, Math. Slovaca, 68, 3 (2018), 591–606.
- [7] J. CHANG, M. FANG AND L. ZALCMAN, Entire functions that share a set with their derivatives, Archiv der Mathematik, 89, (2007), 561–569.
- [8] J. CHANG AND L. ZALCMAN, Meromorphic functions that share a set with their derivatives, J. Math. Anal. Appl., 338, (2008), 1020–1028.
- [9] M. FANG AND L. ZALCMAN, Normal families and uniqueness theorems for entire functions, J. Math. Anal. Appl., 280, (2003), 273–283.
- [10] F. GROSS, Factorization of meromorphic functions and some open problems. Proc. Conf. Univ. Kentucky, Leixngton, Ky (1976); Lecture Notes in Math, 599, (1977), 51–69, Springer (Berlin).
- [11] W. K. HAYMAN, Meromorphic Functions, The Clarendon Press, Oxford (1964).
- [12] I. LAHIRI, Weighted sharing and uniqueness of meromorphic functions, Nagoya Math. J., 161, (2001), 193–206.
- [13] I. LAHIRI, Value distribution of certain differential polynomials, Int. J. Math. Math. Sci., 28, 2 (2001), 83–91.
- [14] I. LAHIRI, Weighted value sharing and uniqueness of meromorphic functions, Complex Var. Theo. Appl., 46, 3 (2001), 241-253.
- [15] W. C. LIN AND H. X. YI, Some further results on meromorphic functions that share two sets, Kyungpook Math. J., 43, 1 (2003), 73–85.



- [16] F. Lü, A note on meromorphic functions that share a set with their derivatives, Arch. Math., 96, (2011), 369–377.
- [17] A. Z. MOKHON'KO, On the Nevanlinna characteristics of some meromoephic function, in "Theory of functions, functional analysis and their applications", Izd-vo Khar'kovsk, Un-ta, 14, (1971), 83–87.
- [18] E. MUES AND N. STEINMETZ, Meromorphe Funktionen die unit ihrer Ableitung Werteteilen, Manuscripta Math., 29, (1979), 195–206.
- [19] H. X. YI, Meromorphic functions that share one or two values II, Kodai Math. J., 22, 2 (1999), 264– 272.
- [20] H. X. YI, Meromorphic functions that share two sets (Chinese), Acta Math. Sinica (Chin. Ser.), 45, 1 (2002), 75–82.
- [21] H. X. YI AND W. R. LÜ, Meromorphic functions that share two sets II, Acta Math. Sci. (Ser. B) Engl. Ed., 24, 1 (2004), 83–90.