

CERTAIN CLASSES OF ANALYTIC FUNCTIONS WITH NEGATIVE COEFFICIENTS ASSOCIATED WITH A CONVOLUTION STRUCTURE

G. MURUGUSUNDARAMOORTHY AND S. B. JOSHI

Abstract. Making use of a convolution structure, we introduce a new class of analytic functions $\mathbb{P}\mathbb{T}_g(\lambda, \alpha, \beta, \gamma)$ defined in the open unit disc and investigate its various characteristics. Further we obtained distortion bounds, extreme points and radii of close-to-convexity, starlikeness and convexity for functions belonging to the class $\mathbb{P}\mathbb{T}_g(\lambda, \alpha, \beta, \gamma)$.

Mathematics subject classification (2000): 30C45.

Keywords and phrases: Analytic, univalent, starlikeness, convexity, Hadamard product (convolution).

REFERENCES

- [1] O. ALTINTAS, *A subclass of analytic functions with negative coefficients*, Bull. Sci-Engg. Hacettepe Univ., **19** (1990), 15–24.
- [2] G. J. BAO AND Y. LING, *On Some classes of analytic functions with negative coefficients*, J. Harbin Inst. Tech., **23** (1991), 100–103.
- [3] B. C. CARLSON AND D. B. SHAFFER, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal., **15** (1984), 737–745.
- [4] N. E. CHO AND T. H. KIM, *Multiplier transformations and strongly close-to-convex functions*, Bull. Korean Math. Soc., **40**, 3 (2003), 399–410.
- [5] N. E. CHO AND H. M. SRIVASTAVA, *Argument estimates of certain analytic functions defined by a class of multiplier transformations*, Math. Comput. Modelling, **37**, 1–2 (2003), 39–49.
- [6] J. DZIOK AND H. M. SRIVASTAVA, *Classes of analytic functions associated with the generalized hypergeometric function*, Appl. Math. Comput., **103**, 1 (1999), 1–13.
- [7] V. P. GUPTA AND P. K. JAIN, *Certain classes of univalent functions with negative coefficients-II*, Bull. Austral. Math. Soc., **15** (1976), 467–473.
- [8] S. OWA AND S. K. LEE, *Certain generalized class of analytic functions with negative coefficients*, Bull. Cal. Math. Soc., **82** (1990), 284–289.
- [9] J. PATEL AND A. K. MOHANTY, *On certain generalized class of analytic functions*, Soochow J. Math., **23**, 4 (1997), 365–379.
- [10] G. Ş. SĂLĂGEAN, *Subclasses of univalent functions*, Lect. Notes in Math. (Springer-Verlag), **1013** (1983), 362–372.
- [11] S. M. SARANGI AND B. A. URALEGADDI, *The radius of convexity and starlikeness for certain classes of analytic functions with negative coefficients*, Atti. Acad. Naz. Lincei Rend. Sc. Fis. Mat. Natur., **45** (1978), 38–42.
- [12] H. SILVERMAN, *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc., **51** (1975), 109–116.
- [13] S. RUSCHEWEYH, *New criteria for univalent functions*, Proc. Amer. Math. Soc., **49** (1975), 109–115.
- [14] T. YAGUCHI, T. SEKINE, H. SAITOH, S. OWA, M. NUNOKAWA AND S. FUKUI, *A generalization class of certain subclasses of analytic functions with negative coefficients*, Proc. Inst. Natur. Sci. Nihon Univ., **25** (1990), 67–80.