

## THE GOLDEN–THOMPSON–SEGAL TYPE INEQUALITIES RELATED TO THE WEIGHTED GEOMETRIC MEAN DUE TO LAWSON–LIM

JUN ICHI FUJII, MASATOSHI FUJII AND YUKI SEO

**Abstract.** In this paper, by using the weighted geometric mean  $G[n, t]$  and the weighted arithmetic one  $A[n, t]$  due to Lawson-Lim for each  $t \in [0, 1]$ , we investigate  $n$ -variable versions of a complement of the Golden-Thompson-Segal type inequality due to Ando-Hiai: Let  $H_1, H_2, \dots, H_n$  be selfadjoint operators such that  $m \leq H_i \leq M$  for  $i = 1, 2, \dots, n$  and some scalars  $m \leq M$ . Then

$$S(e^{p(M-m)})^{-\frac{2}{p}} \| G[n, t](e^{pH_1}, \dots, e^{pH_n})^{\frac{1}{p}} \| \\ \leq \| e^{A[n, t](H_1, \dots, H_n)} \| \leq S(e^{p(M-m)})^{\frac{2}{p}} \| G[n, t](e^{pH_1}, \dots, e^{pH_n})^{\frac{1}{p}} \|$$

for all  $p > 0$  and the both-hand sides of the inequality above converge to the middle-hand side as  $p \downarrow 0$ , where  $S(\cdot)$  is the Specht ratio and  $\|\cdot\|$  stands for the operator norm.

**Mathematics subject classification (2000):** 47A30, 47A63 and 47A64.

**Keywords and phrases:** Positive operator, Specht ratio, Golden-Thompson-Segal inequality, geometric mean, Lie-Trotter formula.

### REFERENCES

- [1] M. ALIĆ, P. S. BULLEN, J. E. PEČARIĆ AND V. VOLENEC, *On the geometric-arithmetic mean inequality for matrices*, Math. Communications, **2** (1997), 125–128.
- [2] T. ANDO AND F. HIAI, *Log-majorization and complementary Golden-Thompson type inequalities*, Linear Alg. Appl., **197/198** (1994), 113–131.
- [3] T. ANDO, C.-K. LI AND R. MATHIAS, *Geometric means*, Linear Alg. Appl., **385** (2004), 305–334.
- [4] J. I. FUJII, M. FUJII, M. NAKAMURA, J. E. PEČARIĆ AND Y. SEO, *A reverse of the weighted geometric mean due to Lawson-Lim*, Linear Alg. Appl., **427** (2007), 272–284.
- [5] M. FUJII AND R. NAKAMOTO, *A geometric mean in the Furuta inequality*, Sci. Math. Japon., **55** (2002), 615–621.
- [6] M. FUJII, Y. SEO AND M. TOMINAGA, *Golden-Thompson type inequalities related to a geometric mean via Specht's ratio*, Math. Inequal. Appl., **5** (2002), 573–582.
- [7] T. FURUTA, J. MIĆIĆ, J. E. PEČARIĆ AND Y. SEO, *Mond-Pečarić Method in Operator Inequalities*, Monographs in Inequalities 1, Element, Zagreb, 2005.
- [8] S. GOLDEN, *Lower bounds for Helmholtz function*, Phys. Rev., **137** (1965), B1127–B1128.
- [9] F. HIAI AND D. PETZ, *The Golden-Thompson trace inequality is complemented*, Linear Algebra Appl., **181** (1993), 153–185.
- [10] F. KUBO AND T. ANDO, *Means of positive linear operators*, Math. Ann., **246** (1980), 205–224.
- [11] J. LAWSON AND Y. LIM, *A general framework for extending means to higher orders*, preprint. ([http://arxiv.org/PS\\_cache/math/pdf/0612/0612293v1.pdf](http://arxiv.org/PS_cache/math/pdf/0612/0612293v1.pdf)).
- [12] J. LAWSON AND Y. LIM, *Higher order weighted matrix means and related matrix inequalities*, preprint.
- [13] I. SEGAL, *Notes toward the construction of nonlinear relativistic quantum fields, III*, Bull. Amer. Math. Soc., **75** (1969), 1390–1395.
- [14] Y. SEO, *Reverses of the Golden-Thompson type inequalities due to Ando-Hiai-Petz*, Banach J. Math. Anal., **2** (2008), 140–149.

- [15] W. SPECHT, *Zur Theorie der elementaren Mittel*, Math. Z., **74** (1960), 91–98.
- [16] K. SYMANZIK, *Proof and refinements of an inequality of Feynman*, J.Math. Phys., **6** (1965), 1155–1156.
- [17] C. J. THOMPSON, *Inequality with applications in statistical mechanics*, J. Math. Phys., **6** (1965), 469–480.
- [18] M. TOMINAGA, *Specht's ratio in the Young inequality*, Sci. Math. Japon., **55** (2002), 585–588.
- [19] A. M. TURING, *Rounding off-errors in matrix processes*, Quart. J. Mech. Appl. Math., **1** (1948), 287–308.
- [20] T. YAMAZAKI AND M. YANAGIDA, *Characterizations of chaotic order associated with Kantorovich inequality*, Sci. Math., **2** (1999), 37–50.