

SHARP BOUNDS FOR SEIFFERT MEANS IN TERMS OF LEHMER MEANS

MIAO-KUN WANG, YE-FANG QIU AND YU-MING CHU

Abstract. In this paper, we establish two sharp inequalities as follows: $P(a,b) > L_{-\frac{1}{6}}(a,b)$ and $T(a,b) < L_{\frac{1}{3}}(a,b)$ for all $a, b > 0$ with $a \neq b$. Here, $L_r(a,b)$, $P(a,b)$ and $T(a,b)$ are the Lehmer, first and second Seiffert means of a and b , respectively.

Mathematics subject classification (2010): 26E60.

Keywords and phrases: Lehmer mean, first Seiffert mean, second Seiffert mean.

REFERENCES

- [1] D. H. LEHMER, *On the compounding of certain means*, J. Math. Anal. Appl., **36** (1971), 183–200.
- [2] H. ALZER, *Über Lehmers Mittelwertfamilie*, Elem. Math., **43**, 2 (1988), 50–54.
- [3] ZH. LIU, *Remark on inequalities between Hölder and Lehmer means*, J. Math. Anal. Appl., **247**, 1 (2000), 309–313.
- [4] H. ALZER, *Bestmögliche Abschätzungen für spezielle Mittelwerte*, Zb. Rad. Prirod. -Mat. Fak. Ser. Mat., **23**, 1 (1993), 331–346.
- [5] K. B. STOLARSKY, *Hölder means, Lehmer means, and $x^{-1} \log \cosh x$* , J. Math. Anal. Appl., **202**, 3 (1996), 810–818.
- [6] H. J. SEIFFERT, *Problem 887*, Nieuw Arch. Wisk. (4), **11** (1993), 176–176.
- [7] H. J. SEIFFERT, *Aufgabe β16*, Die Wurzel, **29** (1995), 221–222.
- [8] P. A. HÄSTÖ, *Optimal inequalities between Seiffert's mean and power mean*, Math. Inequal. Appl., **7**, 1 (2004), 47–53.
- [9] E. NEUMAN, J. SÁNDOR, *On certain means of two arguments and their extensions*, Int. J. Math. Math. Sci., **16** (2003), 981–983.
- [10] E. NEUMAN, J. SÁNDOR, *On the Schwab-Borchardt mean*, Math. Pannon., **14**, 2 (2003), 253–266.
- [11] P. A. HÄSTÖ, *A monotonicity property of ratios of symmetric homogeneous means*, JIPAM. J. Inequal. Pure Appl. Math., **3**, 5 (2002), Article 71, 23 pp.
- [12] P. S. BULLEN, D. S. MITRINOVÍĆ AND P. M. VASIĆ, *Means and their inequalities*, D. Reidel Publishing Co., Dordrecht, 1988.