THE WEIGHTED ESTIMATE FOR THE COMMUTATOR
OF THE GENERALIZED FRACTIONAL INTEGRAL

YUE HU, YANHUI WANG AND YUESHAN WANG

Abstract. Let L be the infinitesimal generator of an analytic semigroup on $L^2(\mathbb{R}^n)$ with Gaussian kernel bound, and let $L^{-\alpha/2}$ be the fractional integral of L for $0 < \alpha < n$. Suppose that b is a locally integral function, then the commutator generated by b and $L^{-\alpha/2}$ is defined by $[b, L^{-\alpha/2}](f) = bL^{-\alpha/2}(f) - L^{-\alpha/2}(bf)$. When b belongs to weighted Lipschitz function space, the boundedness of $[b, L^{-\alpha/2}]$ from $L^p(\omega, \mathbb{R}^n)$ to $L^q(\omega^{1-(1-\alpha/n)q}, \mathbb{R}^n)$ is established, where $1 < p < \infty$, $0 < \beta < 1$ and $1/q = 1/p - (\alpha + \beta)/n$ with $1/p > (\alpha + \beta)/n$.

Mathematics subject classification (2010): 42B25, 42B20, 42B35.
Keywords and phrases: Gaussian upper bound, commutator, fractional integral, Muckenhoupt weight, Lipschitz function.

REFERENCES

[1] X. T. DUONG AND L. X. YAN, *On commutators of fractional integrals*, Proc. Amer. Math. Soc. **132** (2004), 3549–3557.

[2] D. G. DENG, X. T. DUONG, A. SIKORA AND L. X. YAN, *Comparison of the Classical BMO with the BMO spaces associated with operators and applications*, Rev. Mat. Iberoamericana **24** (2008), 267–296.

[3] S. CHANILLO, *A note on commutator*, Indiana Univ. Math. J. **31** (1982), 7–16.

[4] M. PALUSZYŃSKI, *Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss*, Indiana Univ. Math. J. **44** (1995), 1–17.

[5] B. HU AND J. GU, *Necessary and sufficient conditions for boundedness of some commutators with weighted Lipschitz functions*, J. Math. Anal. Appl. **340** (2008), 598–605.

[6] H. X. MO AND S. Z. LU, *Boundedness of multilinear commutators of generalized fractional integrals*, Math. Nachr. **281** (2008), 1328–1340.

[7] P. AUSCHER AND J. M. MARTELL, *Weighted norm inequalities for fractional operators*, Indiana Univ. Math. J. **57** (2008), 1845–1869.

[8] B. MUCKENHOUPT, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165** (1972), 207–226.

[9] B. MUCKENHOUPT, R. WHEEDEN, *Weighted Norm Inequalities for Fractional Integrals*, Trans. Amer. Math. Soc. **192** (1974), 261–274.

[10] J. GARCÍA-CUERVA, *Weighted H^p spaces*, Dissert. Math., 1979, 162.

[11] J. GARCÍA-CUERVA AND J. L. RUBIO DE FRANCIA, *Weighted Norm Inequalities and Related Topics*, North-Holland, Amsterdam, 1985.

[12] J. M. MARTELL, *Sharp maximal functions associated with approximations of the identity in spaces of homogeneous type and applications*, Studia Math. **161** (2004), 113–145.

© ELEMENT, Zagreb
Paper JMI-07-30

Journal of Mathematical Inequalities
www.ele-math.com
jmi@ele-math.com