

## ANOTHER APPROACH TO BECKNER'S INEQUALITY

RYOTARO TANAKA, KICHI-SUKE SAITO AND NAOTO KOMURO

*Abstract.* A classical inequality which was proved by Beckner is an important tool for the study of Banach space geometry. In this note, we present another proof of that inequality.

*Mathematics subject classification (2010):* 46B20.

*Keywords and phrases:* Beckner's inequality.

### REFERENCES

- [1] W. BECKNER, *Inequalities in Fourier analysis*, Ann. of Math., **102** (1975), 159–182.
- [2] K. KURIYAMA, M. MIYAGI, M. OKADA AND T. MIYOSHI, *Elementary proof of Clarkson's inequalities and their generalization*, Memories of the Faculty of Engineering, Yamaguchi Univ., **48** (1997), 119–125.
- [3] J. LINDENSTRAUSS AND L. TZAFRIRI, *Classical Banach spaces II*, Springer-Verlag, Berlin, 1979.
- [4] K.-I. MITANI AND K.-S. SAITO, *Dual of two dimensional Lorentz sequence spaces*, Nonlinear Anal., **71** (2009), 5238–5247.

- [5] K.-I. MITANI, K.-S. SAITO AND T. SUZUKI, *On the calculation of the James constant of Lorentz sequence spaces*, J. Math. Anal. Appl., **343** (2008), 310–314.
- [6] H. MIZUGUCHI AND K.-S. SAITO, *A note on Clarkson's inequality in the real case*, J. Math. Inequal., **4** (2010), 129–132.
- [7] Y. YAMADA, Y. TAKAHASHI AND M. KATO, *On Hanner type inequalities with a weight for Banach spaces*, J. Math. Anal. Appl., **324** (2006), 1228–1241.