

## A SOLUTION TO AN OPEN PROBLEM FOR WILKER–TYPE INEQUALITIES

CHEN-SHIPING\* AND GE-XINYU

*Abstract.* This paper is to solve the open problem for Wilker-Type inequality: what are the best possible for the constants  $c_1$  and  $c_2$  such that the double inequality  $c_1 x^{3a} \tan x < \left(\frac{\sin x}{x}\right)^{2a} + \left(\frac{\tan x}{x}\right)^a - 2 < c_2 x^{3a} \tan x$  holds?

*Mathematics subject classification (2010):* 26D05, 33B10.

*Keywords and phrases:* Wilker-type inequality, trigonometric function, open problem.

### REFERENCES

- [1] J. B. WILKER, *Problem E3306*, The American Mathematical Monthly **96**, 1 (1989), 55.
- [2] J. S. SUMNER, A. A. JAGERS, M. VOWE AND J. ANGLÉSIO, *Inequalities involving trigonometric functions*, The American Mathematical Monthly **98**, 3 (1991), 264–267.
- [3] LING ZHU, *Some New Wilker-Type Inequalities for Circular and Hyperbolic Functions*, Abstract and Applied Analysis, 1 (2009), 9 pages, doi:10.1155/2009/485842.
- [4] EDWARD NEUMAN, *Inequalities Involving Generalized Trigonometric and Hyperbolic Functions*, Journal of Mathematical Inequalities **8**, 4 (2014), 19 pages, doi:10.7153/jmi-08-54.
- [5] LADISLAV MATEJČKA, *Note On Two New Wilker-Type Inequalities*, International Journal of Open Problems in Computer Science & Mathematics **4**, 1 (2011), 7 pages.
- [6] WEI-DONG JIANG, QIU-MING LUO AND FENG QI, *Refinements and Sharpening of some Huygens and Wilker Type Inequalities*, Turkish Journal of Analysis and Number Theory **2**, 4 (2014), 134–139, doi:10.12691/tjant-2-4-6.
- [7] KUANG JICHANG (Eds), *Applied Inequalities*, Shandong Science and Technology Press, 4th ed., Shandong, 2010.