

NONEXISTENCE OF GLOBAL SOLUTIONS OF A QUASILINEAR HYPERBOLIC EQUATION

VURAL BAYRAK, MEHMET CAN AND FAHREDDIN ALIYEV

Abstract. In this work, the nonexistence of the global solutions to a class of initial boundary value problems with dissipative terms in the boundary conditions is considered for a quasilinear hyperbolic equation. The nonexistence proof is achieved by the use of a lemma due to O. Ladyzhenskaya and V. K. Kalantarov and by the usage of the so called concavity method. In this method one writes down a functional which reflects the properties of dissipative boundary conditions and represents the norm of the solution in some sense, then proves that this functional satisfies the hypotheses of Ladyzhenskaya-Kalantarov lemma. Hence from the conclusion of the lemma one concludes that in finite time t_2 , this functional and hence the norm of the solution blows up.

Mathematics subject classification (1991): 35L70, 35B05, 65M06.

Key words and phrases: Quasilinear hyperbolic equations, nonexistence of global solutions, blow up of solutions.

REFERENCES

- [1] R. T. GLASSEY, *Blow-up theorems for nonlinear wave equations*, Math. Z. **132** (1973), 183–203.
- [2] O. A. LADYZHENSKAYA AND V. K. KALANTAROV, *Blow up theorems for quasilinear parabolic and hyperbolic equations*, Zap. Nauchn. SLOMI Steklov **69** (1977), 77–102.
- [3] H. A. LEVINE, *Instability and nonexistence of global solutions to nonlinear wave equations of the form $P u_{tt} = -A u + \mathcal{F}(u)$* , Trans. Am. Math. Soc. **192** (1974), 1–22.
- [4] J. L. LIONS, *Equations Différentielles Opérationnelles et Problème aux Limites*, Springer, 1961.