

DECOMPOSITION OF HOMOGENEOUS MEANS AND CONSTRUCTION OF SOME METRIC SPACES

P. KAHLIG AND J. MATKOWSKI

Abstract. Any (positively) homogeneous mean on $(0, \infty)^2$ can be decomposed multiplicatively into the arithmetic mean A and a one-place function, called A -index function. Index functions characterize a homogeneous mean in many respects, and their graphs are suitable for geometrical comparisons of several properties of homogeneous means. Moreover, index functions can facilitate proofs of inequalities between different types of homogeneous means. With the aid of A -index functions, some metrics are introduced in the set of homogeneous means.

Mathematics subject classification (1991): 26D07.

Key words and phrases: Homogeneous mean, index function of a mean, decomposition of a mean, complete metric space of homogeneous means.

REFERENCES

- [1] J. ACZÉL AND J. DHOMBRES, *Functional equations in several variables*, Encyclopaedia Math. Appl. Vol. 31, Cambridge University Press, New York – New Rochelle – Melbourne – Sydney, 1989.
- [2] P. S. BULLEN, D. S. MITRINOVIĆ AND P. M. VASIĆ, *Means and their inequalities*, D. Reidel Publ. Comp. Dordrecht, 1988.
- [3] T. P. LIN, *The power mean and the logarithmic mean*, Amer. Math. Monthly **81** (1974), 879–883.
- [4] J. MATKOWSKI, *A functional inequality characterizing convex functions, conjugacy and a generalization of Hölder's and Minkowski's inequalities*, Aequationes Math. **40** (1990), 168–180.
- [5] J. MATKOWSKI, *Subadditive functions and a relaxation of the homogeneity condition of seminorms*, Proc. Amer. Math. Soc. **117** (1993), 991–1001.
- [6] J. MATKOWSKI AND J. RÄTZ, *Convexity of power functions with respect to symmetric homogeneous means*, Internat. Ser. Numer. Math. **123** (1997), 231–247.
- [7] Zs. PÁLES, *Inequalities for sums of powers*, J. Math. Anal. Appl. **131** (1988), 265–270.