

MAXIMUM OF GREEN'S FUNCTIONS AND LIAPUNOV INEQUALITIES FOR THE STURM-LIOUVILLE PROBLEM

Laurie Battle

Abstract. First we find the Green's function for the Sturm-Liouville problem with separated boundary conditions. Then we calculate maximum values of the Green's function to determine Liapunov-type inequalities.

Mathematics subject classification (2000): 26D10, 34B27, 34B24, 26D20.

Key words and phrases: Green's function, Liapunov inequality, Sturm-Liouville problem, separated boundary conditions.

REFERENCES

- [1] R. C. BROWN AND D. B. HINTON, *Lyapunov inequalities and their applications*, Survey on Classical Inequalities, T. M. Rassias, ed., Kluwer, Dordrecht (2000), 1–25.
- [2] R. C. BROWN, A. M. FINK AND D. B. HINTON, *Some Opial, Lyapunov, and De la Valée Poussin Inequalities with Nonhomogeneous Boundary Conditions*, Journal of Inequalities and Applications, OPA **5** (2000), 11–37.
- [3] SUI-SANG CHENG, *Lyapunov inequalities for differential and difference equations*, Fasc. Math. **23** (1991), 25–41.
- [4] RANDAL H. COLE, *Theory of Ordinary Differential Equations*, Appleton-Century-Crofts (1968).