

AN APPLICATION OF ALMOST INCREASING SEQUENCES

HÜSEYİN BOR

Abstract. In this paper using any almost increasing sequence a result of Mishra and Srivastava [5] on $|C, 1|_k$ summability factors has been generalized for $|C, \alpha; \delta|_k$ summability factors under weaker conditions.

Mathematics subject classification (2000): 40D15, 40F05, 40G05.

Key words and phrases: Absolute Cesaro summability factors, almost increasing sequences.

REFERENCES

- [1] S. ALJANCIC AND D. ARANDELOVIC, *O*-regularly varying functions, Publ. Inst. Math., **22** (1977), 5–22.
- [2] L. S. BOSANQUET, *A mean value theorem*, J. London Math. Soc., **16** (1941), 146–148.
- [3] T. M. FLETT, *On an extension of absolute summability and some theorems of Littlewood and Paley*, Proc. London Math. Soc., **7** (1957), 113–141.
- [4] T. M. FLETT, *Some more theorems concerning the absolute summability of Fourier series*, Proc. London Math. Soc., **8** (1958), 357–387.
- [5] K. N. MISHRA AND R. S. L. SRIVASTAVA, *On absolute Cesàro summability factors of infinite series*, Portugaliae Math., **42** (1983–1984), 53–61.
- [6] T. PATI, *The summability factors of infinite series*, Duke Math. J. **21** (1954), 271–284.
- [7] I. SZALAY, *On generalized Cesaro summability factors*, Publ. Math. Debrecen, **24** (1977), 343–349.