

INEQUALITIES BETWEEN $f(\|A\|)$ AND $\|f(|A|)\|$

ROMAN DRNOVŠEK AND TOMAŽ KOSEM

Abstract. Let f be a nonnegative concave function on $[0, \infty)$, and let $\|\cdot\|$ be a unitarily invariant norm on the space of $n \times n$ complex matrices. We prove that, for any $n \times n$ complex matrix A , $f(\|A\|) \leq \|f(|A|)\|$ provided the norm $\|\cdot\|$ is normalized. On the other hand, if the norm of the identity matrix is 1, then $f(\|A\|) \geq \|f(|A|)\|$ for any matrix A . These results extend the theorems of F. Hiai and X. Zhan that were proved in the case when f is an operator monotone function.

Mathematics subject classification (2000): 15A60, 15A45.

Key words and phrases: unitarily invariant norms, inequalities, operator monotone functions.

REFERENCES

- [1] R. BHATIA, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [2] F. HANSEN, *An operator inequality*, *Math. Ann.*, **246** 3 (1979/80), 249–250.
- [3] F. HANSEN, *Operator monotone functions of several variables*, *Math. Inequal. Appl.*, **6** (2003), no. 1, 1–17.
- [4] F. HANSEN AND G. K. PEDERSEN, *Jensen's inequality for operators and Löwner's theorem*, *Math. Ann.* **258** 3 (1981/82), 229–241.
- [5] F. HIAI AND X. ZHAN, *Inequalities involving unitarily invariant norms and operator monotone functions*, *Lin. Alg. Appl.*, **341** (2002), 151–169.
- [6] X. ZHAN, *Matrix inequalities*, Springer-Verlag, Berlin, 2002.