

ON EXTENDED SINGULAR SET OF POTENTIALS

DARKO ŽUBRINIĆ

Abstract. We describe a class of potentials $v = G * f$, such that if x_0 is from extended singular set of v , that is, $r^{-N} \int_{B_r(x_0)} v(x) dx \rightarrow +\infty$ for some sequence $r \rightarrow 0$, then necessarily $v(x_0) = \infty$. This class includes Bessel potentials and Riesz potentials. The result was exploited in our previous paper in order to show that singular dimension of the Bessel potential space $L^{\alpha,p}(\mathbf{R}^N)$ (that is, the supremum of Hausdorff's dimension of extended singular sets, taken over all functions from the space) is equal to $N - \alpha p$, provided $\alpha p \leq N$.

Mathematics subject classification (2000): 26A12, 31C40, 46E35.

Key words and phrases: Singular set, potentials.

REFERENCES

- [1] ADAMS, D. R., HEDBERG, L. I., *Function Spaces and Potential Theory*, Springer Verlag, 1996.
- [2] FALCONER, K. J., *Fractal geometry*, John Wiley and Sons, New York, 1990.
- [3] RESHETNYAK, YU. G., *On the concept of capacity in the theory of functions with generalized derivatives*, Sibirskij mat. žurnal, **X**, 5, (1969), 1108–1138 (Russian); Siberian Math. J. **13** (1969), 818–842.
- [4] ZIEMER, W. P., *Weakly differentiable functions; Sobolev spaces and functions of bounded variation*, Graduate texts in mathematics, Springer–Verlag, 1989.
- [5] ŽUBRINIĆ, D., *Singular sets of Sobolev functions*, C. R. Acad. Sci. Paris, Ser. I, **334**, (2002), 539–544.
- [6] ŽUBRINIĆ, D., *Singular sets of Lebesgue integrable functions*, Chaos, Solitons and Fractals, **21**, (2004), 1281–1287.