

RICCI CURVATURE OF LAGRANGIAN SUBMANIFOLDS IN COMPLEX SPACE FORMS

TEODOR OPREA

Abstract. As we showed in [5] and [6], the basic inequalities, involving Riemannian invariants of a Lagrangian submanifold immersed in a complex space form, can be improved using optimization methods. Also in [1] is showed that the improved Chen's inequality from [5] is optimal. In this paper we find another proof for a Chen's inequality, regarding the Ricci curvature [2] and we improve this inequality in the Lagrangian case.

Mathematics subject classification (2010): 53C21, 53C24, 53C25, 49K35.

Keywords and phrases: Constrained maximum, Chen's inequality, Lagrangian submanifolds.

REFERENCES

- [1] J. BOLTON, L. VRANCKEN, *Lagrangian submanifolds attaining equality in the improved Chen's inequality*, Bull. Belg. Math. Soc., **14**, 2 (2007), 311–315.
- [2] B.Y. CHEN, *Mean curvature shape operator of isometric immersions in real-space-forms*, Glasgow Math. J., **38** (1996), 87–97.
- [3] B.Y. CHEN, *Relations between Ricci curvature shape operator for submanifolds with arbitrary codimensions*, Glasgow Math. J., **41** (1999), 33–41.
- [4] T. OPREA, *Optimizations on Riemannian Submanifolds*, Analele Univ. Buc., **LIV**, 1 (2005), 127–136.
- [5] T. OPREA, *Chen's inequality in the Lagrangian case*, Colloq. Math., **108**, 1 (2007), 163–169.
- [6] T. OPREA, *On a Riemannian invariant of Chen type*, Rocky Mountain J. Math., **38**, 2 (2008), 567–581.
- [7] C. UDRIȘTE, *Convex functions and optimization methods on Riemannian manifolds*, Kluwer Academic Publishers Group, Dordrecht, 1994.
- [8] C. UDRIȘTE, O. DOGARU, I. ȚEVY, *Extrema with nonholonomic constraints*, Geometry Balkan Press, Bucharest, 2002.