

INTERPOLATIONS OF SCHWAB–BORCHARDT MEAN

ALFRED WITKOWSKI

Abstract. For two means M, N satisfying $M(x,y) \leq N(x,y)$ we apply the 'borchardtisation' process to obtain a new mean

$$SB_{M,N} = \frac{\sqrt{N^2 - M^2}}{\arccos(M/N)}.$$

We use some geometric ideas to prove inequalities between the three means. In particular some new inequalities for Seiffert means are established.

Mathematics subject classification (2010): 26D15.

Keywords and phrases: Seiffert mean, Schwab-Borchardt mean, power means.

REFERENCES

- [1] B. C. CARLSSON, *Algorithms involving arithmetic and geometric means*, Amer. Math. Monthly **78** (1971), 496–505.
- [2] Y.-M. CHU, Y.-F. QIU, M.-K. WANG, AND G.-D. WANG, *The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean*, J. Inequal. Appl., Volume **2010** (2010), Article ID 436457, 7 pages, doi:10.1155/2010/436457.
- [3] P. A. HÄSTÖ, *A Monotonicity Property of Ratios of Symmetric Homogeneous Means*, J. Inequal. Pure and Appl. Math. **3**, 5 (2002), Article 71.
(Online: <http://www.emis.de/journals/JIPAM/article223.html?sid=223>).
- [4] P. A. HÄSTÖ, *Optimal inequalities between Seiffert's mean and power means*, Math. Inequal. Appl. **7**, 1 (2004), 47–53.
- [5] A. A. JAGER, *Solution of Problem 887*, Nieuw Arch. Wisk. **4**, 12 (1994), 230–231.
- [6] E. NEUMAN, J. SÁNDOR, *On the Schwab-Borchardt Mean*, Math. Pannon. **14**, 2 (2003), 253–266.
(Online: http://ttk.pte.hu/mii/html/pannonica/index_elemei/mp14-2/mp14-2-253-266.pdf).
- [7] H. LIU AND X.-J. MENG, *The Optimal Convex Combination Bounds for Seiffert's Mean*, J. Inequal. Appl. **2011**, 1, 44, doi:10.1186/1029-242X-2011-44.
- [8] J. SÁNDOR, *On certain inequalities for means, II*, J. Math. Anal. Appl. **199**, 2 (1996), 629–635.
- [9] J. SÁNDOR, *On certain inequalities for means, III*, Arch. Math. (Basel) **76** (2001), 34–40.
- [10] J. SÁNDOR, *Über zwei Mittel von Seiffert*, Die Wurzel **5** (2002), 104–107.
- [11] J. SÁNDOR, E. NEUMAN, *On certain means of two arguments and their extensions*, Int. J. Math. Math. Sci. **2003**, 16 (2003), 981–993, doi:10.1155/S0161171203208103 .
- [12] J. SÁNDOR, T. TRIF, *Some new inequalities for means of two arguments*, Int. J. Math. Math. Sci. **25**, 8 (2001), 525–532, doi:10.1155/S0161171201003064
- [13] H.-J. SEIFFERT, *Werte zwischen dem geometrischen und dem arithmetischen Mittel zweier Zahlen*, Elem. Math. **42** (1987), 105–107.
- [14] H.-J. SEIFFERT, *Problem 887*, Nieuw Arch. Wiskd. (Ser. 4) **11** (1993), 196.
- [15] H.-J. SEIFFERT, *Aufgabe β 16*, Die Wurzel **29** (1995), 221–222.
- [16] H.-J. SEIFFERT, *Ungleichungen für einen bestimmten Mittelwert*, Nieuw Arch. Wiskd. (Ser. 4), **13** (1995), 195–198.
- [17] S. WANG, Y. CHU, *The Best Bounds of the Combination of Arithmetic and Harmonic Means for the Seiffert's Mean*, Int. J. Math. Anal. **4**, 22 (2010), 1079–1084.

- [18] C. ZONG, Y. CHU, *An Inequality Among Identric, Geometric and Seiffert's Means*, Int. Math. Forum **5**, 26 (2010), 1297–1302.