STABILITY OF THE *n*-DIMENSIONAL MIXED-TYPE ADDITIVE AND QUADRATIC FUNCTIONAL EQUATION IN NON-ARCHIMEDEAN NORMED SPACES, II

YANG-HI LEE AND SOON-MO JUNG

Abstract. In this paper, we prove the stability of the functional equation

$$\sum_{\leq i,j \leq n, i \neq j} \left(f(x_i + x_j) + f(x_i - x_j) \right) = (n-1) \sum_{i=1}^n \left(3f(x_i) + f(-x_i) \right)$$

in non-Archimedean normed spaces.

1

Mathematics subject classification (2010): 39B82, 46S10, 39B52.

Keywords and phrases: Non-Archimedean space, mixed-type additive and quadratic function, generalized Hyers-Ulam stability.

REFERENCES

- T. AOKI, On the stability of the linear transformation in Banach spaces, J. Math. Soc. Japan 2 (1950), 64–66.
- [2] L. M. ARRIOLA AND W. A. BEYER, Stability of the Cauchy functional equation over p-adic fields, Real Anal. Exchange 31 (2005/2006), 125–132.
- [3] I.-S. CHANG, E.-H. LEE AND H.-M. KIM, On Hyers-Ulam-Rassias stability of a quadratic functional equation, Math. Inequal. Appl. 6 (2003), 87–95.
- [4] S. CZERWIK, Functional Equations and Inequalities in Several Variables, World Scientific, Hackensacks, New Jersey, 2002.
- [5] P. GĂVRUTA, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994), 431–436.
- [6] D. H. HYERS, On the stability of the linear functional equation, Proc. Natl. Acad. Sci. USA 27 (1941), 222–224.
- [7] D. H. HYERS, G. ISAC AND TH. M. RASSIAS, Stability of Functional Equations in Several Variables, Birkhäuser, Boston, 1998.
- [8] K.-W. JUN AND Y.-H. LEE, A Generalization of the Hyers-Ulam-Rassias stability of the Pexiderized quadratic equations II, Kyungpook Math. J. 47 (2007), 91–103.
- [9] S.-M. JUNG, Hyers-Ulam-Rassias stability of Jensen's equation and its application, Proc. Amer. Math. Soc. 126 (1998), 3137–3143.
- [10] S.-M. JUNG, Hyers-Ulam-Rassias Stability of Functional Equations in Nonlinear Analysis, Springer Optimization and Its Applications 48, Springer, New York, 2011.
- G.-H. KIM, On the stability of functional equations with square-symmetric operation, Math. Inequal. Appl. 4 (2001), 257–266.
- [12] H.-M. KIM, On the stability problem for a mixed type of quartic and quadratic functional equation, J. Math. Anal. Appl. 324 (2006), 358–372.
- [13] Y.-H. LEE, On the stability of the monomial functional equation, Bull. Korean Math. Soc. 45 (2008), 397–403.
- [14] Y.-H. LEE AND K.-W. JUN, A generalization of the Hyers-Ulam-Rassias stability of Jensen's equation, J. Math. Anal. Appl. 238 (1999), 305–315.



- [15] Y.-H. LEE AND K.-W. JUN, A generalization of the Hyers-Ulam-Rassias stability of Pexider equation, J. Math. Anal. Appl. 246 (2000), 627–638.
- [16] Y.-H. LEE, S.-M. JUNG AND TH. M. RASSIAS, Stability of the n-dimensional mixed-type additive and quadratic functional equation in non-Archimedean normed spaces, Abstr. Appl. Anal. 2012 (2012), Article ID 401762, 9 pages, (doi: 10.1155/2012/401762).
- [17] A. K. MIRMOSTAFAEE, Approximately additive mappings in non-Archimedean spaces, Bull. Korean Math. Soc. 46 (2009), 387–400.
- [18] M. S. MOSLEHIAN AND TH. M. RASSIAS, Stability of functional equations in non-Archimedean spaces, Appl. Anal. Discrete Math. 1 (2007), 325–334.
- [19] TH. M. RASSIAS, On the stability of the linear mapping in Banach spaces, Proc. Amer. Math. Soc. 72 (1978), 297–300.
- [20] F. SKOF, Local properties and approximations of operators, Rend. Sem. Mat. Fis. Milano 53 (1983), 113–129.
- [21] S. M. ULAM, A Collection of Mathematical Problems, Interscience, New York, 1960.
- [22] M. ABRAMOWITZ AND I. A. STEGUN (Eds), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, National Bureau of Standards, Applied Mathematics Series 55, 9th printing, Washington, 1970.
- [23] W. BURNSIDE, A rapidly convergent series for log N!, Messenger Math., 46, 1 (1917), 157–159.