

## SHARP BOUNDS FOR SEIFFERT MEAN IN TERMS OF WEIGHTED POWER MEANS OF ARITHMETIC MEAN AND GEOMETRIC MEAN

ZHEN-HANG YANG

**Abstract.** For  $a, b > 0$  with  $a \neq b$ , let  $P = (a - b)/(4 \arctan \sqrt{a/b} - \pi)$ ,  $A = (a + b)/2$ ,  $G = \sqrt{ab}$  denote the Seiffert mean, arithmetic mean, geometric mean of  $a$  and  $b$ , respectively. In this paper, we present new sharp bounds for Seiffert  $P$  in terms of weighted power means of arithmetic mean  $A$  and geometric mean  $G$ :

$$\left(\frac{2}{3}A^{p_1} + \frac{1}{3}G^{p_1}\right)^{1/p_1} < P < \left(\frac{2}{3}A^{p_2} + \frac{1}{3}G^{p_2}\right)^{1/p_2},$$

where  $p_1 = 4/5$  and  $p_2 = \log_{\pi/2}(3/2)$  are the best possible constants. Moreover, our sharp bounds for  $P$  are compared with other known ones, which yields a chain of inequalities involving Seiffert mean  $P$ .

**Mathematics subject classification (2010):** Primary 26E60, 26D05; secondary 33B10.

**Keywords and phrases:** Seiffert mean, arithmetic mean, geometric mean, power mean, sharp bound.

### REFERENCES

- [1] P. S. BULLEN, D. S. MITRINOVÍĆ AND P. M. VASIĆ, *Means and Their Inequalities*, Dordrecht 1988.
- [2] CH.-P. CHEN AND W.-S. CHEUNG, *Sharp Cusa and Becker-Stark inequalities*, J. Inequal. Appl., (2011), 136; available online at <http://www.journalofinequalitiesandapplications.com/content/2011/1/136>.
- [3] Y.-M. CHU, Y.-F. QIU, M.-K. WANG, AND G.-D. WANG, *The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean*, J. Inequal. Appl., (2010), Art. ID 436457, 7 pages.
- [4] Y.-M. CHU, Y.-F. QIU, AND M.-K. WANG, *Sharp power mean bounds for the combination of seiffert and geometric means*, Abstr. Appl. Anal., (2010), Art. ID 108920, 12 pages.
- [5] P. A. HÄSTÖ, *A monotonicity property of ratios of symmetric homogeneous means*, J. Inequal. Pure Appl. Math., **3**, 5 (2002), Art. 71, pp. 23.
- [6] P. A. HÄSTÖ, *Optimal inequalities between Seiffert's mean and power mean*, Math. Inequal. Appl., **7**, 1 (2004), 47–53.
- [7] D. HE AND ZH.-J. SHEN, *Advances in research on Seiffert mean*, Communications in inequalities research, **17**, 4 (2010), Art. 26; available online: <http://old.irgoc.org/Article/UploadFiles/201010/20101026104515652.pdf>.
- [8] C. HUYGENS, *Oeuvres Complètes 1888-1940*, Société Hollondaise des Science, Haga.
- [9] A. A. JAGERS, *Solution of problem 887*, Nieuw Arch. Wisk., **12** (1994), 2 30–231.
- [10] R. KLEŃ, M. VISURI AND M. VUORINEN, *On Jordan type inequalities for hyperbolic functions*, J. Inequal. Appl., (2010), Art. ID 362548, 14 pages.
- [11] O. KOUBA, *New bounds for the identric mean of two arguments*, J. Inequal. Pure Appl. Math., **9**, 3 (2008), Art. 71, 6 pages.
- [12] H. LIU AND X.-J. MENG, *The optimal convex combination bounds for Seiffert's mean*, J. Inequal. Appl., (2011), Art. ID 686834, 9 pages.
- [13] C. MORTITI, *The natural approach of Wilker-Cusa-Huygens inequalities*, Math. Inequal. Appl., **14**, 3 (2011), 535–541.
- [14] E. NEUMAN AND J. SÁNDOR, *On the Schwab-Borchardt mean*, Math. Pannon., **17**, 1 (2006), 49–59.

- [15] E. NEUMAN, *On Wilker and Huygens inequalities*, Math. Inequal. Appl., **15**, 2 (2012), 271–279.
- [16] J. SÁNDOR, *On certain inequalities for means III*, Arch. Math., **76** (2001), 34–40.
- [17] H.-J. SEIFFERT, *Werte zwischen dem geometrischen und dem arithmetischen Mittel zweier Zahlen*, Elem. Math., **42** (1987), 105–107.
- [18] H.-J. SEIFFERT, *Problem 887*, Nieuw Arch. Wisk., **4**, 11 (1993), 176.
- [19] H.-J. SEIFFERT, *Aufgabe 16*, Die Wurzel, **29** (1995), 221–222.
- [20] S.-S. WANG AND Y.-M. CHU, *The best bounds of the combination of arithmetic and harmonic means for the Seiffert's mean*, Int. J. Math. Anal. (Ruse), **4**, 21–24 (2010), 1079–1084.
- [21] M.-K. WANG, Y.-F. QIU, AND Y.-M. CHU, *Sharp bounds for Seiffert means in terms of Lehmer means*, J. Math. Inequal., **4**, 4 (2010), 581–586.
- [22] L. ZHU, *Inequalities for hyperbolic functions and their applications*, J. Inequal. Appl., **2010** (2010), Art. ID 130821, 10 pages.