

ON MULTIPLE FOURIER COEFFICIENTS OF FUNCTIONS OF ϕ - Λ -BOUNDED VARIATION

R. G. VYAS AND K. N. DARJI

Abstract. Here, we have estimated the order of magnitude of multiple Fourier coefficients of functions of $\phi(\Lambda^1, \dots, \Lambda^N)BV([0, 2\pi]^N)$.

Mathematics subject classification (2010): 42B05, 26B30, 26D15.

Keywords and phrases: Order of magnitude of multiple Fourier coefficients, functions of $\phi(\Lambda^1, \dots, \Lambda^N)BV$.

REFERENCES

- [1] A. N. BAKHVALOV, *Fourier coefficients of functions from many-dimensional classes of bounded Λ -variation*, Moscow Univ. Math. Bulletin, **66**, 1 (2011), 8–16.
- [2] V. FÜLÖP AND F. MÓRICZ, *Order of magnitude of multiple Fourier coefficients of functions of bounded variation*, Acta Math. Hungar., **104**, 1–2 (2004), 95–104.
- [3] S. R. GHORPADE AND B. V. LIMAYE, *A Course in Multivariable Calculus and Analysis*, Springer, 2010.
- [4] F. MÓRICZ AND A. VERES, *On the absolute convergence of multiple Fourier series*, Acta Math. Hungar., **117**, 3 (2007), 275–292.
- [5] M. SCHRAMM AND D. WATERMAN, *On the magnitude of Fourier coefficients*, Proc. Amer. Math. Soc., **85**, (1982), 407–410.
- [6] R. G. VYAS, *On the absolute convergence of small gaps Fourier series of functions of $\phi\Lambda BV$* , JIPAM. J. Inequal. Pure Appl. Math. **6** (2005), no. 4, Article 94, 5 pp.
- [7] R. G. VYAS AND K. N. DARJI, *On multiple Walsh Fourier coefficients*, J. Indian Math. Soc., **79**, 1–4 (2012), 219–228.