

## A NOTE ON ESTIMATES OF THE NEWTONIAN POTENTIAL ON BOUNDED DOMAINS

TRAN VU KHANH\* AND ANDREW RAICH

*Abstract.* In this paper, we investigate the regularity of the operator  $K$  on a smooth bounded domain in  $\mathbb{R}^d$  given by convolution against the Newtonian potential. We show that the gain in  $L^p$ -Sobolev spaces agrees with elliptic regularity. We also establish  $L^p$ -Sobolev to  $L^q$ -Sobolev bounds as well as bounds from  $L^\infty$ -Sobolev spaces to Hölder spaces.

*Mathematics subject classification (2020):* 42B20, 35B65, 42B37, 35C15, 35J05, 32A26, 32A50.

*Keywords and phrases:* Newtonian potential, Sobolev regularity, non-characteristic formula, Bochner-Martinelli-Koppelman kernel, Hölder regularity.

### REFERENCES

- [1] Z. CHEN, S. KRANTZ, AND D. MA, *Optimal  $L^p$  estimates for the  $\bar{\partial}$ -equation on complex ellipsoids in  $\mathbb{C}^n$* , *Manuscripta Math.*, **80** (1993), pp. 131–149.
- [2] L. EVANS, *Partial differential equations*, vol. 19 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, Second ed., 2010.
- [3] G. FOLLAND AND E. STEIN, *Estimates for the  $\bar{\partial}_b$ -complex and analysis on the Heisenberg group*, *Comm. Pure and Appl. Math.*, **27** (1974), pp. 429–522.
- [4] G. B. FOLLAND, *Real Analysis : Modern techniques and their applications*, Pure and Applied Mathematics (New York), John Wiley & Sons Inc., New York, Second ed., 1999.
- [5] D. GILBARG AND N. S. TRUDINGER, *Elliptic partial differential equations of second order*, *Classics in Mathematics*, Springer-Verlag, Berlin, 2001, reprint of the 1998 edition.
- [6] K. HA, T. KHANH, AND A. RAICH,  *$L^p$ -estimates for the  $\bar{\partial}$ -equation on a class of infinite type domains*, *Internat. J. Math.*, **25** (2014), p. 1450106 (15 pages), doi:10.1142/S0129167X14501067.
- [7] P. HARRINGTON AND A. RAICH, *Defining functions for unbounded  $C^m$  domains*, *Rev. Mat. Iberoam.*, **29** (2013), pp. 1405–1420.
- [8] L. HÖRMANDER,  *$L^2$  estimates and existence theorems for the  $\bar{\partial}$  operator*, *Acta Math.*, **113** (1965), pp. 89–152.
- [9] C. KENIG, *Harmonic analysis techniques for second order elliptic boundary value problems*, vol. 83 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1994.
- [10] T. KHANH, *Supnorm and  $f$ -Hölder estimates for  $\bar{\partial}$  on convex domains of general type in  $\mathbb{C}^2$* , *J. Math. Anal. Appl.*, **403** (2013), pp. 522–531.
- [11] T. KHANH AND A. RAICH,  *$L^p$ -estimates for the  $\bar{\partial}_b$ -equation on a class of infinite type domains*, *Math. Nachr.*, **294** (2021), pp. 82–97.
- [12] M. RANGE, *Integral kernels and Hölder estimates for  $\bar{\partial}$  on pseudocconvex domains of finite type in  $\mathbb{C}^2$* , *Math. Ann.*, **288** (1990), pp. 63–74.
- [13] E. STRAUBE, *Lectures on the  $\mathcal{L}^2$ -Sobolev Theory of the  $\bar{\partial}$ -Neumann Problem*, *ESI Lectures in Mathematics and Physics*, European Mathematical Society (EMS), Zürich, 2010.