STRUCTURED DECOMPOSITIONS FOR MATRIX TRIPLES: SVD–LIKE CONCEPTS FOR STRUCTURED MATRICES

Christian Mehl, Volker Mehrmann and Hongguo Xu

Abstract. Canonical forms for matrix triples $(A, G, \hat{G})$, where A is arbitrary rectangular and G, $\hat{G}$ are either real symmetric or skew symmetric, or complex Hermitian or skew Hermitian, are derived. These forms generalize classical singular value decompositions. In [1] a similar canonical form has been obtained for the complex case. In this paper, we provide an alternative proof for the complex case which is based on the construction of a staircase-like form with the help of a structured QR-like decomposition. This approach allows generalization to the real case.

REFERENCES

[1] Y. Bolschakov and B. Reichstein, *Unitary equivalence in an indefinite scalar product: an analogue of singular-value decomposition*, Linear Algebra Appl., **222** (1995), 155–226.

[2] Y. Bolschakov, C.V.M. van der Mee, A.C.M. Ran, B. Reichstein, and L. Rodman, *Polar decompositions in finite-dimensional indefinite scalar product spaces: general theory*, Linear Algebra Appl., **261** (1997), 91–141.

[3] D.Z. Djokovic, J. Patera, P. Winternitz, and H. Zassenhaus, *Normal forms of elements of classical real and complex Lie and Jordan algebras*, J. Math. Phys., **24** (1983), 1363–1374.

[4] N.M. Dobrovol'skaya and V.A. Ponomarev, *A pair of counter-operators* (in Russian), Uspehi Mat. Nauk, **20** (1965), 80–86.

[5] H. Fassbender, D. S. Mackey, N. Mackey, and H. Xu, *Hamiltonian square roots of skew–Hamiltonian matrices*, Linear Algebra Appl., **287** (1999), 125–159.

[6] H. Flanders, *Elementary divisors of AB and BA*, Proc. Amer. Math. Soc., **2** (1951), 871–874.

[7] I. Gohberg, P. Lancaster, and L. Rodman, *Matrices and Indefinite Scalar Products*, Birkhäuser, Basel, 1983.

[8] I. Gohberg, P. Lancaster, and L. Rodman, *Indefinite Linear Algebra and Applications*, Birkhäuser, Basel, 2005.

[9] G. H. Golub and C. F. Van Loan, *Matrix Computations*, Johns Hopkins University Press, 3rd edition, Baltimore, 1996.

[10] S. Hassi, *Matriisin yleisen singulaariarvohajoitelman ominaisuuksia ja tilastollisia sovelluksia*, Master's thesis, University of Helsinki, 1984.

[11] S. Hassi, *A singular value decomposition of matrices in a space with an indefinite scalar product*, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes, **79**, 1990.

[12] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.

[13] R. A. Horn and D. Merino, *Contragredient equivalence: a canonical form and some applications*, Linear Algebra Appl., **214** (1995), 43–92.

[14] T. Kapitula, P.G. Kevrekidis, and B. Sandstede, *Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems*, Physica D: Nonlinear Phenomena, **195** (2004), 263–282.

[15] J. Kes and A.C.M. Ran, *On the relation between $XX^{[*]}$ and $X^{[*]}X$ in an indefinite inner product space*, Oper. Matrices, **1** (2007), 181–197.

[16] P. Lancaster and L. Rodman, *The Algebraic Riccati Equation*, Oxford University Press, Oxford, 1995.

[17] P. Lancaster and L. Rodman, *Canonical forms for Hermitian matrix pairs under strict equivalence and congruence*, SIAM Rev., **47**, (2005), 407–443.

[18] P. Lancaster and L. Rodman, *Canonical forms for symmetric/skew symmetric real pairs under strict equivalence and congruence*, Linear Algebra Appl., **406** (2005), 1–76.

[19] C. Mehl, V. Mehrmann, and H. Xu, *Singular-value-like decompositions for complex matrix triples*, J. Comput. Appl. Math., to appear, 2009.
`http://dx.doi.org/10.1016/j.cam.2008.02.017`.

[20] C. Mehl, A.C.M. Ran, and L. Rodman, *Polar decompositions of normal operators in indefinite inner product spaces*, Oper. Theory Adv. Appl., **162** (2006), 277–292.

[21] V. Mehrmann and H. Xu, *Structured Jordan canonical forms for structured matrices that are Hermitian, skew Hermitian or unitary with respect to indefinite inner products*, Electron J. Linear Algebra, **5** (1999), 67–103.

[22] J. Patera and C. Rousseau, *Versal deformations of elements of classical Jordan algebras*, J. Math. Phys., **24** (1983), 1375–1379.

[23] A. C. M. Ran and M. Wojtylak, *Analysis of spectral points of the operators $T^{[*]}T$ and $TT^{[*]}$ in a Krein space*, submitted, 2008.

[24] V.V. Sergeichuk, *Computation of canonical matrices for chains and cycles of linear mappings*, Linear Algebra Appl., **376** (2004), 235–263.

[25] R. C. Thompson, *The characteristic polynomial of a principal submatrix of a Hermitian pencil*, Linear Algebra Appl., **14** (1976), 135–177.

[26] R. C. Thompson, *Pencils of complex and real symmetric and skew matrices*, Linear Algebra Appl., **147** (1991), 323–371.

[27] H. Xu, *An SVD-like matrix decomposition and its applications*, Linear Algebra Appl., **368** (2003), 1–24.

[28] H. Xu, *A numerical method for computing an SVD-like decomposition*, SIAM J. Matrix Anal. Appl., **26** (2005), 1058–1082.

Operators and Matrices
www.ele-math.com
oam@ele-math.com