

## BISHOP'S PROPERTY ( $\beta$ ) FOR PARANORMAL OPERATORS

ATSUSHI UCHIYAMA AND KOTARO TANAHASHI

*Abstract.* For an operator  $T$  on a separable complex Hilbert space  $\mathcal{H}$ , we say that  $T$  has Bishop's property ( $\beta$ ) if for any open subset  $\mathcal{D} \subset \mathbb{C}$  and any sequence of analytic functions  $f_n : \mathcal{D} \rightarrow \mathcal{H}$  such as  $\|(T - z)f_n(z)\| \rightarrow 0$  as  $n \rightarrow \infty$  uniformly on every compact subset  $\mathcal{K} \subset \mathcal{D}$ , then  $f_n \rightarrow 0$  uniformly on  $\mathcal{K}$ . It is a very important property in spectral theory. It is well-known that every normal operator ( $T^*T = TT^*$ ) has Bishop's property ( $\beta$ ). Now, many mathematicians attempt to extend this result to non-normal operators.

In this paper, we shall show that every paranormal operator ( $\|T^2x\|\|x\| \geq \|Tx\|^2$  for all  $x \in \mathcal{H}$ ) has Bishop's property ( $\beta$ ).

*Mathematics subject classification (2000):* 47B20.

*Keywords and phrases:* Bishop's property ( $\beta$ ), paranormal operator.

### REFERENCES

- [1] A. ALUTHGE, *On  $p$ -hyponormal operators for  $0 < p < 1$* , Integral Equations and Operator Theory, **13** (1990), 307–315.
- [2] A. ALUTHGE AND D. WANG,  *$w$ -hyponormal operators*, Integral Equations and Operator Theory, **36** (2000), 1–10.
- [3] M. CHŌ AND T. HURUYA,  *$p$ -hyponormal operators for  $0 < p < \frac{1}{2}$* , Comment Math., **33** (1993), 23–29.
- [4] M. CHŌ AND T. YAMAZAKI, *An operator transform class  $A$  to the class of hyponormal operators and its application*, Integral Equations and Operator Theory, **53** (2005), 497–508.
- [5] T. FURUTA, *On the class of paranormal operators*, Proc. Japan Acad., **43** (1967), 594–598.
- [6] T. FURUTA, M. ITO AND T. YAMAZAKI, *A subclass of paranormal operators including class of log-hyponormal and several related classes*, Sci. Math., **1** (1998), 389–403.
- [7] V. ISTRĂȚESCU, T. SAITŌ AND T. YOSHINO, *On a class of operators*, Tôhoku Math. J., (2), **18** (1966), 410–413.
- [8] F. KIMURA, *Analysis of non-normal operators via Aluthge transformation*, Integral Equations and Operator Theory, **50** (2004), 375–384.
- [9] K. B. LAUSEN, *Operators with finite ascent*, Pacific J. Math., **152** (1992), 323–336.
- [10] A. UCHIYAMA, *Weyl's theorem for class  $A$  operators*, Math. Inequal. & Appl., **4** (2001), 143–150.
- [11] A. UCHIYAMA, K. TANAHASHI AND J. I. LEE, *Spectrum of class  $A(s,t)$  operators*, Acta Sci. Math. (Szeged), **70** (2004), 279–287.
- [12] D. XIA, *On the non-normal operators—semihyponormal operators*, Sci. Sinica., **23** (1980), 700–713.