

ACCUMULATION OF COMPLEX EIGENVALUES OF AN INDEFINITE STURM—LIOUVILLE OPERATOR WITH A SHIFTED COULOMB POTENTIAL

MICHAEL LEVITIN AND MARCELLO SERI

Abstract. For a particular family of long-range potentials V , we prove that the eigenvalues of the indefinite Sturm–Liouville operator $A = \text{sign}(x)(-\Delta + V(x))$ accumulate to zero asymptotically along specific curves in the complex plane. Additionally, we relate the asymptotics of complex eigenvalues to the two-term asymptotics of the eigenvalues of associated self-adjoint operators.

Mathematics subject classification (2010): 34B24, 34L15, 47E05, 47B50, 33C15.

Keywords and phrases: Linear operator pencils, non-self-adjoint operators, Sturm–Liouville problem, Coulomb potential, complex eigenvalues, Kummer functions.

REFERENCES

- [1] T. YA. AZIZOV AND I. S. IOKHVIDOV, *Linear operators in spaces with an indefinite metric*, John Wiley, 1989
- [2] J. BEHRNDT, *On the spectral theory of singular indefinite Sturm–Liouville operators*, J. Math. Anal. Appl. **334** (2007), 1439–1449.
- [3] J. BEHRNDT, *An open problem: accumulation of nonreal eigenvalues of indefinite Sturm–Liouville operators*, Integral Equations Operator Theory **77** (2013), no. 3, 299–301.
- [4] J. BEHRNDT AND Q. KATATBEH AND C. TRUNK, *Accumulation of complex eigenvalues of indefinite Sturm–Liouville operators*, J. Phys. A **41** (2008), no. 24, 244003.
- [5] J. BEHRNDT AND F. PHILIPP AND C. TRUNK, *Bounds for the non-real spectrum of differential operators with indefinite weights*, Math. Ann. **357** (2013), 185–213.
- [6] J. BEHRNDT AND C. TRUNK, *On the negative squares of indefinite Sturm–Liouville operators*, Journal of Differential Equations **238** (2007), 491–519.
- [7] B. ČURGUS AND H. LANGER, *A Krein space approach to symmetric ordinary differential operators with an indefinite weight function*, J. Differential Equations **79** (1989), 31–61.
- [8] B. ČURGUS AND B. NAJMAN, *The operator $(\text{sgn} x) \frac{d^2}{dx^2}$ is similar to a self-adjoint operator in $L^2(\mathbb{R})$* , Proc. Amer. Math. Soc. **123** (1995), 1125–1128.
- [9] E. B. DAVIES AND M. LEVITIN, *Spectra of a class of non-self-adjoint matrices*, Linear Algebra Appl. **448** (2014), 55–84.
- [10] D. M. ELTON AND M. LEVITIN AND I. POLTEROVICH, *Eigenvalues of a one-dimensional Dirac operator pencil*, Annales H. Poincaré **15** (2014), 2321–2377.
- [11] M. M. FADDEEV AND R. G. SHTERENBERG, *On the similarity of some singular differential operators to selfadjoint operators*, Zap. Nauchn. Sem. POMI **270** (2000), 336–349.
- [12] A. FLEIGE, *The critical point infinity associated with indefinite Sturm–Liouville problems*, in: *Operator Theory*, D. ALPAY (ED). Springer, Basel (2015) (to appear)
- [13] F. GESZTESY, *On the one-dimensional Coulomb Hamiltonian*, J. Phys. A: Math. Gen. **13** (1980), 867–875.
- [14] N. J. HIGHAM AND F. TISSEUR AND P. M. VAN DOOREN, *Detecting a definite Hermitian pair and a hyperbolic or elliptic quadratic eigenvalue problem, and associated nearness problems*, Lin. Alg. and its App. **351** (2002), 455–474.

- [15] I. M. KARABASH, *A functional model, eigenvalues, and finite singular critical points for indefinite Sturm–Liouville operators*, Oper. Theory Adv. Appl. **203** (2010), 247–287.
- [16] I. M. KARABASH AND A. S. KOSTENKO AND M. M. MALAMUD, *The similarity problem for J -nonnegative Sturm–Liouville operators*, J. Differential Equations **246** (2009), no. 3, 964–997.
- [17] I. M. KARABASH AND M. M. MALAMUD, *On the similarity of the J -selfadjoint Sturm–Liouville operator with a finite-gap potential to a selfadjoint one*, (Russian) Dokl. Akad. Nauk **395** (2004), no. 3, 303–307.
- [18] I. M. KARABASH AND M. M. MALAMUD, *Indefinite Sturm–Liouville operators $(\operatorname{sgn}x)(-\frac{d^2}{dx^2} + q(x))$ with finite-zone potentials*, Oper. Matrices **1** (2007), no. 3, 301–368.
- [19] I. M. KARABASH AND C. TRUNK, *Spectral properties of singular Sturm–Liouville operators with indefinite weight $\operatorname{sgn} x$* , Proc. Royal Soc. Edinb.: Sect. A **139** (2009), no. 3, 483–503.
- [20] A. KOSTENKO, *The similarity problem for indefinite Sturm–Liouville operators and the HELP inequality*, Adv. Math. **246** (2013), 368–413.
- [21] H. LANGER, *Krein space*, in: *Encyclopedia of Mathematics*, available at http://www.encyclopediaofmath.org/index.php?title=Krein_space&oldid=18988
- [22] A. S. MARKUS, *Introduction to the Spectral Theory of Polynomial Operator Pencils*, Amer. Math. Soc., Providence, RI, 1988.
- [23] F. W. J. OLVER AND D. W. LOZIER AND R. F. BOISVERT AND C. W. CLARK (EDS), *NIST handbook of mathematical functions*, U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010, available at <http://dlmf.nist.gov>
- [24] S. G. PYATKOV, *Maximal semidefinite invariant subspaces for J -dissipative operators*, Oper. Theory Adv. Appl. **221** (2012), 549–570.
- [25] N. M. TEMME, *Asymptotic Methods for Integrals*, World Scientific, Singapore, 2015.
- [26] H. VAN HAERINGEN, *The bound states for the symmetric shifted Coulomb potential*, J. Math. Phys. **19** (1978), no. 10, 2165–2170.
- [27] O. VERDIER, *Reductions of operator pencils*, Math. Comp. **83** (2014), 189–214.
- [28] D. R. YAFAEV, *The low energy scattering for slowly decreasing potentials*, Comm. Math. Phys. **85** (1982), no. 2, 177–196.