

OPERATOR MONOTONICITY OF A 2-PARAMETER FAMILY OF FUNCTIONS AND $\exp\{f(x)\}$ RELATED TO THE STOLARSKY MEAN

YOICHI UDAGAWA

Abstract. We consider operator monotonicity of a 2-parameter family of functions including the representing function of the Stolarsky mean, which is constructed by integration of the function $[(1-\alpha)+\alpha x^p]^{\frac{1}{p}}$, representing the weighted power mean, of $\alpha \in [0,1]$. We also think about operator monotonicity of $\exp\{f(x)\}$ for a continuous function f(x) defined on $(0,\infty)$.

Mathematics subject classification (2010): 15A60, 47A64.

Keywords and phrases: Operator mean, operator monotone function, Stolarsky mean, identric mean.

REFERENCES

- [1] F. HANSEN, Selfadjoint means and operator monotone functions, Math. Ann. 256 (1981), 29–35.
- [2] F. HIAI AND H. KOSAKI, Means for matrices and comparison of their norms, Indiana Univ. Math. J. 48 (1999), 899–936.
- [3] F. HIAI AND D. PETZ, Introduction to Matrix Analysis and Applications, Universitext, Springer, 2014.
- [4] F. Kubo and T. Ando, Means of positive linear operators, Math. Ann. 246 (1979/80), 205-224.
- [5] M. NAGISA, S. WADA, Operator monotonicity of some functions, Linear Algebra Appl., 486 (2015), 389–408.
- [6] Y. NAKAMURA, Classes of operator monotone functions and Stieltjes functions, The Gohberg anniversary collection, vol. II (Calgary, AB, 1988), 395–404, Oper. Theory Adv. Appl. 41, Birkhäuser, Basel 1989
- [7] Y. UDAGAWA, S. WADA, T. YAMAZAKI AND M. YANAGIDA, On a family of operator means involving the power difference means, Linear Algebra Appl. 485 (2015), 124–131.

