

## WEAK SUBNORMALITY OF INFINITE 4-BANDED MATRICES

IN SUNG HWANG, IN HYOUN KIM AND SUMIN KIM

*Abstract.* In this paper, we consider a class of operators whose matrix representations comprise 4-banded matrices, i.e., sparse matrices whose non-zero entries are confined to four diagonals. In particular, we focus on the hyponormality and weak subnormality when each diagonal forms a hyponormal weighted shift.

*Mathematics subject classification (2010):* 47B20, 47B37, 47A63.

*Keywords and phrases:* weak subnormality, hyponormality, 4-banded matrices, weighted shifts.

### REFERENCES

- [1] C. COWEN AND J. LONG, *Some subnormal Toeplitz operators*, J. Reine Angew. Math. **351** (1984), 216–220.
- [2] R. E. CURTO, I. B. JUNG, AND S. S. PARK, *A characterization of  $k$ -hyponormality via weak subnormality*, J. Math. Anal. Appl. **279** (2) (2003), 556–568.
- [3] R. E. CURTO, S. H. LEE, AND W. Y. LEE, *A new criterion for  $k$ -hyponormality via weak subnormality*, Proc. Amer. Math. Soc. **133** (6) (2004), 1805–1816.
- [4] R. E. CURTO AND W. Y. LEE, *Towards a model theory for 2-hyponormal operators*, Integral Equations Operator Theory **44** (2002), 290–315.
- [5] C. GU, I. S. HWANG AND W. Y. LEE, *Hyponormality of Toeplitz operators in several variables by the weighted shifts approach*, Linear and Multilinear Algebra **68** (8) (2020), 1695–1720.
- [6] I. S. HWANG, *Hyponormality of Toeplitz operators on the Bergman space*, J. Korean Math. Soc. **45** (2008), 1027–1041.
- [7] J. LEE, *Hyponormality of Toeplitz operators on the weighted Bergman space with matrix-valued circulant symbols*, Linear Algebra Appl. **576** (2019), 35–50.