

ANALYSIS OF THE BOUNDARY VALUE PROBLEM ASSOCIATED WITH THE NONRELATIVISTIC THOMAS–FERMI EQUATION FOR HEAVY ATOMS IN INTENSE MAGNETIC FIELDS

CHRISTOPHER C. TISDELL AND MARK HOLZER

Abstract. This article presents a firm mathematical foundation for the boundary value problem associated with the nonrelativistic Thomas-Fermi equation for heavy atoms in intense magnetic fields. Our approach uses an application of differential inequalities and ideas from nonlinear analysis, including: the technique of lower and upper solutions; and fixed-point theory. We present new results that ensure existence, uniqueness, location and approximation of solutions. We thus establish that the Thomas-Fermi model leads to a robust theory of heavy atoms in intense magnetic fields in spite of the severe approximations that it employs.

A YouTube video from the first author that is designed to complement this paper can be found here <http://tinyurl.com/ThomasFermi>.

Mathematics subject classification (2010): 34B60, 34B15.

Keywords and phrases: Thomas-Fermi equation, boundary value problem, heavy atoms in intense magnetic fields, qualitative analysis of solutions, existence, uniqueness and location of solutions.

REFERENCES

- [1] R.P. AGARWAL, *Boundary value problems for higher order differential equations*. World Scientific, Singapore, 1986.
- [2] E. AKKERMANS, S. GHOSH, Z.H MUSSIMANI, *Numerical study of one-dimensional and interacting Bose-Einstein condensates in a random potential*, J. Phys. B: At. Mol. Opt. Phys., **41** (2008), no. 4, Article ID 045302 (12pp).
- [3] D.R. BALLANTYNE, *Reflection spectra from an accretion disc illuminated by a neutron star X-ray burst*, Mon. Not. R. Astron. Soc., **351** (2004), 57–62.
- [4] M.A. BAUTISTA, *Electronic correlations and polarizability of the Thomas-Fermi-Dirac-Amaldi potential: applications to the singly ionized iron-peak species*, J. Phys. B: At. Mol. Opt. Phys. **41** (2008), no. 6, Article ID 065701 (8pp).
- [5] PH. BÉNILAN, H. BRÉZIS, *Nonlinear problems related to the Thomas-Fermi equation. Dedicated to Philippe Bénilan*, J. Evol. Equ., **3** (2003), no. 4, 673–770.
- [6] R. BENGURIA, H. BRÉZIS, E.H. LIEB, *The Thomas-Fermi-von Weizsäcker theory of atoms and molecules*, Comm. Math. Phys., **79** (1981), no. 2, 167–180.
- [7] R. BENGURIA, E.H. LIEB, *Many-body atomic potentials in Thomas-Fermi theory*, Ann. Phys. NY **110**, (1978) 34–45.
- [8] R. BENGURIA, E.H. LIEB, *The positivity of the pressure in Thomas-Fermi theory*, Comm. Math. Phys., **63** (1978), 193–218.
- [9] R.K. BHADURI, V.M.N. MURTHY, M. BRACK, *Fermionic ground state at unitarity and Haldane exclusion statistics*, J. Phys. B: At. Mol. Opt. Phys., **41** (2008), no. 11, Article ID 115301 (4pp).
- [10] H. BRÉZIS, *Nonlinear problems related to the Thomas-Fermi equation. Contemporary developments in continuum mechanics and partial differential equations (Proc. Internat. Sympos., Inst. Mat., Univ. Fed. Rio de Janeiro, Rio de Janeiro, 1977)*, pp. 81–89, North-Holland Math. Stud., **30**, North-Holland, Amsterdam-New York, 1978.

- [11] H. BRÉZIS, *Nonlinear equations of Thomas Fermi type*. Proceedings of the First Franco–Southeast Asian Mathematical Conference (Nanyang Univ., Singapore, 1979). Southeast Asian Bull. Math. 1979, Special Issue, 167–171.
- [12] H. BRÉZIS, *Some variational problems of the Thomas-Fermi type. Variational inequalities and complementarity problems* (Proc. Internat. School, Erice, 1978), pp. 53–73, Wiley, Chichester, 1980.
- [13] H. BRÉZIS, *A free boundary problem in quantum mechanics: Thomas-Fermi equation*. Free boundary problems, Vol. II (Pavia, 1979), pp. 85–91, Ist. Naz. Alta Mat. Francesco Severi, Rome, 1980.
- [14] H. BRÉZIS, E.H. LIEB, *Long range atomic potentials in Thomas-Fermi theory*, Comm. Math. Phys., **65** (1979), no. 3, 231–246.
- [15] M.A. CARACANHAS, J.A. SEMAN, E.R.F. RAMOS, E.A.L. HENN, K.M.F. MAGALHAES, K. HELMERSON, V.S. BAGNATO, *Finite temperature correction to the Thomas-Fermi approximation for a Bose-Einstein condensate: comparison between theory and experiment*, J. Phys. B: At. Mol. Opt. Phys., **42** (2009), no. 14, Article ID 145304, (8pp).
- [16] C. DE COSTER, P. HABETS, *The lower and upper solutions method for boundary value problems*. Handbook of differential equations, 69–160, Elsevier North–Holland, Amsterdam, 2004.
- [17] P.A.M. DIRAC, *Note on the exchange phenomena in the Thomas atom*, Proc. Cambridge Philos. Soc., **26** (1930), 376.
- [18] E. FERMI, *Un metodo statistico per la determinazione di alcune priorietà dell'atome*, Rend. Addad. Naz. Lincei **6** (1927), 602.
- [19] M.D. GLOSSMAN, E.A. CASTRO, *Approximate analytical solution of the Thomas-Fermi equation for positive ions in a strong magnetic field*, Z. Phys. D - Atoms, Molecules and Clusters, **6** (1987), 81–82.
- [20] A. GRANAS, R.B. GUENTHER, J.W. LEE, *A note on the Thomas-Fermi equation*, Z. Angew. Math. Mech., **61** (1981), no. 3, 204–205.
- [21] M. GUPTA, K.R. KRISHNA, *An analytical approach for the nonlinear modified Thomas-Fermi equation to derive the ground-state and dynamic properties of a spherically and cylindrically trapped Bose-Einstein condensate*, J. Phys. B: At. Mol. Opt. Phys., **41** (2008), no. 19, Article ID 195302, (8pp).
- [22] P. HARTMAN, *Ordinary differential equations*. Corrected reprint of the second (1982) edition. With a foreword by Peter Bates. Classics in Applied Mathematics, 38. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2002.
- [23] S.H. HILL, P.J. GROUT, N.H. MARCH, *Relativistic Thomas-Fermi equation in an extremely high magnetic field, and its solution for atomic ions*, J. Phys. B: At. Mol. Phys., **18** (1985), 4665–4673.
- [24] S.H. HILL, P.J. GROUT, N.H. MARCH, *Chemical potential and total energy of heavy positive ions in extremely strong magnetic fields, near the weak ionisation limit*, J. Phys. B: At. Mol. Phys., **16** (1983), 2301–2307.
- [25] E. HILLE, *On the Thomas-Fermi equation*, Proc. Nat. Acad. Sci. U.S.A., **62** (1969), 7–10.
- [26] A.I. IBRAHIM, S. SAFI-HARB, J.H. SWANK, W. PARKE, S. ZANE, R. TUROLLA, *Discovery of cyclotron resonance features in the soft gamma repeater SGR 1806–20*. The Astrophysical Journal, 574:L51–L55, (2002).
- [27] B.B. KADOMSTEV, *Heavy atom in an ultrastrong magnetic field*, Soviet Physics JETP, **31** (1970), no. 5, 945–947.
- [28] A.S. KORNEV, B.A. ZON, *Rydberg spectra of atoms and positive ions in the Thomas-Fermi model*, J. Phys. B: At. Mol. Opt. Phys., **36** (2003), no. 19, 4027–4034.
- [29] E.H. LIEB, J.P. SOLOVEJ, J. YNGVASON, *Asymptotics of Heavy atoms in High magnetic Fields: I. Lowest Landau Band Regions*, Commun. Pure. Appl. Math., **47** (1994), 513–591.
- [30] E.H. LIEB, J.P. SOLOVEJ, J. YNGVASON, *Asymptotics of Heavy Atoms in High Magnetic Fields: II. Semiclassical Regions*, Commun. Math. Phys., **161** (1994), 77–124.
- [31] E.H. LIEB, THOMAS-FERMI AND HARTREE-FOCK THEORY. *Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974)*, Vol. 2, pp. 383–386. Canad. Math. Congress, Montreal, Que., 1975.
- [32] E.H. LIEB, *The stability of matter*, Rev. Mod. Phys., **48** (1976), 553–569.
- [33] E.H. LIEB, B. SIMON, *Thomas-Fermi theory revisited*, Phys. Rev. Lett., **31** (1973), 681–683.
- [34] E.H. LIEB, B. SIMON, *The Thomas-Fermi theory of atoms, molecules and solids*, Adv. Math., **23**, (1977), 22–116.

- [35] R.O. MUELLER, A.R.P. RAU, L. SPRUCH, *Statistical Model of Atoms in Intense Magnetic Fields*, Phys. Rev. Lett., **26** (1971), 1136–1139.
- [36] E. TELLER, *On the stability of molecules in the Thomas-Fermi theory*, Rev. Mod. Phys., **34** (1962), 627–631.
- [37] A.Z. MSEZANE, Z. FELFLI, D. SOKOLOVSKI, *Near-threshold resonances in electron elastic scattering cross sections for Au and Pt atoms: identification of electron affinities*, J. Phys. B: At. Mol. Opt. Phys., **41** (2008), no. 10, Article ID 105201, (6pp).
- [38] D. O'REGAN, THEORY OF SINGULAR BOUNDARY VALUE PROBLEMS. World Scientific Publishing Co., Inc., River Edge, NJ, 1994.
- [39] A.L.A. PENNA, J.B. DINIZ, F.A. OLIVERIA, *Charge degeneracy removal in the screened hydrogen atom*, J. Phys. B: At. Mol. Opt. Phys., **41** (2008), no. 19, Article ID 195302, (8pp).
- [40] M.H. PROTTER, H.F. WEINBERGER, *Maximum principles in differential equations*. Corrected reprint of the 1967 original. Springer-Verlag, New York, 1984.
- [41] L.H. THOMAS, *The calculation of atomic fields*, Proc. Cambridge Phil. Soc., **23** (1927), 542.
- [42] C. TISDELL, *What is a Lipschitz condition?* Available on Dr Chris Tisdell's YouTube Channel, 14/02/2014, Accessed on 04/11/2014 <http://www.youtube.com/watch?v=Cnc83B3C2pY>
- [43] C. TISDELL, *Banach fixed point theorem and differential equations*. Available on Dr Chris Tisdell's YouTube Channel, 08/09/2014, Accessed on 04/11/2014 <http://www.youtube.com/watch?v=RhAjh8QpkDY>
- [44] E. ZEIDLER, *Nonlinear functional analysis and its applications. I. Fixed-point theorems*. Translated from the German by Peter R. Wadsack. Springer-Verlag, New York, 1986.