

EXISTENCE OF SOLUTIONS FOR A COUPLED SYSTEM OF THREE-POINT FRACTIONAL ORDER BOUNDARY VALUE PROBLEMS

K.R. PRASAD AND B.M.B. KRUSHNA

(Communicated by Meiqiang Feng)

Abstract. In this paper, we establish the existence of a solution for a coupled system of three-point fractional order boundary value problems by applying the Schauder fixed point theorem in a Banach space under suitable conditions.

1. Introduction

This paper is concerned with establishing the existence of a solution to the coupled system of three-point fractional order boundary value problems,

$$D_{0+}^{\alpha} u(t) = f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)), \quad t \in (0, 1), \quad (1)$$

$$D_{0+}^{\beta} v(t) = f_2(t, u(t), D_{0+}^{q_1} u(t), D_{0+}^{q_2} u(t)), \quad t \in (0, 1), \quad (2)$$

$$u(0) = u'(0) = 0, \quad u'(1) - \xi u'(\eta) = 0, \quad (3)$$

$$v(0) = v'(0) = 0, \quad v'(1) - \xi v'(\eta) = 0, \quad (4)$$

where $2 < \alpha, \beta \leq 3$, $\eta \in (0, 1)$, $\xi \in \mathbb{R}$, $f_1, f_2 : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ are given functions and $D_{0+}^{\alpha}, D_{0+}^{\beta}, D_{0+}^{p_i}, D_{0+}^{q_i}$, $i = 1, 2$ are the standard Riemann–Liouville fractional order derivatives.

The study of fractional order differential equations has emerged as an important area of mathematics. It has wide range of applications in various fields of science and engineering such as physics, mechanics, control systems, flow in porous media, electromagnetics and viscoelasticity. For some of the recent developments in fractional calculus, we refer to Miller and Ross [13], Podlubny [15], Kilbas, Srivastava and Trujillo [7], Kilbas and Trujillo [8], Lakshmikantham and Vatsala [11] and the references therein.

Recently, much interest has been created in establishing the existence of solutions for two-point and multi-point fractional order boundary value problems. To mention the related papers along these lines, we refer to Bai and Lü [2], Kauffman and Mboumi

Mathematics subject classification (2010): 34A34, 34B15.

Keywords and phrases: Coupled system, Green's function, fractional derivative, boundary value problem.

[6], Benchohra, Henderson, Ntouyas and Ouahab [3], Su and Zhang [19] for fractional order differential equations, Su [18], Khan, Rehman and Henderson [9], Prasad and Krushna [16] for systems of fractional order differential equations.

Ahmad and Nieto [1] obtained an existence result for a coupled system of nonlinear fractional order boundary value problem,

$$\begin{aligned} D^\alpha u(t) &= f(t, v(t), D^p v(t)), \quad t \in (0, 1), \\ D^\beta v(t) &= g(t, u(t), D^q u(t)), \quad t \in (0, 1), \\ u(0) = 0, u(1) &= \gamma u(\eta), \quad v(0) = 0, v(1) = \gamma v(\eta), \end{aligned}$$

where D is the standard Riemann–Liouville fractional order derivative, $1 < \alpha, \beta < 2$, p, q, η, γ satisfy certain conditions, by applying the Schauder fixed point theorem.

In [10], the authors studied the existence and uniqueness of solutions to the boundary value problem,

$$\begin{aligned} D^\alpha u(t) &= f(t, u(t), u'(t)), \quad t \in (0, 1), \\ u(0) = 0, D^p u(1) &= \delta D^p u(\eta), \end{aligned}$$

where D is the Caputo fractional order derivative, $1 < \alpha \leq 2$, $0 < \delta < p < 1$, $0 < \eta \leq 1$, by using some fixed point theorem.

We assume that the following conditions hold throughout the paper:

- (A1) $f_1, f_2 : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ are continuous,
- (A2) $\alpha - q_i \geq 1, \beta - p_i \geq 1$, for $i = 1, 2$,
- (A3) $\xi \eta^{\alpha-2} < 1, \xi \eta^{\beta-2} < 1, \eta \in (0, 1)$ and $\xi \in \mathbb{R}$,
- (A4) there exists a nonnegative function $l(t) \in C(0, 1)$ such that $|f_1(t, x, y, z)| \leq l(t) + \varepsilon_1 |x|^{\theta_1} + \varepsilon_2 |y|^{\theta_2} + \varepsilon_3 |z|^{\theta_3}$, where $\varepsilon_1, \varepsilon_2, \varepsilon_3 > 0$ and $0 < \theta_1, \theta_2, \theta_3 < 1$,
- (A5) there exists a nonnegative function $l^*(t) \in C(0, 1)$ such that $|f_2(t, x, y, z)| \leq l^*(t) + \delta_1 |x|^{\psi_1} + \delta_2 |y|^{\psi_2} + \delta_3 |z|^{\psi_3}$, where $\delta_1, \delta_2, \delta_3 > 0$ and $0 < \psi_1, \psi_2, \psi_3 < 1$.

1.1. Preliminaries

In this section, we present some definitions and lemmas that are useful in the proof of our main results.

DEFINITION 1. [15] The Riemann–Liouville fractional integral of order $p > 0$ of a function $f : [0, +\infty) \rightarrow \mathbb{R}$ is given by

$$I_{0+}^p f(t) = \frac{1}{\Gamma(p)} \int_0^t (t - \tau)^{p-1} f(\tau) d\tau,$$

provided the right-hand side is defined.

DEFINITION 2. [15] The Riemann–Liouville fractional derivative of order $p > 0$ of a function $f : [0, +\infty) \rightarrow \mathbb{R}$ is given by

$$D_{0+}^p f(t) = \frac{1}{\Gamma(k-p)} \frac{d^k}{dt^k} \int_0^t (t-\tau)^{k-p-1} f(\tau) d\tau, \quad (k-1 \leq p < k),$$

provided the right-hand side is defined.

LEMMA 1. [15] If $f(t) \in L_1(0, T)$, then the equation

$$D_{0+}^{\sigma_n} y(t) = f(t)$$

has the unique solution $y(t) \in L_1(0, T)$, which satisfies the initial conditions

$$\left[D_{0+}^{\sigma_k} y(t) \right]_{t=0} = b_k, \quad k = 1, 2, \dots, n,$$

is

$$y(t) = \frac{1}{\Gamma(\sigma_n)} \int_0^t (t-\tau)^{\sigma_n-1} f(\tau) d\tau + \sum_{i=1}^n \frac{b_i}{\Gamma(\sigma_i)} t^{\sigma_i-1}.$$

The rest of the paper is organized as follows. In Section 2, the Green’s function for the homogeneous fractional order boundary value problem is constructed and sufficient lemmas are estimated. In Section 3, the existence of a solution for a coupled system of fractional order boundary value problem (1)-(4) is established using the Schauder fixed point theorem. In Section 4, as an application, the results are demonstrated with an example.

2. Green’s Function and Lemmas

In this section, the Green’s function for the homogeneous fractional order boundary value problem is constructed and sufficient lemmas are estimated, which are needed to establish the main results.

Let

$$X = \left\{ u : u, D_{0+}^{q_1} u, D_{0+}^{q_2} u \in C[0, 1] \right\}$$

be the Banach space equipped with the norm,

$$\|u\|_X = \max_{t \in I} |u(t)| + \max_{t \in I} \left| D_{0+}^{q_1} u(t) \right| + \max_{t \in I} \left| D_{0+}^{q_2} u(t) \right|,$$

and

$$Y = \left\{ v : v, D_{0+}^{p_1} v, D_{0+}^{p_2} v \in C[0, 1] \right\}$$

be the Banach space equipped with the norm,

$$\|v\|_Y = \max_{t \in I} |v(t)| + \max_{t \in I} \left| D_{0+}^{p_1} v(t) \right| + \max_{t \in I} \left| D_{0+}^{p_2} v(t) \right|.$$

LEMMA 2. Let $d = (1 - \xi \eta^{\alpha-2})\Gamma(\alpha) \neq 0$. If $h(t) \in C[0, 1]$, then the fractional order differential equation,

$$D_{0+}^{\alpha} u(t) = h(t), \quad t \in (0, 1), \tag{5}$$

with (3) has a unique solution,

$$u(t) = \int_0^1 G_1(t, s)h(s)ds,$$

where $G_1(t, s)$ is the Green's function for the boundary value problem (5),(3) and is given by

$$G_1(t, s) = \begin{cases} G_1(t, s) = \begin{cases} G_{11}(t, s), & 0 < s < t \leq \eta < 1, \\ G_{12}(t, s), & 0 \leq t < s < \eta < 1, \\ G_{13}(t, s), & 0 \leq t < \eta < s < 1, \end{cases} \\ G_1(t, s) = \begin{cases} G_{14}(t, s), & 0 < \eta < s < t \leq 1, \\ G_{15}(t, s), & 0 < \eta \leq t < s < 1, \\ G_{16}(t, s), & 0 \leq s < \eta < t < 1, \end{cases} \end{cases} \tag{6}$$

$$G_{11}(t, s) = \frac{1}{d} \left[(t-s)^{\alpha-1} (1 - \xi \eta^{\alpha-2}) - t^{\alpha-1} (1-s)^{\alpha-2} + \xi t^{\alpha-1} (\eta-s)^{\alpha-2} \right],$$

$$G_{12}(t, s) = \frac{1}{d} \left[\xi t^{\alpha-1} (\eta-s)^{\alpha-2} - t^{\alpha-1} (1-s)^{\alpha-2} \right],$$

$$G_{13}(t, s) = \frac{1}{d} \left[-t^{\alpha-1} (1-s)^{\alpha-2} \right],$$

$$G_{14}(t, s) = \frac{1}{d} \left[(t-s)^{\alpha-1} (1 - \xi \eta^{\alpha-2}) - t^{\alpha-1} (1-s)^{\alpha-2} + \xi t^{\alpha-1} (\eta-s)^{\alpha-2} \right],$$

$$G_{15}(t, s) = \frac{1}{d} \left[-t^{\alpha-1} (1-s)^{\alpha-2} \right],$$

$$G_{16}(t, s) = \frac{1}{d} \left[(t-s)^{\alpha-1} (1 - \xi \eta^{\alpha-2}) - t^{\alpha-1} (1-s)^{\alpha-2} \right].$$

Proof. Let $u(t) \in C[0, 1]$ be the solution of fractional order boundary value problem (5), (3) and is uniquely expressed as

$$\begin{aligned} u(t) &= \frac{1}{d} \left\{ \int_0^t \left[(1 - \xi \eta^{\alpha-2})(t-s)^{\alpha-1} + \left(\xi (\eta-s)^{\alpha-2} - (1-s)^{\alpha-2} \right) t^{\alpha-1} \right] h(s) ds \right. \\ &\quad + \int_t^{\eta} \left[\xi t^{\alpha-1} (\eta-s)^{\alpha-2} - t^{\alpha-1} (1-s)^{\alpha-2} \right] h(s) ds \\ &\quad \left. - \int_{\eta}^1 t^{\alpha-1} (1-s)^{\alpha-2} h(s) ds \right\} \\ &= \int_0^1 G_1(t, s)h(s)ds. \end{aligned}$$

In a similar manner, we construct the Green’s function $G_2(t, s)$ for the homogeneous fractional order boundary value problem corresponding to (2) and (4).

Let $T : X \times Y \rightarrow X \times Y$ be the operator defined as

$$T(u, v)(t) = (T_1 v(t), T_2 u(t)), \tag{7}$$

where

$$T_1 v(t) = \int_0^1 G_1(t, s) f_1(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s)) ds,$$

$$T_2 u(t) = \int_0^1 G_2(t, s) f_2(s, u(s), D_{0+}^{q_1} u(s), D_{0+}^{q_2} u(s)) ds.$$

In view of the continuity of G_1, G_2, f_1 and f_2 , it follows that T is continuous.

Let $B = X \times Y$ be the Banach space defined as

$$B = \{(u, v) : \|(u, v)\| \leq R, t \in I\}, \tag{8}$$

where

$$\|(u, v)\| = \max \{ \|u\|_X, \|v\|_Y \},$$

and

$$R \geq \max \left\{ \left(4\Lambda_1 \varepsilon_1 \right)^{\frac{1}{1-\theta_1}}, \left(4\Lambda_1 \varepsilon_2 \right)^{\frac{1}{1-\theta_2}}, \left(4\Lambda_1 \varepsilon_3 \right)^{\frac{1}{1-\theta_3}}, 4\gamma, \right. \\ \left. \left(4\Lambda_2 \delta_1 \right)^{\frac{1}{1-\psi_1}}, \left(4\Lambda_2 \delta_2 \right)^{\frac{1}{1-\psi_2}}, \left(4\Lambda_2 \delta_3 \right)^{\frac{1}{1-\psi_3}}, 4\nu \right\}.$$

LEMMA 3. [1] *Assume that (A1) is satisfied. Then $(u, v) \in B$ is a solution of the fractional order boundary value problem (1)-(4) if and only if $(u, v) \in B$ is a solution of (7).*

3. Existence of a solution

In this section, we establish the existence of a solution for a coupled system of nonlinear fractional order boundary value problem (1)-(4) by applying the Schauder fixed point theorem.

For our convenience, we denote

$$\Lambda_1 = \frac{1 - \xi \eta^{\alpha-2} + \alpha \xi (\eta^{\alpha-2} - \eta^{\alpha-1})}{(\alpha - 1)(1 - \xi \eta^{\alpha-2})\Gamma(\alpha + 1)} + \frac{(\alpha - 1)(2 - \xi \eta^{\alpha-2}) + \xi \eta^{\alpha-1}(\alpha - q_1)}{(\alpha - 1)(1 - \xi \eta^{\alpha-2})\Gamma(\alpha - q_1 + 1)} \\ + \frac{(\alpha - 1)(2 - \xi \eta^{\alpha-2}) + \xi \eta^{\alpha-1}(\alpha - q_2)}{(\alpha - 1)(1 - \xi \eta^{\alpha-2})\Gamma(\alpha - q_2 + 1)},$$

$$\Lambda_2 = \frac{1 - \xi \eta^{\beta-2} + \alpha \xi (\eta^{\beta-2} - \eta^{\alpha-1})}{(\beta - 1)(1 - \xi \eta^{\beta-2})\Gamma(\beta + 1)} + \frac{(\beta - 1)(2 - \xi \eta^{\beta-2}) + \xi \eta^{\beta-1}(\beta - p_1)}{(\beta - 1)(1 - \xi \eta^{\beta-2})\Gamma(\beta - p_1 + 1)}$$

$$\times \frac{(\beta - 1)(2 - \xi \eta^{\beta-2}) + \xi \eta^{\beta-1}(\beta - p_2)}{(\beta - 1)(1 - \xi \eta^{\beta-2})\Gamma(\beta - p_2 + 1)},$$

$$\begin{aligned} \gamma = & \max_{t \in [0,1]} \int_0^1 |G_1(t,s)l(s)ds| \\ & + \frac{2 - \xi \eta^{\alpha-2}}{1 - \xi \eta^{\alpha-2}} \left\{ \int_0^1 \left[\frac{(1-s)^{\alpha-q_1-1}}{\Gamma(\alpha - q_1)} + \frac{(1-s)^{\alpha-q_2-1}}{\Gamma(\alpha - q_2)} \right] l(s)ds \right\} \\ & + \frac{1}{(1 - \xi \eta^{\alpha-2})} \left[\frac{1}{\Gamma(\alpha - q_1)} + \frac{1}{\Gamma(\alpha - q_2)} \right] \int_0^\eta (\eta - s)^{\alpha-2} l(s)ds, \end{aligned}$$

and

$$\begin{aligned} v = & \max_{t \in [0,1]} \int_0^1 |G_2(t,s)l^*(s)ds| \\ & + \frac{2 - \xi \eta^{\beta-2}}{1 - \xi \eta^{\beta-2}} \left\{ \int_0^1 \frac{(1-s)^{\beta-p_1-1}}{\Gamma(\beta - p_1)} l^*(s)ds + \int_0^1 \frac{(1-s)^{\beta-p_2-1}}{\Gamma(\beta - p_2)} l^*(s)ds \right\} \\ & + \frac{1}{(1 - \xi \eta^{\beta-2})} \left[\frac{1}{\Gamma(\beta - p_1)} + \frac{1}{\Gamma(\beta - p_2)} \right] \int_0^\eta (\eta - s)^{\beta-2} l^*(s)ds. \end{aligned}$$

To establish the existence of a solution to the fractional order boundary value problem (1)-(4) by utilizing the Schauder fixed point theorem.

LEMMA 4. (Schauder fixed point theorem) *Let E be a Banach space and Ω be any nonempty convex and closed subset of E. If M is a continuous mapping of Ω into itself and MΩ is relatively compact, then the mapping M has at least one fixed point.*

THEOREM 1. *Assume that the conditions (A1)-(A5) hold. Then there exists a solution for the coupled system of fractional order boundary value problem (1)-(4).*

Proof. Using the assumptions (A1)-(A5) together with the results

$$D_{0^+}^{q_1} t^{\alpha-1} = \frac{\Gamma(\alpha)}{\Gamma(\alpha - q_1)} t^{\alpha-q_1-1} \text{ and } D_{0^+}^{q_2} t^{\alpha-1} = \frac{\Gamma(\alpha)}{\Gamma(\alpha - q_2)} t^{\alpha-q_2-1}.$$

Firstly, we prove that $T : B \rightarrow B$, where B is given in (8). We have,

$$\begin{aligned} |T_1 v(t)| &= \left| \int_0^1 G_1(t,s) f_1 \left(s, v(s), D_{0^+}^{p_1} v(s), D_{0^+}^{p_2} v(s) \right) ds \right| \\ &\leq \int_0^1 |G_1(t,s)l(s)| ds + \sum_{i=1}^3 \left(\varepsilon_i |R|^{\theta_i} \right) \int_0^1 |G_1(t,s)| ds \\ &\leq \int_0^1 |G_1(t,s)l(s)| ds + \sum_{i=1}^3 \left(\varepsilon_i |R|^{\theta_i} \right) \left[- \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} ds \right] \end{aligned}$$

$$\begin{aligned}
 & + \int_0^1 \frac{t^{\alpha-1}(1-s)^{\alpha-2}}{(1-\xi\eta^{\alpha-2})} ds - \frac{\xi t^{\alpha-1}}{(1-\xi\eta^{\alpha-2})} \int_0^\eta \frac{(\eta-s)^{\alpha-2}}{\Gamma(\alpha)} ds \Big] \\
 \leq & \int_0^1 |G_1(t,s)l(s)| ds + \sum_{i=1}^3 (\varepsilon_i |R|^{\theta_i}) \left[\frac{1-\xi\eta^{\alpha-2} + \alpha\xi(\eta^{\alpha-2} - \eta^{\alpha-1})}{(\alpha-1)\Gamma(\alpha+1)(1-\xi\eta^{\alpha-2})} \right],
 \end{aligned}$$

and

$$\begin{aligned}
 |D_{0+}^{q_1} T_1 v(t)| & = \left| D_{0+}^{q_1} I_{0+}^\alpha f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)) \right. \\
 & \quad - \frac{1}{(1-\xi\eta^{\alpha-2})} \left[I_{0+}^\alpha f_1(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1)) \right. \\
 & \quad \quad \left. - \xi I_{0+}^\alpha f_1(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta)) \right] D_{0+}^{q_1} t^{\alpha-1} \Big| \\
 & = \left| I_{0+}^{\alpha-q_1} f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)) \right. \\
 & \quad - \frac{1}{(1-\xi\eta^{\alpha-2})} \left[I_{0+}^\alpha f_1(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1)) \right. \\
 & \quad \quad \left. - \xi I_{0+}^\alpha f_1(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta)) \right] \frac{\Gamma\alpha}{\Gamma(\alpha-q_1)} t^{\alpha-q_1-1} \Big| \\
 & \leq \frac{1}{\Gamma(\alpha-q_1)(1-\xi\eta^{\alpha-2})} \left[(2-\xi\eta^{\alpha-2}) \int_0^1 (1-s)^{\alpha-q_1-1} l(s) ds \right. \\
 & \quad \left. + \xi \int_0^\eta (\eta-s)^{\alpha-2} l(s) ds \right] + (\varepsilon_1 |R|^{\theta_1} + \varepsilon_2 |R|^{\theta_2} + \varepsilon_3 |R|^{\theta_3}) \\
 & \quad \times \left[\frac{(\alpha-1)(2-\xi\eta^{\alpha-2}) + \xi\eta^{\alpha-1}(\alpha-q_1)}{(\alpha-1)(1-\xi\eta^{\alpha-2})\Gamma(\alpha-q_1+1)} \right]
 \end{aligned}$$

and

$$\begin{aligned}
 |D_{0+}^{q_2} T_1 v(t)| & = \left| D_{0+}^{q_2} I_{0+}^\alpha f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)) \right. \\
 & \quad - \frac{1}{(1-\xi\eta^{\alpha-2})} \left[I_{0+}^\alpha f_1(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1)) \right. \\
 & \quad \quad \left. - \xi I_{0+}^\alpha f_1(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta)) \right] D_{0+}^{q_2} t^{\alpha-1} \Big| \\
 & = \left| I_{0+}^{\alpha-q_2} f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)) \right. \\
 & \quad - \frac{1}{(1-\xi\eta^{\alpha-2})} \left[I_{0+}^\alpha f_1(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1)) \right.
 \end{aligned}$$

$$\begin{aligned}
& -\xi I_{0^+}^{\alpha} f_1\left(\eta, v(\eta), D_{0^+}^{p_1} v(\eta), D_{0^+}^{p_2} v(\eta)\right)\left] \frac{\Gamma \alpha}{\Gamma(\alpha-q_2)} t^{\alpha-q_2-1}\right| \\
\leq & \frac{1}{\Gamma(\alpha-q_2)}\left[\int_0^t(t-s)^{\alpha-q_2-1} l(s) d s+\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right)\right. \\
& \times \int_0^t(t-s)^{\alpha-q_2-1} d s+\frac{1}{(1-\xi \eta^{\alpha-2})}\left\{\int_0^1(1-s)^{\alpha-2} l(s) d s\right. \\
& +\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \int_0^1(1-s)^{\alpha-2} d s \\
& +\xi \int_0^{\eta}(\eta-s)^{\alpha-2} l(s) d s+\xi\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \\
& \times \int_0^1(1-s)^{\alpha-2} d s+\xi \int_0^{\eta}(\eta-s)^{\alpha-2} l(s) d s \\
& \left.+\xi\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \int_0^{\eta}(\eta-s)^{\alpha-2} d s\right\}\left. \right] \\
\leq & \frac{1}{\Gamma(\alpha-q_2)(1-\xi \eta^{\alpha-2})}\left[\left(2-\xi \eta^{\alpha-2}\right) \int_0^1(1-s)^{\alpha-q_2-1} l(s) d s\right. \\
& +\xi \int_0^{\eta}(\eta-s)^{\alpha-2} l(s) d s+\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \\
& \times\left.\left\{\left(2-\xi \eta^{\alpha-2}\right) \int_0^1(1-s)^{\alpha-q_2-1} d s+\xi \int_0^{\eta}(\eta-s)^{\alpha-2} d s\right\}\right] \\
\leq & \frac{1}{\Gamma(\alpha-q_2)(1-\xi \eta^{\alpha-2})}\left[\left(2-\xi \eta^{\alpha-2}\right) \int_0^1(1-s)^{\alpha-q_2-1} l(s) d s\right. \\
& +\xi \int_0^{\eta}(\eta-s)^{\alpha-2} l(s) d s\left. \right]+\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \\
& \times\left[\frac{(\alpha-1)\left(2-\xi \eta^{\alpha-2}\right)+\xi \eta^{\alpha-1}(\alpha-q_2)}{(\alpha-1)(1-\xi \eta^{\alpha-2}) \Gamma(\alpha-q_2+1)}\right].
\end{aligned}$$

Thus,

$$\begin{aligned}
\|T_1 v(t)\|_X & =\max _{t \in[0,1]}|T_1 v(t)|+\max _{t \in[0,1]}\left|D_{0^+}^{q_1} T_1 v(t)\right|+\max _{t \in[0,1]}\left|D_{0^+}^{q_2} T_2 v(t)\right| \\
& \leq \int_0^1\left|G_1(t, s) l(s) d s\right|+\left(\varepsilon_1|R|^{\theta_1}+\varepsilon_2|R|^{\theta_2}+\varepsilon_3|R|^{\theta_3}\right) \\
& \quad \times\left[\frac{1-\xi \eta^{\alpha-2}+\alpha \xi\left(\eta^{\alpha-2}-\eta^{\alpha-1}\right)}{(\alpha-1) \Gamma(\alpha+1)(1-\xi \eta^{\alpha-2})}\right] \\
& \quad +\frac{1}{\Gamma(\alpha-q_1)(1-\xi \eta^{\alpha-2})}\left[\left(2-\xi \eta^{\alpha-2}\right) \int_0^1(1-s)^{\alpha-q_1-1} l(s) d s\right.
\end{aligned}$$

$$\begin{aligned}
 & + \xi \int_0^\eta (\eta - s)^{\alpha-2} l(s) ds \Big] + \left(\varepsilon_1 |R|^{\theta_1} + \varepsilon_2 |R|^{\theta_2} + \varepsilon_3 |R|^{\theta_3} \right) \\
 & \quad \times \left[\frac{(\alpha - 1)(2 - \xi \eta^{\alpha-2}) + \xi \eta^{\alpha-1}(\alpha - q_1)}{(\alpha - 1)(1 - \xi \eta^{\alpha-2})\Gamma(\alpha - q_1 + 1)} \right] \\
 & + \frac{1}{\Gamma(\alpha - q_2)(1 - \xi \eta^{\alpha-2})} \left[(2 - \xi \eta^{\alpha-2}) \int_0^1 (1 - s)^{\alpha - q_2 - 1} l(s) ds \right. \\
 & \quad \left. + \xi \int_0^\eta (\eta - s)^{\alpha-2} l(s) ds \right] + \left(\varepsilon_1 |R|^{\theta_1} + \varepsilon_2 |R|^{\theta_2} + \varepsilon_3 |R|^{\theta_3} \right) \\
 & \quad \times \left[\frac{(\alpha - 1)(2 - \xi \eta^{\alpha-2}) + \xi \eta^{\alpha-1}(\alpha - q_2)}{(\alpha - 1)(1 - \xi \eta^{\alpha-2})\Gamma(\alpha - q_2 + 1)} \right] \\
 & \leq \gamma + \left(\varepsilon_1 |R|^{\theta_1} + \varepsilon_2 |R|^{\theta_2} + \varepsilon_3 |R|^{\theta_3} \right) \Lambda_1 \\
 & \leq \frac{R}{4} + \frac{R}{4} + \frac{R}{4} + \frac{R}{4} \\
 & = R.
 \end{aligned}$$

Similarly, it can be shown that

$$\|T_2 u(t)\|_Y \leq \nu + \left(\delta_1 |R|^{\psi_1} + \delta_2 |R|^{\psi_2} + \delta_3 |R|^{\psi_3} \right) \Lambda_2 \leq R.$$

Hence, we conclude that

$$\|T(u, v)\|_{X \times Y} \leq R.$$

Since $T_1 v(t)$, $T_2 u(t)$, $D_{0+}^{q_1} T_1 v(t)$, $D_{0+}^{p_1} T_2 u(t)$, $D_{0+}^{q_2} T_1 v(t)$ and $D_{0+}^{p_2} T_2 u(t)$ are continuous on $[0, 1]$. Therefore, $T : B \rightarrow B$.

Now, we show that T is a completely continuous operator. For this, we let

$$\begin{aligned}
 M &= \max_{t \in [0, 1]} \left| f_1(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t)) \right|, \\
 N &= \max_{t \in [0, 1]} \left| f_2(t, u(t), D_{0+}^{q_1} u(t), D_{0+}^{q_2} u(t)) \right|.
 \end{aligned}$$

For $(u, v) \in B$, $t, \tau \in [0, 1]$, $t < \tau$, we have

$$\begin{aligned}
 |T_1 v(t) - T_1 v(\tau)| &= \left| \int_0^1 \left(G_1(t, s) - G_1(\tau, s) \right) f_1 \left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s) \right) ds \right| \\
 &\leq \frac{M}{\Gamma \alpha (1 - \xi \eta^{\alpha-2})} \left[\int_0^t [(\tau - s)^{\alpha-1} - (t - s)^{\alpha-1}] (1 - \xi \eta^{\alpha-2}) \right. \\
 &\quad \left. + (\tau^{\alpha-1} - t^{\alpha-1}) (1 - s)^{\alpha-2} + \xi (\tau^{\alpha-1} - t^{\alpha-1}) (\eta - s)^{\alpha-2} \right] ds \\
 &\quad + \int_t^\tau (\tau^{\alpha-1} - t^{\alpha-1}) \left[(1 - s)^{\alpha-2} + \xi (\eta - s)^{\alpha-2} \right] ds
 \end{aligned}$$

$$\begin{aligned}
& + \int_{\eta}^1 (\tau^{\alpha-1} - t^{\alpha-1})(1-s)^{\alpha-2} ds \Big] \\
& \leq M \left[\frac{\tau^{\alpha} - t^{\alpha}}{\Gamma(\alpha+1)} + \frac{(\tau^{\alpha-1} - t^{\alpha-1})(1 - \xi \eta^{\alpha-1})}{(\alpha-1)\Gamma(\alpha)(1 - \xi \eta^{\alpha-2})} \right],
\end{aligned}$$

and

$$\begin{aligned}
& \left| D_{0+}^{q_1} T_1 v(t) - D_{0+}^{q_1} T_1 v(\tau) \right| \\
& = \left| I_{0+}^{\alpha-q_1} f_1 \left(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t) \right) \right. \\
& \quad - \frac{\Gamma \alpha}{\Gamma(\alpha - q_1)(1 - \xi \eta^{\alpha-2})} \left\{ I_{0+}^{\alpha} f_1 \left(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1) \right) \right. \\
& \quad \quad \left. - \xi I_{0+}^{\alpha} f_1 \left(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta) \right) \right\} t^{\alpha-q_1-1} \\
& \quad - I_{0+}^{\alpha-q_1} f_1 \left(\tau, v(\tau), D_{0+}^{p_1} v(\tau), D_{0+}^{p_2} v(\tau) \right) \\
& \quad + \frac{\Gamma \alpha}{\Gamma(\alpha - q_1)(1 - \xi \eta^{\alpha-2})} \left\{ I_{0+}^{\alpha} f_1 \left(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1) \right) \right. \\
& \quad \quad \left. - \xi I_{0+}^{\alpha} f_1 \left(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta) \right) \right\} \tau^{\alpha-q_1-1} \Big| \\
& \leq \frac{1}{\Gamma(\alpha - q_1)} \left[\left| \int_0^t (t-s)^{\alpha-q_1-1} f_1 \left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s) \right) ds \right. \right. \\
& \quad \left. \left. - \int_0^{\tau} (\tau-s)^{\alpha-q_1-1} f_1 \left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s) \right) ds \right| \right. \\
& \quad \left. + \frac{M(\tau^{\alpha-q_1-1} - t^{\alpha-q_1-1})}{\Gamma(\alpha - q_1)(1 - \xi \eta^{\alpha-2})} \left| \left\{ \int_0^1 (1-s)^{\alpha-1} ds - \xi \int_0^{\eta} (\eta-s)^{\alpha-1} ds \right\} \right| \right] \\
& \leq \left[\frac{M(\tau^{\alpha-q_1} - t^{\alpha-q_1})}{\Gamma(\alpha - q_1 + 1)} + \frac{M(1 - \xi \eta^{\alpha})(\tau^{\alpha-q_1-1} - t^{\alpha-q_1-1})}{(1 - \xi \eta^{\alpha-2})\Gamma(\alpha - q_1)\alpha} \right]
\end{aligned}$$

and

$$\begin{aligned}
& \left| D_{0+}^{q_2} T_1 v(t) - D_{0+}^{q_2} T_1 v(\tau) \right| \\
& = \left| I_{0+}^{\alpha-q_2} f_1 \left(t, v(t), D_{0+}^{p_1} v(t), D_{0+}^{p_2} v(t) \right) \right. \\
& \quad - \frac{\Gamma \alpha}{\Gamma(\alpha - q_2)(1 - \xi \eta^{\alpha-2})} \left\{ I_{0+}^{\alpha} f_1 \left(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1) \right) \right. \\
& \quad \quad \left. - \xi I_{0+}^{\alpha} f_1 \left(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta) \right) \right\} t^{\alpha-q_2-1}
\end{aligned}$$

$$\begin{aligned}
 & -I_{0+}^{\alpha-q_2} f_1\left(\tau, v(\tau), D_{0+}^{p_1} v(\tau), D_{0+}^{p_2} v(\tau)\right) \\
 & + \frac{\Gamma \alpha}{\Gamma(\alpha-q_2)(1-\xi \eta^{\alpha-2})} \left\{ I_{0+}^{\alpha} f_1\left(1, v(1), D_{0+}^{p_1} v(1), D_{0+}^{p_2} v(1)\right) \right. \\
 & \quad \left. - \xi I_{0+}^{\alpha} f_1\left(\eta, v(\eta), D_{0+}^{p_1} v(\eta), D_{0+}^{p_2} v(\eta)\right) \right\} \tau^{\alpha-q_2-1} \Big| \\
 \leq & \frac{1}{\Gamma(\alpha-q_2)} \left[\left| \int_0^t (t-s)^{\alpha-q_2-1} f_1\left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s)\right) ds \right. \right. \\
 & \quad \left. \left. - \int_0^{\tau} (\tau-s)^{\alpha-q_2-1} f_1\left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s)\right) ds \right| \right] \\
 & + \frac{M}{\Gamma(\alpha-q_2)(1-\xi \eta^{\alpha-2})} \left| \left(\tau^{\alpha-q_2-1} - t^{\alpha-q_2-1} \right) \right. \\
 & \quad \left. \times \left\{ \int_0^1 (1-s)^{\alpha-1} ds - \xi \int_0^{\eta} (\eta-s)^{\alpha-1} ds \right\} \right| \\
 \leq & \frac{1}{\Gamma(\alpha-q_2)} \left[\left| \int_0^t (t-s)^{\alpha-q_2-1} f_1\left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s)\right) ds \right. \right. \\
 & \quad \left. \left. - \int_0^{\tau} (\tau-s)^{\alpha-q_2-1} f_1\left(s, v(s), D_{0+}^{p_1} v(s), D_{0+}^{p_2} v(s)\right) ds \right| \right] \\
 & + \frac{M(1-\xi \eta^{\alpha})}{\Gamma(\alpha-q_2)(1-\xi \eta^{\alpha-2}) \alpha} \left(\tau^{\alpha-q_2-1} - t^{\alpha-q_2-1} \right) \\
 \leq & \frac{M}{\Gamma(\alpha-q_2)} \left[\int_0^{\tau} (t-s)^{\alpha-q_2-1} - (\tau-s)^{\alpha-q_2-1} ds \right. \\
 & \quad \left. + \int_t^{\tau} (\tau-s)^{\alpha-q_2-1} ds \right] + \frac{M(1-\xi \eta^{\alpha})(\tau^{\alpha-q_2-1} - t^{\alpha-q_2-1})}{(1-\xi \eta^{\alpha-2}) \Gamma(\alpha-q_2) \alpha} \\
 \leq & \left[\frac{M(\tau^{\alpha-q_2} - t^{\alpha-q_2})}{\Gamma(\alpha-q_2+1)} + \frac{M(1-\xi \eta^{\alpha})(\tau^{\alpha-q_2-1} - t^{\alpha-q_2-1})}{(1-\xi \eta^{\alpha-2}) \Gamma(\alpha-q_2) \alpha} \right].
 \end{aligned}$$

Analogously, it can be proved that

$$\left| T_2 u(t) - T_2 u(\tau) \right| \leq N \left[\frac{\tau^{\beta} - t^{\beta}}{\Gamma(\beta+1)} + \frac{(\tau^{\beta-1} - t^{\beta-1})(1-\xi \eta^{\beta-1})}{(\beta-1) \Gamma(\beta)(1-\xi \eta^{\beta-2})} \right],$$

$$\begin{aligned}
 & \left| D_{0+}^{p_1} T_2 u(t) - D_{0+}^{p_1} T_2 u(\tau) \right| \\
 & \leq N \left[\frac{(\tau^{\beta-p_1} - t^{\beta-p_1})}{\Gamma(\beta-p_1+1)} + \frac{(1-\xi \eta^{\beta})(\tau^{\beta-p_1-1} - t^{\beta-p_1-1})}{(1-\xi \eta^{\beta-2}) \Gamma(\beta-p_1) \beta} \right]
 \end{aligned}$$

and

$$\begin{aligned} & \left| D_{0+}^{p_2} T_2 u(t) - D_{0+}^{p_2} T_2 u(\tau) \right| \\ & \leq N \left[\frac{(\tau^{\beta-p_2} - t^{\beta-p_2})}{\Gamma(\beta - p_2 + 1)} + \frac{(1 - \xi \eta^\beta)(\tau^{\beta-p_2-1} - t^{\beta-p_2-1})}{(1 - \xi \eta^{\beta-2})\Gamma(\beta - p_2)\beta} \right]. \end{aligned}$$

Therefore, it follows from the above estimates that T_B is an equicontinuous. Also it is uniformly bounded as $T_B \subset B$. Thus, we conclude that T is a completely continuous operator. Hence by the Schauder fixed point theorem, there exists a solution to the coupled system of fractional order boundary value problem (1)-(4). This completes the proof.

4. An example

In this section, as an application, we demonstrate our results with an example. Consider the system of fractional order differential equations,

$$D_{0+}^{\frac{23}{9}} u(t) = \frac{1}{\sqrt{t(1-t^2)}} \left[\left(D_{0+}^{\frac{3}{2}} v(t) \right)^{\theta_3} + \left(D_{0+}^{\frac{1}{5}} v(t) \right)^{\theta_2} + (v(t))^{\theta_1} \right] + \left. \begin{aligned} & \frac{99}{100}, \quad t \in (0, 1), \end{aligned} \right\} \tag{9}$$

$$D_{0+}^{\frac{16}{7}} v(t) = \frac{1}{\sqrt{t(1-t^2)}} \left[\left(D_{0+}^{\frac{4}{3}} u(t) \right)^{\psi_3} + \left(D_{0+}^{\frac{1}{7}} u(t) \right)^{\psi_2} + (u(t))^{\psi_1} \right] + \left. \begin{aligned} & \frac{1}{2}, \quad t \in (0, 1), \end{aligned} \right\} \tag{10}$$

satisfying the three-point boundary conditions,

$$u(0) = u'(0) = 0, \quad u'(1) - \frac{7}{9}u'\left(\frac{5}{7}\right) = 0, \tag{11}$$

$$v(0) = v'(0) = 0, \quad v'(1) - \frac{7}{9}v'\left(\frac{5}{7}\right) = 0, \tag{12}$$

where $0 < \theta_i, \psi_i < 1, i = 1, 2, 3$. By direct calculations, one can determine

$$\Lambda_1 = 29.285308, \Lambda_2 = 15.356694, \gamma = 0.475942, \nu = 0.811456$$

and

$$\begin{aligned} f_1(t, x, y, z) &= \frac{1}{\sqrt{t(1-t^2)}} \left[(x(t))^{\theta_1} + (y(t))^{\theta_2} + (z(t))^{\theta_3} \right] + \frac{99}{100}, \\ f_2(t, x, y, z) &= \frac{1}{\sqrt{t(1-t^2)}} \left[(x(t))^{\psi_1} + (y(t))^{\psi_2} + (z(t))^{\psi_3} \right] + \frac{1}{2}, \end{aligned}$$

for $t \in (0, 1)$, we have

$$\begin{aligned} |f_1(t, x, y, z)| &\leq l + \varepsilon_1 |x|^{\theta_1} + \varepsilon_2 |y|^{\theta_2} + \varepsilon_3 |z|^{\theta_3}, \\ |f_2(t, x, y, z)| &\leq l^* + \delta_1 |x|^{\psi_1} + \delta_2 |y|^{\psi_2} + \delta_3 |z|^{\psi_3}, \end{aligned}$$

where l, l^* are constants, $\varepsilon_1, \varepsilon_2, \varepsilon_3 > 0$ and $\delta_1, \delta_2, \delta_3 > 0$. Applying the Theorem 1, the fractional order boundary value problem (9)-(12) has a solution $(u, v) \in B$, where B is determined from (8).

Acknowledgements. The authors would like to thank the editor and referees for their valuable comments and suggestions.

REFERENCES

- [1] B. AHMED AND J.J. NIETO, *Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions*, Comput. Math. Appl., **58** (2009), 1838–1843.
- [2] Z. BAI AND H. LÜ, *Positive solutions for boundary value problems of nonlinear fractional differential equations*, J. Math. Anal. Appl., **311** (2005), 495–505.
- [3] M. BENCHOHRA, J. HENDERSON, S. K. NTOYUAS AND A. OUAHAB, *Existence results for fractional order functional differential equations with infinite delay*, J. Math. Anal. Appl., **338** (2008), 1340–1350.
- [4] M. FENG, X. LIU AND H. FENG, *The existence of positive solution to a nonlinear fractional differential equation with integral boundary conditions*, Adv. Difference Equ., **2011** (2011), Article ID 546038, 1–14.
- [5] M. FENG, X. ZHANG AND W. GE, *New existence results for higher-order nonlinear fractional differential equation with integral boundary conditions*, Bound. Value Probl., **2011** (2011), Article ID 720702, 1–20.
- [6] E.R. KAUFFMAN AND E. MBOUMI, *Positive solutions of a boundary value problem for a nonlinear fractional differential equation*, Elec. J. Qual. Theory Diff. Equ., **2008** (2008), 1–11.
- [7] A.A. KILBAS, H.M. SRIVASTHAVA AND J. J. TRUJILLO, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, **204**, Elsevier Science, Amsterdam, 2006.
- [8] A. A. KILBAS AND J. J. TRUJILLO, *Differential equations of fractional order methods, results, problems*, Appl. Anal., **78** (2001), 153–192.
- [9] R.A. KHAN, M. REHMAN AND J. HENDERSON, *Existence and uniqueness of solutions for nonlinear fractional differential equations with integral boundary conditions*, Fract. Differ. Calc., **1** (2011), 29–43.
- [10] R.A. KHAN AND H. KHAN, *Existence of solution for a three-point boundary value problem of fractional differential equation*, Journal of Fractional Calculus and Applications, **5** (2014), 156–164.
- [11] V. LAKSHMIKANTHAM AND A.S. VATSALA, *Basic theory of fractional differential equations*, Nonlinear Anal., **69** (2008), 2677–2682.
- [12] N.I. MAHMUDOV AND S. UNUL, *Existence of solutions of $\alpha \in (2, 3]$ order fractional three-point boundary value problems with integral conditions*, Abst. Appl. Anal., **2014** (2014), Article ID 198632, 1–12.
- [13] K.S. MILLER AND B. ROSS, *An Introduction to Fractional Calculus and Fractional Differential Equations*, John Wiley and Sons, New York, 1993.
- [14] Z. OUYANG AND G. LI, *Existence of the solutions for a class of nonlinear fractional order three-point boundary value problems with resonance*, Bound. Value Probl., **2012** (2012), Article ID 68, 1–13.
- [15] I. PODULBNY, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [16] K. R. PRASAD AND B. M. B. KRUSHNA, *Multiple positive solutions for a coupled system of Riemann–Liouville fractional order two-point boundary value problems*, Nonlinear Stud., **20** (2013), 501–511.

- [17] M. REHMAN AND R.A. KHAN, *A note on boundary value problems for a coupled system of fractional differential equations*, *Comput. Math. Appl.*, **61** (2011), 2630–2637.
- [18] X. SU, *Boundary value problem for a coupled system of non-autonomous fractional differential equations*, *Appl. Math. Lett.*, **22** (2009), 64–69.
- [19] X. SU AND S. ZHANG, *Solutions to boundary value problems for nonlinear differential equations of fractional order*, *Electron. J. Differential Equations*, **2009** (2009), 1–15.
- [20] G. WANG, L. ZHANG AND S.K. NTUYAS, *Multiplicity of positive solutions for fractional order three-point boundary value problems*, *Commun. Appl. Nonlinear Anal.*, **20** (2013), 41–53.
- [21] Y. ZHANG AND Z. BAI, *Existence of solutions for nonlinear fractional three-point boundary value problems at resonance*, *J. Appl. Math. Comput.*, **36** (2011), 417–440.
- [22] Y. ZHANG, Z. BAI AND T. FENG, *Existence results for a coupled system of nonlinear fractional three-point boundary value problems at resonance*, *Comput. Math. Appl.*, **61** (2011), 1032–1047.

(Received November 25, 2014)

(Revised April 19, 2015)

K.R. Prasad
Department of Applied Mathematics
Andhra University
Visakhapatnam, 530 003
India
e-mail: rajendra92@rediffmail.com

B.M.B. Krushna
Department of Mathematics
MVGR College of Engineering
Vizianagaram, 535 005
India
e-mail: muraleebalu@yahoo.com