

A PROBLEM INVOLVING THE p -LAPLACIAN OPERATOR

RATAN KR. GIRI AND D. CHOUDHURI

Abstract. Using a variational technique we guarantee the existence of a solution to the resonant Lane-Emden problem $-\Delta_p u = \lambda |u|^{q-2} u$, $u|_{\partial\Omega} = 0$ if and only if a solution to $-\Delta_p u = \lambda |u|^{q-2} u + f$, $u|_{\partial\Omega} = 0$, $f \in L^{p'}(\Omega)$ (p' being the conjugate of p), exists for $q \in (p, p^*)$ under certain condition on λ , where p^* is the Sobolev conjugate of p .

Mathematics subject classification (2010): 35A15, 35A01.

Keywords and phrases: p -Laplacian, elliptic PDE, Palais-Smale condition, Sobolev space.

REFERENCES

- [1] A. AMBROSETTI AND P. H. RABINOWITZ, *Dual Variational Methods in Critical Point Theory and Applications*, J. Funct. Anal., **14** (1973), 349–381.
- [2] B. KAWOHL, *Symmetry results for functions yielding best constants in Sobolev-type inequalities*, Discrete Contin. Dynam. Systems, **6** (2000), 683–690.
- [3] C. GRUMIAU AND E. PARINI, *On the asymptotics of solutions of the Lane-Emden problem for the p -Laplacian*, Archiv der Mathematik, **91**, 4 (2008), 354–365.
- [4] G. BACHMAN AND L. NARICI, *Functional Analysis*, Dover Publications, Mineola, New York, 1966.
- [5] GREY ERCOLE, *On the resonant Lane-Emden problem for the p -Laplacian*, Communications in Contemporary Mathematics, **16**, 4 (2014), 1350033-1–22.
- [6] GREY ERCOLE, *On the resonant Lane-Emden problem for the p -Laplacian*, Communications in Contemporary Mathematics, **16**, 4 (2014), 1492001-1.
- [7] IDRISSE LY, *The first eigenvalue for the p -laplacian operator*, Journal of inequalities of pure and applied Math., **6**, 3 (2005), 1–28.
- [8] J. GARCÍA AZORERO AND I. PERAL ALONSO, *On limits of solutions of elliptic problems with nearly critical exponent*, Comm. Partial Differential Equations, **17** (1992), 2113–2126.
- [9] J. I. DÍAZ AND J. E. SAA, *Existence et unicité de solutions positives pour certaines équations elliptiques quasilinéaires*, C. R. Acad. Sci. Paris, **305** (1987), 521–524.
- [10] S. KESAVAN, *Functional Analysis and applications*, New age international pvt. ltd., 2003.
- [11] L.C. EVANS, *Partial Differential Equations*, Amer. Math. Soc., 2009.
- [12] M. F. BIDAUT-VÉRON, N. Q. HUNG AND LAURENT VÉRON, *Quasilinear Lane-Emden equations with absorption and measure data*, J. Math. Pures Appl., **102**, 2 (2014), 315–337.
- [13] M. ÔTANI, *Existence and nonexistence of nontrivial solutions of some nonlinear degenerate elliptic equations*, J. Funct. Anal., **76** (1988), 140–159.
- [14] N. DUNFORD AND J. SCHWARTZ, *Linear Operators. Part 2: Spectral theory*, Wiley (Interscience), New York, 1958.
- [15] NIKOS E. MASTORAKIS, HASSAN FATHABADI, *On the solution of p -Laplacian for non-newtonian fluid flow*, W.S.E.A.S. transactions on Math., **8**, 6 (2009), 238–245.
- [16] P. DRÁBEK, *A note on the nonuniqueness for some quasilinear eigenvalue problem*, Appl. Math. Lett., **13** (2000), 39–41.
- [17] PHILIPPE G. CIARLET, *The finite element method for elliptic problems*, Studies in Mathematics and its Applications, Vol. 4, North-Holland Publishing Co., Amsterdam, 1978.
- [18] P. JUNTINEN, P. LINDQVIST, *On the higher eigenvalues for the ∞ -eigenvalue problem*, Calc. Var. partial differential equations, **23** (2005), 169–192.

- [19] RATAN KR. GIRI AND D. CHOUDHURI, Reduced limit approach to semilinear elliptic PDEs with measure data (arXiv:1605.00870 [math.AP]).
- [20] T. BHATTACHARYA, E. D. BENEDETTO AND J. MANFREDI, *Limits as $p \rightarrow \infty$ of $\Delta_p u = f$ and related extremal problems*, Rend. Sem. Mat. Univ. Politec. Torino, special issue (1991), 15–68.