

## A REMARK ON THE LOCAL WELL-POSEDNESS FOR A COUPLED SYSTEM OF MKDV TYPE EQUATIONS IN $H^s \times H^k$

XAVIER CARVAJAL

*Abstract.* We consider the initial value problem associated to a system consisting modified Korteweg-de Vries type equations

$$\begin{cases} \partial_t v + \partial_x^3 v + \partial_x(vw^2) = 0, & v(x, 0) = \phi(x), \\ \partial_t w + \alpha \partial_x^3 w + \partial_x(v^2 w) = 0, & w(x, 0) = \psi(x), \end{cases}$$

and using only bilinear estimates of the type  $\|J^\gamma F_{b_1}^1 \cdot J^\beta F_{b_2}^2\|_{L_x^2 L_t^2}$ , where  $J$  is the Bessel potential and  $F_{b_j}^j$ ,  $j = 1, 2$  are multiplication operators, we prove the local well-posedness results for given data in low regularity Sobolev spaces  $H^s(\mathbb{R}) \times H^k(\mathbb{R})$  for  $\alpha \neq 0, 1$ . In this work we improve the previous result in [6], extending the LWP region from  $|s - k| < 1/2$  to  $|s - k| < 1$ . This result is sharp in the region of the LWP with  $s \leq 0$  and  $k \leq 0$ , in the sense of the trilinear estimates fails to hold.

*Mathematics subject classification (2010):* 35Q35, 35Q53.

*Keywords and phrases:* Korteweg-de Vries equation, Cauchy problem, local well-posedness.

### REFERENCES

- [1] M. ABLOWITZ, D. KAUP, A. NEWELL AND H. SEGUR, *Nonlinear evolution equations of physical significance*, Phys. Rev. Lett. **31** (2) (1973) 125–127.
- [2] E. ALARCON, J. ANGULO AND J. F. MONTENEGRO, *Stability and instability of solitary waves for a nonlinear dispersive system*, Nonlinear Analysis **36** (1999) 1015–1035.
- [3] B. ALVAREZ, X. CARVAJAL, *On the local well-posedness for some systems of coupled KdV equations*, Nonlinear Analysis **69**, (2008) 692–715.
- [4] D. BEKIRANOV, T. OGAWA, G. PONCE, *Weak solvability and well-posedness of a coupled Schrödinger-Korteweg de Vries equation for capillary-gravity wave interactions*, Proceedings of the American Mathematical Society **125** (1997) 2907–2919.
- [5] J. BOURGAIN, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, II, The KdV-equation*, Geom. Funct. Anal. **3** (1993) 209–262.
- [6] X. CARVAJAL, L. ESQUIVEL, R. SANTOS, *Sharp well-posedness for a coupled system of mkdv type equations*, (2020) arXiv:2003.12619v1.
- [7] X. CARVAJAL, M. PANTHEE, *Sharp well-posedness for a coupled system of mkdv type equations*, J. Evol. Equ. **19**, (2019), 1167–1197.
- [8] X. CARVAJAL, *Sharp global well-posedness for a higher order Schrödinger equation*, J. Fourier Anal. Appl. **12** (2006) no. 1, 53–70.
- [9] X. CARVAJAL, *Local well-posedness for a higher order nonlinear Schrödinger equation in Sobolev spaces of negative indices*, Electron. J. Differential Equations **13** (2004) 1–13.
- [10] M. CHEN, B. GUO, *Local Well and Ill Posedness for the Modified KdV Equations in Subcritical Modulation Spaces*, (2018) arXiv:1811.05182.
- [11] M. CHRIST, J. HOLMER, D. TATARU, *Low regularity a priori bounds for the modified Korteweg-de Vries equation*, Libertas Mathematica **32** (2012) no. 1 51–75.
- [12] A. CORCHO, M. PANTHEE, *Global well-posedness for a coupled modified KdV system*, Bull Braz Math Soc, New Series **43**, (2012) 27–57.

- [13] J. GINIBRE, Y. TSUTSUMI, G. VELO, *On the Cauchy problem for the Zakharov system*, J. Funct. Anal. **151** (1997), no. 2, 384–436.
- [14] Z. GUO, *Global well-posedness of Korteweg-de Vries equation in  $H^{-3/4}(\mathbb{R})$* , J. Math. Pures Appl. **91** (2009) 583–597.
- [15] T. OH, *Diophantine conditions in well-posedness theory of coupled KdV-type systems: local theory*, Int. Math. Res. Not. IMRN (2009), no. 18, 3516–3556.
- [16] C. E. KENIG, G. PONCE, L. VEGA, *A bilinear estimate with applications to the KdV equation*, J. Amer. Math. Soc. **92** (1996) 573–603.
- [17] C. E. KENIG, G. PONCE, L. VEGA, *Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle*, Comm. Pure Appl. Math. **46** (1993) 527–620.
- [18] C. E. KENIG, G. PONCE, L. VEGA, *The Cauchy problem for the Korteweg-de Vries equation in Sobolev spaces of negative indices*, Duke Math. J. **71** (1993), no. 1, 1–21.
- [19] R. KILLIP, M. VISAN, *KdV is well-posed in  $H^{-1}$* , Ann. of Math. **190**, no. 1 (2019) 249–305.
- [20] NOBU KISHIMOTO, *Well-posedness of the Cauchy problem for the Korteweg-de Vries equation at the critical regularity*, Differential Integral Equations **22** (2009), no. 5–6, 447–464.
- [21] T. TAO, *Multilinear weighted convolution of  $L^2$ -functions, and applications to nonlinear dispersive equations*, Amer. J. Math. **123** (2001) no. 5, 839–908.
- [22] J. F. MONTENEGRO, *Sistemas de equações de evolução não lineares; Estudo local, global e estabilidade de ondas solitárias*, Ph. D. Thesis, IMPA, Rio de Janeiro (1995).
- [23] H. TAKAOKA, *Well-posedness for the higher order nonlinear Schrödinger equation*, Adv. Math. Sci. Appl. **10** (2000) 149–171.