

EXISTENCE OF SOLUTIONS FOR A COUPLED SYSTEM OF CAPUTO TYPE FRACTIONAL-ORDER DIFFERENTIAL INCLUSIONS WITH NON-SEPARATED BOUNDARY CONDITIONS ON MULTIVALUED MAPS

B. M. B. KRUSHNA*, K. R. PRASAD AND P. VEERAIAH

(Communicated by C. Goodrich)

Abstract. Sufficient conditions for the existence of solutions to a coupled system of fractional-order differential inclusions associated with fractional non-separated boundary conditions for multivalued maps are established, by employing the nonlinear alternative of Leray–Schauder type. We emphasize that the methods of fixed point theory used in our analysis are standard, although their application to a system of fractional-order differential inclusions is new.

1. Introduction

Fractional-order differential equations (FDEs) arise in the mathematical modeling of several real-world phenomena and have now attained immense value due to their broad applications in the fields of engineering and science. Recently, there has been shown a great deal of interest in the study of fractional-order boundary value problems (FBVPs). Fractional-order models are more realistic and beneficial than their corresponding integer-order counterparts. See, for example, [18, 26] for additional information and explanations.

The existence of solutions for FDEs has been studied and used by authors under different initial or boundary conditions. For the recent development on the topic, we refer the reader to the works [15, 27, 28, 21, 35, 22] and the references cited therein. In particular, coupled systems of FDEs have aroused significant attention in the mathematical modeling of physical phenomena such as disease models [12, 25], chaos synchronization [14, 34], ecological effects [17], anomalous diffusion [32], etc. For some prior theoretical findings on coupled systems of FDEs [31, 33].

Differential inclusions play an essential role as a tool in the analysis of various dynamical processes such as sweeping processes [2, 24, 29], nonlinear dynamics of wheeled vehicles [6], control problems [20], granular systems [30], etc. Information on the critical issues in stochastic processes, control, differential games, optimization and their application to the economy, manufacturing, queuing and climate control can be found in [19] and also we refer [11] for application of fractional-order differential

Mathematics subject classification (2010): 34A08, 34A60, 34B15.

Keywords and phrases: Fractional derivative, differential inclusions, boundary value problem, multi-valued maps, system.

* Corresponding author.

inclusions in synchronization processes. In [5], the existence of solutions for the BVP of coupled fractional-order differential inclusions supplemented by coupled boundary conditions studied.

To the best of our knowledge, the fractional-order coupled system with nonlocal and non-separated boundary conditions has not yet been studied thoroughly. We emphasize here that our problem is new in the sense of implemented the non-separated coupled boundary conditions. The objective of this work is to develop criteria for the existence of solutions for a coupled system of fractional-order differential inclusions associated with coupled boundary conditions for multi-valued maps F and G that include convex and non-convex values by using the non-linear alternative of Leray–Schauder type. In this article, we consider a coupled system of fractional-order differential inclusions

$$\begin{cases} {}^cD_{0+}^{q_1}y_1(t) \in F(t, y_1(t), y_2(t)), & 0 < t < 1, \\ {}^cD_{0+}^{q_2}y_2(t) \in G(t, y_1(t), y_2(t)), & 0 < t < 1, \end{cases} \quad (1)$$

subject to the fractional non-separated boundary conditions

$$\begin{cases} y_1(0) = \vartheta_{11}y_2(1), & {}^cD_{0+}^{q_3}y_1(1) = \vartheta_{12}{}^cD_{0+}^{q_3}y_2(\eta), \\ y_2(0) = \vartheta_{21}y_1(1), & {}^cD_{0+}^{q_3}y_2(1) = \vartheta_{22}{}^cD_{0+}^{q_3}y_1(\eta), \end{cases} \quad (2)$$

where $q_1, q_2 \in (1, 2]$, $0 < \eta, q_3 < 1$, ${}^cD^{q_i}$, for $i = 1, 2, 3$ are the Caputo type fractional-order derivatives and $\vartheta_{11}, \vartheta_{12}, \vartheta_{21}$ and ϑ_{22} are real constants.

Throughout the article, we assume the following conditions hold:

- (A0) $\Delta = 1 - \vartheta_{12}\vartheta_{22}\eta^{2(1-q_3)} \neq 0$ and $\vartheta_{21}\vartheta_{11} \neq 1$,
- (A1) $F, G : [0, 1] \times \mathbb{R}^2 \rightarrow \mathcal{P}(\mathbb{R})$ are given multivalued maps, $\mathcal{P}(\mathbb{R})$ is the family of all nonempty subsets of \mathbb{R} ,
- (A2) the maps $F, G : [0, 1] \times \mathbb{R}^2 \rightarrow \mathcal{P}(\mathbb{R})$ are L^1 –Carathéodory and have convex values,
- (A3) there exist continuous functions $b_1, b_2 : [0, 1] \rightarrow \mathbb{R}^+$ and continuous nondecreasing functions $\Psi_1, \Psi_2, \Phi_1, \Phi_2 : [0, \infty) \rightarrow \mathbb{R}^+$ such that

$$\|F(t, y_1, y_2)\|_{\mathcal{P}} = \sup \left\{ |f| : f \in F(t, y_1, y_2) \right\} \leq b_1(t) (\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|)),$$

$$\|G(t, y_1, y_2)\|_{\mathcal{P}} = \sup \left\{ |g| : g \in G(t, y_1, y_2) \right\} \leq b_2(t) (\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|)),$$

for each $(t, y_1, y_2) \in [0, 1] \times \mathbb{R}^2$,

- (A4) there exist positive constants m_1, m_2 and N such that

$$\frac{m_1 \|b_1\| (\Psi_1(N) + \Phi_1(N)) + m_2 \|b_2\| (\Psi_2(N) + \Phi_2(N))}{N} < 1.$$

The rest of this article is organized as follows. Section 2 consists some preliminary results. The main results are presented in Section 3. In Section 4, as an application, we demonstrate our results with an example.

2. Preliminaries

We recall some basic definitions of fractional calculus.

DEFINITION 1. [26] The fractional integral of a function $f : [a, +\infty) \rightarrow \mathbb{R}$ is given by

$${}_a D_t^{-p} f(t) = {}_a I_t^p f(t) = \frac{1}{\Gamma(p)} \int_a^t (t-\tau)^{p-1} f(\tau) d\tau,$$

provided the right-hand side is defined and $p > 0$.

DEFINITION 2. The Caputo derivative of a function $f : [0, \infty) \rightarrow \mathbb{R}$ can be expressed as

$${}^c D_{0+}^m f(t) = D_{0+}^m \left[f(t) - \sum_{j=0}^{n-1} \frac{t^j}{j!} f^{(j)}(0) \right], \quad t > 0, \quad n-1 < m < n.$$

For the sake of convenience, we will use ${}^c D^m$ instead of ${}^c D_{0+}^m$ in the rest of this paper.

REMARK 1. If $f(t) \in C^n[0, \infty)$, then

$${}^c D^m f(t) = \frac{1}{\Gamma(n-m)} \int_0^t (t-\tau)^{n-m-1} f^{(n)}(\tau) d\tau, \quad t > 0, \quad n-1 < m < n.$$

Some basic concepts of multivalued analysis are listed here [1, 10, 16]. Let $(X, \|\cdot\|)$ be a Banach space and K be a subset of X . Then

- (i) $\mathcal{P}_{cl}(X) = \{K \subset \mathcal{P}(X) : K \text{ is closed}\},$
- (ii) $\mathcal{P}_{cv}(X) = \{K \subset \mathcal{P}(X) : K \text{ is convex}\},$
- (iii) $\mathcal{P}_b(X) = \{K \subset \mathcal{P}(X) : K \text{ is bounded}\},$
- (iv) $\mathcal{P}_{cp}(X) = \{K \subset \mathcal{P}(X) : K \text{ is compact}\},$
- (v) $\mathcal{P}_{cv, cp}(X) = \mathcal{P}_{cv}(X) \cap \mathcal{P}_{cp}(X).$

A multivalued map $G : X \rightarrow \mathcal{P}(X)$ is

- (a) convex (closed) valued if $G(x)$ is convex (closed) for all $x \in X$,
- (b) upper semicontinuous (u.s.c.) on X if the set $G(x_0)$ is a nonempty, closed subset of X for each open set U of X containing $G(x_0)$ is a nonempty, \exists an open neighborhood V of x_0 such that $G(V) \subset U$.
- (c) lower semicontinuous (l.s.c.) if the set $\{x \in X : G(x) \cap V \neq \emptyset\}$ is open for any open set V in X .
- (d) completely continuous if $G(\mathcal{B})$ is relatively compact for every $\mathcal{B} \in \mathcal{P}_b(X)$.

REMARK 2. If the multivalued map G is completely continuous with nonempty compact values then G is u.s.c. $\iff G$ has a closed graph, i.e., $x_n \rightarrow x_*$, $y_n \rightarrow y_*$, $y_n \in G(x_n)$ imply $y_* \in G(x_*)$. The set $Gr(G) = \{(x, y) \in X \times Y, y \in G(x)\}$ defines the graph of G .

DEFINITION 3. A multivalued map $G : [c, d] \rightarrow \mathcal{P}_{cl}(\mathbb{R})$ is said to be measurable if the function $t \mapsto d(y_1, G(t)) = \inf \{|y_1 - y_2| : y_2 \in G(t)\}$ is measurable for every $y_1 \in \mathbb{R}$.

DEFINITION 4. A multivalued map $G : [c, d] \times \mathbb{R}^2 \rightarrow \mathcal{P}(\mathbb{R})$ is said to be Carathéodory if

- (i) $t \mapsto G(t, x, y)$ is measurable $\forall x, y \in \mathbb{R}$,
- (ii) $(x, y) \mapsto G(t, x, y)$ is u.s.c. for almost all $t \in [c, d]$.

DEFINITION 5. A Carathéodory function G is said to be L^1 -Carathéodory if for each $\rho > 0$, $\exists \varphi_\rho \in L^1([c, d], \mathbb{R}^+)$ such that $\|G(t, x, y)\|_{\mathcal{P}} = \sup \{|g| : g \in G(t, x, y)\} \leq \varphi_\rho(t) \quad \forall x, y \in \mathbb{R}$ with $\|x\|, \|y\| \leq \rho$ and for a.e. $t \in [c, d]$.

LEMMA 1. ([10, Proposition 1.2]) *If $G : X \rightarrow \mathcal{P}_{cl}(Y)$ is u.s.c., then $Gr(G)$ is a closed subset of $X \times Y$, i.e., for every sequence $\{x_n\}_{n \in \mathbb{N}} \subset X$ and $\{y_n\}_{n \in \mathbb{N}} \subset Y$, if $x_n \rightarrow x_*$, $y_n \rightarrow y_*$ when $n \rightarrow \infty$ and $y_n \in G(x_n)$, then $y_* \in G(x_*)$. Conversely, if G is completely continuous and has a closed graph, then it is upper semicontinuous.*

LEMMA 2. ([23]) *Let X be a separable Banach space. Let $G : [0, 1] \times \mathbb{R}^2 \rightarrow \mathcal{P}_{cp, cv}(\mathbb{R})$ be an L^1 -Carathéodory multivalued map and let χ be a linear continuous mapping from $L^1([0, 1], \mathbb{R})$ to $C([0, 1], \mathbb{R})$. Then the operator*

$$\chi \circ S_{G,x} : C([0, 1], \mathbb{R}) \rightarrow \mathcal{P}_{cp, cv}(C([0, 1], \mathbb{R})), \quad x \mapsto (\chi \circ S_{G,x})(x) = \chi(S_{G,x})$$

is a closed graph operator in $C([0, 1], \mathbb{R}) \times C([0, 1], \mathbb{R})$.

LEMMA 3. (Nonlinear alternative for Kakutani maps [13]) *Let \mathcal{E}_1 be a closed convex subset of a Banach space \mathcal{E} , and \mathcal{U} be an open subset of \mathcal{E}_1 with $0 \in \mathcal{U}$. Suppose that $F : \overline{\mathcal{U}} \rightarrow \mathcal{P}_{cp, cv}(\mathcal{E}_1)$ is an upper semicontinuous compact map. Then either*

- (i) F has a fixed point in $\overline{\mathcal{U}}$, or
- (ii) there is a $u \in \partial \mathcal{U}$ and $\lambda \in (0, 1)$ with $u \in \lambda F(u)$.

We now present an auxiliary lemma, which will play a key role in the forthcoming analysis.

LEMMA 4. *Let $\phi, \psi \in C([0, 1], \mathbb{R})$ and Suppose that the condition (A0) is fulfilled. Then the solution of the associated linear FDEs*

$$\begin{cases} {}^cD_{0+}^{q_1} y_1(t) = \phi(t), & t \in [0, 1], \\ {}^cD_{0+}^{q_2} y_2(t) = \psi(t), & t \in [0, 1], \end{cases} \quad (3)$$

supplemented with the boundary conditions (2) is given by

$$\left. \begin{aligned} y_1(t) = & \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] (\vartheta_{22}A_3 - B_3) \\ & + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}+\vartheta_{22}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{12}B_2 - A_2) \\ & + \frac{\vartheta_{11}(\vartheta_{21}A_1+B_1)}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} \phi(s) ds \end{aligned} \right\} \quad (4)$$

and

$$\left. \begin{aligned} y_2(t) = & \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}+\vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{22}A_3 - B_3) \\ & + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] (\vartheta_{12}B_2 - A_2) \\ & + \frac{\vartheta_{21}(A_1+\vartheta_{11}B_1)}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} \psi(s) ds, \end{aligned} \right\} \quad (5)$$

where

$$A_1 = \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} \phi(s) ds, \quad B_1 = \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} \psi(s) ds, \quad (6)$$

$$A_2 = \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} \phi(s) ds, \quad B_2 = \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} \psi(s) ds, \quad (7)$$

$$A_3 = \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} \phi(s) ds, \quad B_3 = \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} \psi(s) ds. \quad (8)$$

DEFINITION 6. A function $(y_1, y_2) \in C^2([0, 1], \mathbb{R}) \times C^2([0, 1], \mathbb{R})$ is a solution of the coupled system of FBVP (1)–(2) if it satisfies the coupled boundary conditions (2) and there exist functions $f, g \in L^1([0, 1], \mathbb{R})$ such that $f(t) \in F(t, y_1(t), y_2(t))$, $g(t) \in G(t, y_1(t), y_2(t))$ a.e. on $[0, 1]$ and

$$\left. \begin{aligned} y_1(t) = & \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ & + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}+\vartheta_{22}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ & + \frac{\vartheta_{11}(\vartheta_{21}A_{1f}+B_{1g})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \end{aligned} \right\} \quad (9)$$

and

$$\left. \begin{aligned} y_2(t) = & \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ & + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ & + \frac{\vartheta_{21}(\vartheta_{11}B_{1g} + A_{1f})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds, \end{aligned} \right\} \quad (10)$$

where

$$\left. \begin{aligned} A_{1f} &= \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds, & B_{1g} &= \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds, \\ A_{2f} &= \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds, & B_{2g} &= \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds, \\ A_{3f} &= \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds, & B_{3g} &= \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds. \end{aligned} \right\}$$

3. Main results

Consider the Banach space $\mathcal{B} = X \times X$, where $X = \{x : x \in C[0, 1]\}$ equipped with the norm $\|(y_1, y_2)\| = \|y_1\| + \|y_2\|$, for $(y_1, y_2) \in \mathcal{B}$ and the norm is defined as

$$\|y_1\| = \sup_{t \in [0, 1]} |y_1(t)|.$$

Define the selection sets of F, G by

$$\begin{aligned} S_{F,(y_1,y_2)} &= \left\{ f \in L^1([0, 1], \mathbb{R}) : f(t) \in F(t, y_1(t), y_2(t)) \text{ for a.e. } t \in [0, 1] \right\}, \\ S_{G,(y_1,y_2)} &= \left\{ g \in L^1([0, 1], \mathbb{R}) : g(t) \in G(t, y_1(t), y_2(t)) \text{ for a.e. } t \in [0, 1] \right\}. \end{aligned}$$

In view of Lemma 4, we define the operators $\mathcal{A}_1, \mathcal{A}_2 : \mathcal{B} \rightarrow \mathcal{P}(\mathcal{B})$ by

$$\left. \begin{aligned} \mathcal{A}_1(y_1, y_2) &= \left\{ h_1 \in \mathcal{B} : \exists f \in S_{F,(y_1,y_2)}, g \in S_{G,(y_1,y_2)} \text{ such that} \right. \\ &\quad \left. h_1(y_1, y_2)(t) = Q_1(t, y_1, y_2), \forall t \in [0, 1] \right\} \\ \mathcal{A}_2(y_1, y_2) &= \left\{ h_2 \in \mathcal{B} : \exists f \in S_{F,(y_1,y_2)}, g \in S_{G,(y_1,y_2)} \text{ such that} \right. \\ &\quad \left. h_2(y_1, y_2)(t) = Q_2(t, y_1, y_2), \forall t \in [0, 1] \right\}, \end{aligned} \right\} \quad (11)$$

where

$$Q_1(y_1, y_2)(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21} + \vartheta_{22}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ + \frac{\vartheta_{11}(\vartheta_{21}A_{1f} + B_{1g})}{1 - \vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \end{cases}$$

and

$$Q_2(y_1, y_2)(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ + \frac{\vartheta_{21}(\vartheta_{11}B_{1g} + A_{1f})}{1 - \vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds. \end{cases}$$

Then we define an operator $\mathcal{A} : \mathcal{B} \rightarrow \mathcal{P}(\mathcal{B})$ by

$$\mathcal{A}(y_1, y_2)(t) = \begin{pmatrix} \mathcal{A}_1(y_1, y_2)(t) \\ \mathcal{A}_2(y_1, y_2)(t) \end{pmatrix},$$

where \mathcal{A}_1 and \mathcal{A}_2 are respectively defined by (11).

For the sake of computational convenience, we set

$$\mathcal{M}_1 = \left\{ \begin{array}{l} \left[\frac{|\vartheta_{11}|(|\vartheta_{21}\vartheta_{12}\eta^{1-q_3}| + 1) + |\vartheta_{12}|}{|1 - \vartheta_{11}\vartheta_{21}|} + \frac{|\vartheta_{22}\eta^{q_1-q_3}|}{|\Gamma(q_1 - q_3 + 1)|} \right] \\ \times \frac{|\Gamma(2 - q_3)|}{|\Delta|} + \left[\frac{|\vartheta_{11}|(|\vartheta_{21}| + |\vartheta_{22}\eta^{1-q_3}|)}{|1 - \vartheta_{11}\vartheta_{21}|} + 1 \right] \frac{|\Gamma(2 - q_3)|}{|\Delta|} \\ \times \frac{1}{|\Gamma(q_1 - q_3 + 1)|} + \left[\frac{|\vartheta_{11}\vartheta_{21}|}{|1 - \vartheta_{21}\vartheta_{11}|} + 1 \right] \frac{1}{|\Gamma(q_1 + 1)|}, \end{array} \right\} \quad (12)$$

$$\mathcal{M}_2 = \left\{ \begin{array}{l} \left[\frac{|\vartheta_{11}|(|\vartheta_{21}\vartheta_{12}\eta^{1-q_3}| + 1) + |\vartheta_{12}|}{|1 - \vartheta_{11}\vartheta_{21}|} + \frac{1}{\eta^{q_3-1}} \right] \frac{1}{\Gamma(q_2 - q_3 + 1)} \\ \times \frac{|\Gamma(2 - q_3)|}{|\Delta|} + \left[\frac{|\vartheta_{11}|(|\vartheta_{21}| + |\vartheta_{22}\eta^{1-q_3}|)}{|1 - \vartheta_{11}\vartheta_{21}|} + 1 \right] \frac{|\Gamma(2 - q_3)|}{|\Delta|} \\ \times \frac{|\vartheta_{12}\eta^{q_2-q_3}|}{|\Gamma(q_2 - q_3 + 1)|} + \frac{|\vartheta_{11}|}{|1 - \vartheta_{21}\vartheta_{11}|} \frac{1}{|\Gamma(q_2 + 1)|}, \end{array} \right\} \quad (13)$$

$$\mathcal{M}_3 = \left\{ \begin{array}{l} \left[\frac{|\vartheta_{21}|(|\vartheta_{11}\vartheta_{22}\eta^{1-q_3}| + 1) + |\vartheta_{22}|}{|1 - \vartheta_{11}\vartheta_{21}|} + \frac{1}{\eta^{q_3-1}} \right] \frac{1}{\Gamma(q_1 - q_3 + 1)} \\ \times \frac{|\Gamma(2 - q_3)|}{|\Delta|} + \left[\frac{|\vartheta_{21}|(|\vartheta_{11}| + |\vartheta_{12}\eta^{1-q_3}|)}{|1 - \vartheta_{11}\vartheta_{21}|} + 1 \right] \frac{|\Gamma(2 - q_3)|}{|\Delta|} \\ \times \frac{|\vartheta_{22}\eta^{q_1-q_3}|}{|\Gamma(q_1 - q_3 + 1)|} + \frac{|\vartheta_{21}|}{|1 - \vartheta_{21}\vartheta_{11}|} \frac{1}{|\Gamma(q_1 + 1)|}, \end{array} \right\} \quad (14)$$

$$\mathcal{M}_4 = \left\{ \begin{aligned} & \left[\frac{|\vartheta_{21}|(|\vartheta_{11}\vartheta_{22}\eta^{1-q_3}|+1)}{|1-\vartheta_{11}\vartheta_{21}|} + \left| \frac{\vartheta_{22}}{\eta^{q_3-1}} \right| \right] \frac{|\vartheta_{12}\eta^{q_2-q_3}|}{|\Gamma(q_2-q_3+1)|} \\ & \times \frac{|\Gamma(2-q_3)|}{|\Delta|} + \left[\frac{|\vartheta_{21}|(|\vartheta_{11}|+|\vartheta_{12}\eta^{1-q_3}|)}{|1-\vartheta_{11}\vartheta_{21}|} + 1 \right] \frac{|\Gamma(2-q_3)|}{|\Delta|} \\ & \times \frac{1}{|\Gamma(q_2-q_3+1)|} + \left[\frac{|\vartheta_{11}\vartheta_{21}|}{|1-\vartheta_{21}\vartheta_{11}|} + 1 \right] \frac{1}{|\Gamma(q_2+1)|}. \end{aligned} \right\} \quad (15)$$

THEOREM 1. Suppose the conditions (A0)–(A4) are fulfilled. Then the coupled system of FBVP (1)–(2) has at least one solution on $[0, 1]$.

Proof. Consider the operators $\mathcal{A}_1, \mathcal{A}_2 : \mathcal{B} \rightarrow \mathcal{P}(\mathcal{B})$ defined by (11). From (A2), it follows that the sets $S_{F,(y_1,y_2)}$ and $S_{G,(y_1,y_2)}$ are nonempty for each $(y_1, y_2) \in \mathcal{B}$. Then, for $f \in S_{F,(y_1,y_2)}$, $g \in S_{G,(y_1,y_2)}$ for $(y_1, y_2) \in \mathcal{B}$, we have

$$h_1(y_1, y_2)(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}+\vartheta_{22}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ + \frac{\vartheta_{11}(\vartheta_{21}A_{1f} + B_{1g})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds, \end{cases}$$

$$h_2(y_1, y_2)(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}+\vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{22}A_{3f} - B_{3g}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] (\vartheta_{12}B_{2g} - A_{2f}) \\ + \frac{\vartheta_{21}(\vartheta_{11}B_{1g} + A_{1f})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds, \end{cases}$$

where $h_1 \in \mathcal{A}_1(y_1, y_2)$, $h_2 \in \mathcal{A}_2(y_1, y_2)$ and so $(h_1, h_2) \in \mathcal{A}(y_1, y_2)$.

Now we verify that the operator \mathcal{A} satisfies the assumptions of the nonlinear alternative of Leray–Schauder type. This can be done in a couple of steps. In the first step, we will show that $\mathcal{A}(y_1, y_2)$ is convex valued. Let $(h_i, \bar{h}_i) \in (\mathcal{A}_1, \mathcal{A}_2)$, $i = 1, 2$. Then there exist $f_i \in S_{F,(y_1,y_2)}$, $g_i \in S_{G,(y_1,y_2)}$, $i = 1, 2$ such that, for each $t \in [0, 1]$, we have

$$h_i(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] (\vartheta_{22}A_{3f_i} - B_{3g_i}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}+\vartheta_{22}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{12}B_{2g_i} - A_{2f_i}) \\ + \frac{\vartheta_{11}(\vartheta_{21}A_{1f_i} + B_{1g_i})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f_i(s) ds, \end{cases}$$

$$\bar{h}_i(t) = \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] (\vartheta_{22}A_{3f_i} - B_{3g_i}) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] (\vartheta_{12}B_{2g_i} - A_{2f_i}) \\ + \frac{\vartheta_{21}(\vartheta_{11}B_{1g_i} + A_{1f_i})}{1-\vartheta_{21}\vartheta_{11}} + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g_i(s) ds, \end{cases}$$

where

$$\begin{aligned} A_{1f_i} &= \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f_i(s) ds, \\ B_{1g_i} &= \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g_i(s) ds, \\ A_{2f_i} &= \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_i(s) ds, \\ B_{2g_i} &= \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_i(s) ds, \\ A_{3f_i} &= \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_i(s) ds, \\ B_{3g_i} &= \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_i(s) ds. \end{aligned}$$

Let $0 \leq \omega \leq 1$. Then, for each $t \in [0, 1]$, we have

$$\begin{aligned} &[\omega h_1 + (1-\omega)h_2](t) \\ &= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} [\omega f_1(s) \right. \\ \left. + (1-\omega)f_2(s)] ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} [\omega g_1(s) + (1-\omega)g_2(s)] ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} \right. \\ \times [\omega g_1(s) + (1-\omega)g_2(s)] ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} [\omega f_1(s) \\ + (1-\omega)f_2(s)] ds \left. \right) + \frac{\vartheta_{21}}{1-\vartheta_{21}\vartheta_{11}} \left(\vartheta_{11} \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} [\omega g_1(s) \right. \\ \left. + (1-\omega)g_2(s)] ds + \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} [\omega f_1(s) + (1-\omega)f_2(s)] ds \right) \\ &\quad + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} [\omega g_1(s) + (1-\omega)g_2(s)] ds, \end{cases} \end{aligned}$$

$$\begin{aligned}
& [\omega \bar{h}_1 + (1 - \omega) \bar{h}_2](t) \\
= & \left\{ \begin{aligned}
& \frac{\Gamma(2 - q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \left(\vartheta_{22} \int_0^\eta \frac{(\eta - s)^{q_1 - q_3 - 1}}{\Gamma(q_1 - q_3)} [\omega f_1(s) \right. \\
& \left. + (1 - \omega)f_2(s)] ds - \int_0^1 \frac{(1 - s)^{q_2 - q_3 - 1}}{\Gamma(q_2 - q_3)} [\omega g_1(s) + (1 - \omega)g_2(s)] ds \right) \\
& + \frac{\Gamma(2 - q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] \left(\vartheta_{12} \int_0^\eta \frac{(\eta - s)^{q_2 - q_3 - 1}}{\Gamma(q_2 - q_3)} \right. \\
& \times [\omega g_1(s) + (1 - \omega)g_2(s)] ds - \int_0^1 \frac{(1 - s)^{q_1 - q_3 - 1}}{\Gamma(q_1 - q_3)} [\omega f_1(s) \\
& \left. + (1 - \omega)f_2(s)] ds \right) + \frac{\vartheta_{21}}{1 - \vartheta_{21}\vartheta_{11}} \left(\vartheta_{11} \int_0^1 \frac{(1 - s)^{q_2 - 1}}{\Gamma(q_2)} [\omega g_1(s) \right. \\
& \left. + (1 - \omega)g_2(s)] ds + \int_0^1 \frac{(1 - s)^{q_1 - 1}}{\Gamma(q_1)} [\omega f_1(s) + (1 - \omega)f_2(s)] ds \right) \\
& + \int_0^t \frac{(t - s)^{q_2 - 1}}{\Gamma(q_2)} [\omega g_1(s) + (1 - \omega)g_2(s)] ds.
\end{aligned} \right.
\end{aligned}$$

Since F, G are considered as convex valued, we conclude that $S_{F, (y_1, y_2)}$, $S_{G, (y_1, y_2)}$ are convex valued. Clearly $\omega h_1 + (1 - \omega)h_2 \in \mathcal{A}_1$, $\omega \bar{h}_1 + (1 - \omega) \bar{h}_2 \in \mathcal{A}_2$ and hence $\omega(h_1, \bar{h}_1) + (1 - \omega)(h_2, \bar{h}_2) \in \mathcal{A}$. Now we show that \mathcal{A} maps bounded sets into bounded sets in \mathcal{B} . For a positive number r , let $B_r = \{(y_1, y_2) \in \mathcal{B} : \| (y_1, y_2) \| \leq r\}$ be a bounded set in \mathcal{B} . Then, $\exists f \in S_{F, (y_1, y_2)}, g \in S_{G, (y_1, y_2)}$ such that

$$\begin{aligned}
& |h_1(y_1, y_2)(t)| \\
\leq & \left\{ \begin{aligned}
& \frac{|\Gamma(2 - q_3)|}{|\Delta|} \left(\frac{|\vartheta_{11}|(|\vartheta_{21}\vartheta_{12}\eta^{1-q_3}| + 1)}{|1 - \vartheta_{11}\vartheta_{21}|} + \frac{|\vartheta_{12}|}{|\eta^{q_3-1}|} \right) \left(\frac{|\vartheta_{22}\eta^{q_1 - q_3}|}{|\Gamma(q_1 - q_3 + 1)|} \right. \\
& \times ||b_1||(\Psi_1(r) + \Phi_1(r)) + \frac{1}{|\Gamma(q_2 - q_3 + 1)|} ||b_2||(\Psi_2(r) + \Phi_2(r)) \Big) \\
& + \frac{|\Gamma(2 - q_3)|}{|\Delta|} \left(\frac{|\vartheta_{11}|(|\vartheta_{21}| + |\vartheta_{22}\eta^{1-q_3}|)}{|1 - \vartheta_{11}\vartheta_{21}|} + 1 \right) \left(\frac{|\vartheta_{12}\eta^{q_2 - q_3}|}{|\Gamma(q_2 - q_3 + 1)|} \right. \\
& \times ||b_2||(\Psi_2(r) + \Phi_2(r)) + \frac{1}{|\Gamma(q_1 - q_3 + 1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) \Big) \\
& + \frac{|\vartheta_{11}|}{|1 - \vartheta_{21}\vartheta_{11}|} \left(\frac{|\vartheta_{21}|}{|\Gamma(q_1 + 1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) + \frac{1}{|\Gamma(q_2 + 1)|} \right. \\
& \times ||b_2||(\Psi_2(r) + \Phi_2(r)) \Big) + \frac{1}{|\Gamma(q_1 + 1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) \\
\leq & \mathcal{M}_1 ||b_1||(\Psi_1(r) + \Phi_1(r)) + \mathcal{M}_2 ||b_2||(\Psi_2(r) + \Phi_2(r)).
\end{aligned} \right.
\end{aligned}$$

and

$$\begin{aligned}
 & |h_2(y_1, y_2)(t)| \\
 & \leq \left\{ \begin{array}{l} \frac{|\Gamma(2-q_3)|}{|\Delta|} \left(\frac{|\vartheta_{21}|(|\vartheta_{11}| + |\vartheta_{12}\eta^{1-q_3}|)}{|1-\vartheta_{11}\vartheta_{21}|} + 1 \right) \\ \times \left(\frac{|\vartheta_{22}\eta^{q_1-q_3}|}{|\Gamma(q_1-q_3+1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) \right. \\ \left. + \frac{1}{|\Gamma(q_2-q_3+1)|} ||b_2||(\Psi_2(r) + \Phi_2(r)) \right) \\ + \frac{|\Gamma(2-q_3)|}{|\Delta|} \left(\frac{|\vartheta_{21}|(|\vartheta_{11}\vartheta_{22}\eta^{1-q_3}| + 1)}{|1-\vartheta_{11}\vartheta_{21}|} + \left| \frac{\vartheta_{22}}{\eta^{q_3-1}} \right| \right) \\ \times \left(\frac{1}{|\Gamma(q_1-q_3+1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) \right. \\ \left. + \frac{|\vartheta_{12}\eta^{q_2-q_3}|}{|\Gamma(q_2-q_3+1)|} ||b_2||(\Psi_2(r) + \Phi_2(r)) \right) \\ + \frac{|\vartheta_{21}|}{|1-\vartheta_{21}\vartheta_{11}|} \left(\frac{|\vartheta_{11}|}{|\Gamma(q_2+1)|} ||b_2||(\Psi_2(r) + \Phi_2(r)) \right. \\ \left. + \frac{1}{|\Gamma(q_2+1)|} ||b_1||(\Psi_1(r) + \Phi_1(r)) \right) \\ \left. + \frac{1}{|\Gamma(q_2+1)|} ||b_2||(\Psi_2(r) + \Phi_2(r)) \right) \\ \leq \mathcal{M}_3 ||b_1||(\Psi_1(r) + \Phi_1(r)) + \mathcal{M}_4 ||b_2||(\Psi_2(r) + \Phi_2(r)), \end{array} \right.
 \end{aligned}$$

where $\mathcal{M}_1, \mathcal{M}_2, \mathcal{M}_3$ and \mathcal{M}_4 are given in (12), (13), (14) and (15) respectively. Thus,

$$\|h_1(y_1, y_2)\| \leq \mathcal{M}_1 ||b_1||(\Psi_1(r) + \Phi_1(r)) + \mathcal{M}_2 ||b_2||(\Psi_2(r) + \Phi_2(r)),$$

and

$$\|h_2(y_1, y_2)\| \leq \mathcal{M}_3 ||b_1||(\Psi_1(r) + \Phi_1(r)) + \mathcal{M}_4 ||b_2||(\Psi_2(r) + \Phi_2(r)).$$

Hence we obtain

$$\begin{aligned}
 \|h_1, h_2\| &= \|h_1(y_1, y_2)\| + \|h_2(y_1, y_2)\| \\
 &\leq m_1 ||b_1||(\Psi_1(r) + \Phi_1(r)) + m_2 ||b_2||(\Psi_2(r) + \Phi_2(r)) \\
 &= \ell,
 \end{aligned}$$

where

$$m_1 = \mathcal{M}_1 + \mathcal{M}_3 \text{ and } m_2 = \mathcal{M}_2 + \mathcal{M}_4 \quad (16)$$

and

$$\ell = m_1 \|b_1\| (\Psi_1(r) + \Phi_1(r)) + m_2 \|b_2\| (\Psi_2(r) + \Phi_2(r)) \quad (17)$$

are constants. Next, we show that \mathcal{A} is equicontinuous. Let $t_1, t_2 \in [0, 1]$ with $t_1 < t_2$. Then, there exist $f \in S_{F, (y_1, y_2)}, g \in S_{G, (y_1, y_2)}$ such that

$$\begin{aligned} & |h_1(y_1, y_2)(t_2) - h_1(y_1, y_2)(t_1)| \\ & \leq \left\{ \begin{array}{l} \|b_1\| (\Psi_1(r) + \Phi_1(r)) \frac{|\eta^{q_1-q_3}|}{|\Gamma(q_1-q_3+1)|} \frac{|\vartheta_{22}\vartheta_{12}\eta^{1-q_3}\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_2\| (\Psi_2(r) + \Phi_2(r)) \frac{1}{|\Gamma(q_2-q_3+1)|} \frac{|\vartheta_{22}\eta^{1-q_3}\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_1\| (\Psi_1(r) + \Phi_1(r)) \frac{1}{|\Gamma(q_1-q_3+1)|} \frac{|\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_2\| (\Psi_2(r) + \Phi_2(r)) \frac{|\eta^{q_2-q_3}|}{|\Gamma(q_2-q_3+1)|} \frac{|\vartheta_{22}\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_1\| (\Psi_1(r) + \Phi_1(r)) \frac{(|t_2^{q_1}-t_1^{q_1}|)}{|\Gamma(q_1+1)|}. \end{array} \right. \end{aligned}$$

Analogously, we can obtain

$$\begin{aligned} & |h_2(y_1, y_2)(t_2) - h_2(y_1, y_2)(t_1)| \\ & \leq \left\{ \begin{array}{l} \|b_1\| (\Psi_1(r) + \Phi_1(r)) \frac{|\eta^{q_1-q_3}|}{|\Gamma(q_1-q_3+1)|} \frac{|\vartheta_{22}\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_2\| (\Psi_2(r) + \Phi_2(r)) \frac{1}{|\Gamma(q_2-q_3+1)|} \frac{|\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_1\| (\Psi_1(r) + \Phi_1(r)) \frac{1}{|\Gamma(q_1-q_3+1)|} \frac{|\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_2\| (\Psi_2(r) + \Phi_2(r)) \frac{|\eta^{q_2-q_3}|}{|\Gamma(q_2-q_3+1)|} \frac{|\vartheta_{12}\vartheta_{22}\eta^{1-q_3}\Gamma(2-q_3)|}{|\Delta|} (|t_2-t_1|) \\ + \|b_2\| (\Psi_2(r) + \Phi_2(r)) \frac{(|t_2^{q_2}-t_1^{q_2}|)}{|\Gamma(q_2+1)|}. \end{array} \right. \end{aligned}$$

It follows from the above arguments that the operator $\mathcal{A}(y_1, y_2)$ is equicontinuous, and so it is completely continuous by the Arzelá–Ascoli theorem.

For our next step, we show that the operator $\mathcal{A}(y_1, y_2)$ has a closed graph. Let $(y_{1n}, y_{2n}) \rightarrow (y_{1*}, y_{2*})$, $(h_n, \bar{h}_n) \in \mathcal{A}(y_{1n}, y_{2n})$ and $(h_n, \bar{h}_n) \rightarrow (h_*, \bar{h}_*)$, then we have to show $(h_*, \bar{h}_*) \in \mathcal{A}(y_{1*}, y_{2*})$. Observe that $(h_n, \bar{h}_n) \in \mathcal{A}(y_{1n}, y_{2n})$ implies that there exist $f_n \in S_{F, (y_{1n}, y_{2n})}$ and $g_n \in S_{G, (y_{1n}, y_{2n})}$ such that

$$h_n(y_{1n}, y_{2n})(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_n(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_n(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21} + \vartheta_{22}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_n(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_n(s) ds \right) \\ + \left[\frac{\vartheta_{11}}{1 - \vartheta_{21}\vartheta_{11}} \right] \left(\vartheta_{21} \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f_n(s) ds + \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g_n(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f_n(s) ds \end{cases}$$

and

$$\bar{h}_n(y_{1n}, y_{2n})(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} \times f_n(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_n(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21} + \frac{\vartheta_{22}t}{\eta^{q_3-1}}} \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_n(s) ds \right. \\ \left. - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_n(s) ds \right) + \frac{\vartheta_{21}}{1 - \vartheta_{21}\vartheta_{11}} \\ \times \left(\vartheta_{11} \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g_n(s) ds + \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f_n(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g_n(s) ds. \end{cases}$$

Let us consider the continuous linear operators $\Theta_1, \Theta_2 : L^1([0, 1], \mathcal{B}) \rightarrow C([0, 1], \mathcal{B})$

given by

$$\Theta_1(y_1, y_2)(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21} + \vartheta_{22}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds \right) \\ + \left[\frac{\vartheta_{11}}{1 - \vartheta_{21}\vartheta_{11}} \right] \left(\vartheta_{21} \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds + \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \end{cases}$$

and

$$\Theta_2(y_1, y_2)(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds \right) \\ + \frac{\vartheta_{21}}{1 - \vartheta_{21}\vartheta_{11}} \left(\vartheta_{11} \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds + \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds. \end{cases}$$

From Lemma 2, we know that $(\Theta_1, \Theta_2) \circ (S_F, S_G)$ is a closed graph operator. Further, we have $(h_n, \bar{h}_n) \in (\Theta_1, \Theta_2) \circ (S_{F,(y_{1n}, y_{2n})}, S_{G,(y_{1n}, y_{2n})})$ for all n . Since $(y_{1n}, y_{2n}) \rightarrow (y_{1*}, y_{2*})$, $(h_n, \bar{h}_n) \rightarrow (h_*, \bar{h}_*)$, it follows that $f_* \in S_{F,(y_1, y_2)}$ and $g_* \in S_{G,(y_1, y_2)}$ such that

$$h_*(y_{1*}, y_{2*})(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_*(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_*(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21} + \vartheta_{22}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_*(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_*(s) ds \right) \\ + \left[\frac{\vartheta_{11}}{1 - \vartheta_{21}\vartheta_{11}} \right] \left(\vartheta_{21} \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f_*(s) ds + \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g_*(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f_*(s) ds \end{cases}$$

and

$$\bar{h}_*(y_{1*}, y_{2*})(t)$$

$$= \begin{cases} \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11} + \vartheta_{12}\eta^{1-q_3})}{1 - \vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_*(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_*(s) ds \right) \\ + \frac{\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3} + 1)}{1 - \vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g_*(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f_*(s) ds \right) \\ + \frac{\vartheta_{21}}{1 - \vartheta_{21}\vartheta_{11}} \left(\vartheta_{11} \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g_*(s) ds + \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f_*(s) ds \right) \\ + \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g_*(s) ds. \end{cases}$$

that is, $(h_n, \bar{h}_n) \in \mathcal{A}(y_{1*}, y_{2*})$.

Eventually, we discuss a priori bounds on solutions. Let $(y_1, y_2) \in \vartheta \mathcal{A}(y_1, y_2)$ for $\vartheta \in (0, 1)$. Then there exist $f \in S_{F, (y_1, y_2)}$ and $g \in S_{G, (y_1, y_2)}$ such that

$$y_1(t) = \begin{cases} \frac{\vartheta\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}\vartheta_{12}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{12}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds \right) \\ + \frac{\vartheta\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{11}(\vartheta_{21}+\vartheta_{22}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds \right) \\ + \left[\frac{\vartheta\vartheta_{11}}{1-\vartheta_{21}\vartheta_{11}} \right] \left(\vartheta_{21} \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds + \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds \right) \\ + \vartheta \int_0^t \frac{(t-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \end{cases}$$

and

$$y_2(t) = \begin{cases} \frac{\vartheta\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}+\vartheta_{12}\eta^{1-q_3})}{1-\vartheta_{11}\vartheta_{21}} + t \right] \\ \times \left(\vartheta_{22} \int_0^\eta \frac{(\eta-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} \times f(s) ds - \int_0^1 \frac{(1-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds \right) \\ + \frac{\vartheta\Gamma(2-q_3)}{\Delta} \left[\frac{\vartheta_{21}(\vartheta_{11}\vartheta_{22}\eta^{1-q_3}+1)}{1-\vartheta_{11}\vartheta_{21}} + \frac{\vartheta_{22}t}{\eta^{q_3-1}} \right] \\ \times \left(\vartheta_{12} \int_0^\eta \frac{(\eta-s)^{q_2-q_3-1}}{\Gamma(q_2-q_3)} g(s) ds - \int_0^1 \frac{(1-s)^{q_1-q_3-1}}{\Gamma(q_1-q_3)} f(s) ds \right) \\ + \frac{\vartheta\vartheta_{21}}{1-\vartheta_{21}\vartheta_{11}} \left(\vartheta_{11} \int_0^1 \frac{(1-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds + \int_0^1 \frac{(1-s)^{q_1-1}}{\Gamma(q_1)} f(s) ds \right) \\ + \vartheta \int_0^t \frac{(t-s)^{q_2-1}}{\Gamma(q_2)} g(s) ds. \end{cases}$$

Using the arguments employed in Step 2, for each $t \in [0, 1]$, we obtain

$$\|y_1\| \leqslant \mathcal{M}_1 \|b_1\| \left(\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|) \right) + \mathcal{M}_2 \|b_2\| \left(\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|) \right),$$

and

$$\|y_2\| \leqslant \mathcal{M}_3 \|b_1\| \left(\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|) \right) + \mathcal{M}_4 \|b_2\| \left(\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|) \right).$$

In consequence, we have

$$\begin{aligned} \|(y_1, y_2)\| &= \|y_1\| + \|y_2\| \\ &\leqslant m_1 \|b_1\| \left(\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|) \right) + m_2 \|b_2\| \left(\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|) \right) \end{aligned}$$

which implies that

$$\frac{\|(y_1, y_2)\|}{m_1 \|b_1\| (\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|)) + m_2 \|b_2\| (\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|))} \leq 1.$$

where m_1 and m_2 are given in (16). In view of (A4), $\exists N$ such that $\|(y_1, y_2)\| \neq N$. Let us set

$$U = \{(y_1, y_2) \in \mathcal{B} : \|(y_1, y_2)\| < N\}.$$

Note that the operator $\mathcal{A} : \overline{U} \rightarrow \mathcal{P}_{cp, cv}(X) \times \mathcal{P}_{cp, cv}(X)$ is upper semicontinuous and completely continuous. From the choice of U , there is no $(y_1, y_2) \in \partial U$ such that $(y_1, y_2) \in \vartheta \mathcal{A}(y_1, y_2)$ for some $\vartheta \in (0, 1)$. Consequently, by the nonlinear alternative of Leray–Schauder type [13], we conclude that \mathcal{A} has a fixed point $(y_1, y_2) \in \overline{U}$ which is a solution of the coupled system of FBVP (1)–(2). The proof is completed. \square

4. Example

In this section, as an application, we demonstrate our results with an example. Consider the FBVP,

$$\begin{cases} {}^cD_{0+}^{\frac{8}{3}}y_1(t) \in F(t, y_1, y_2), & 0 < t < 1, \\ {}^cD_{0+}^{\frac{8}{3}}y_2(t) \in G(t, y_1, y_2), & 0 < t < 1, \end{cases} \quad (18)$$

$$\begin{cases} y_1(0) = \frac{6}{11}y_2(1), & {}^cD_{0+}^{\frac{3}{5}}y_1(1) = \frac{7}{23}{}^cD_{0+}^{\frac{3}{5}}y_2\left(\frac{13}{50}\right), \\ y_2(0) = \frac{27}{64}y_1(1), & {}^cD_{0+}^{\frac{3}{5}}y_2(1) = \frac{17}{28}{}^cD_{0+}^{\frac{3}{5}}y_1\left(\frac{13}{50}\right). \end{cases} \quad (19)$$

Here $q_1 = q_2 = \frac{8}{5}$, $\eta = \frac{13}{50}$, $q_3 = \frac{3}{5}$, $\vartheta_{11} = \frac{6}{11}$, $\vartheta_{12} = \frac{7}{23}$, $\vartheta_{21} = \frac{27}{64}$ and $\vartheta_{22} = \frac{17}{28}$. And also we have $\Gamma(q_1 - q_3 + 1) = \Gamma(q_2 - q_3 + 1) = 1$, $\Gamma(2 - q_3) = 0.8872$, $\Gamma(q_1 + 1) = \Gamma(q_2 + 1) = 1.4296$. $F, G : [0, 1] \times \mathbb{R}^2 \rightarrow \mathcal{P}(\mathbb{R})$ are multivalued maps given by

$$\begin{aligned} F(t, y_1, y_2) &= \left[\sin t + e^{-|y_1|} - \frac{|y_2|}{1 + |y_2|}, \quad \frac{|y_1|}{1 + y_1^2} + \cos(y_2) + 7t^2 + 9 \right], \\ G(t, y_1, y_2) &= \left[e^{-y_1^2} + \frac{|y_2|}{1 + |y_2|} + t^2, \quad \frac{|y_1|}{1 + |y_1|} + t + 3\cos(y_2) + 16 \right]. \end{aligned}$$

Then we have $\|F(t, y_1, y_2)\|_{\mathcal{P}} = \sup \{|f| : f \in F(t, y_1, y_2)\} \leq 18 = b_1(t)(\Psi_1(\|y_1\|) + \Phi_1(\|y_2\|))$ with $b_1(t) = 1$, $\Psi_1(\|y_1\|) = 3$, $\Phi_1(\|y_2\|) = 15$ and $\|G(t, y_1, y_2)\|_{\mathcal{P}} = \sup \{|g| : g \in G(t, y_1, y_2)\} \leq 21 = b_2(t)(\Psi_2(\|y_1\|) + \Phi_2(\|y_2\|))$ with $b_2(t) = 1$, $\Psi_2(\|y_1\|) = 3$, $\Phi_2(\|y_2\|) = 18$. From all information and also by (12), (13), (14), (15), we can

find that $\Delta = 1 - \left(\frac{7}{23}\right)\left(\frac{17}{28}\right)\left(\frac{13}{50}\right)^{2(1-\frac{3}{5})} = 0.8887$, $\mathcal{M}_1 = 3.4546$, $\mathcal{M}_2 = 1.5571$, $\mathcal{M}_3 = 2.2717$, $\mathcal{M}_4 = 2.1580$, $m_1 = \mathcal{M}_1 + \mathcal{M}_3 = 5.7263$ and $m_2 = \mathcal{M}_2 + \mathcal{M}_4 = 3.7151$. Using the condition (A4) we find that $N > 181.0905$. Hence, by applying Theorem (1), we deduce that the FBVP (18)–(19) has at least one solution on $[0, 1]$.

Acknowledgements. The authors express their profound gratitude to the editor and anonymous referees for their insightful comments and constructive suggestions which led to the improvement of this article.

REFERENCES

- [1] S. ABBAS, M. BENCHOHRA, J. R. GRAEF, J. HENDERSON, *Implicit fractional differential and integral equations: existence and stability*, **26**, Walter de Gruyter GmbH and Co KG, 2018.
- [2] S. ADLY, T. HADDAD, L. THIBAULT, *Convex sweeping process in the framework of measure differential inclusions and evolution variational inequalities*, Math. Program. Ser. B, **148** (2014), 5–47.
- [3] B. AHMAD, S. K. NTOUYAS, *Existence results for a coupled system of Caputo type sequential fractional differential equations with nonlocal integral boundary conditions*, Appl. Math. Comput., **266** (2015), 615–622.
- [4] R. P. AGARWAL, B. AHMAD, D. GAROUT, A. ALSAEDI, *Existence results for coupled nonlinear fractional differential equations equipped with nonlocal coupled flux and multi-point boundary conditions*, Chaos Solitons Fractals, **102** (2017), 149–161.
- [5] H. H. ALSULAMI, S. K. NTOUYAS, R. P. AGARWAL, B. AHMAD, A. ALSAEDI, *A study of fractional-order coupled systems with a new concept of coupled non-separated boundary conditions*, Bound. Value Probl., (2017) **2017:68**, 1–11.
- [6] J. BASTIEN, *Study of a driven and braked wheel using maximal monotone differential inclusions: applications to the nonlinear dynamics of wheeled vehicles*, Archive of Applied Mechanics, **84** (2014), 851–880.
- [7] A. BRESSAN, G. COLOMBO, *Extensions and selections of maps with decomposable values*, Studia Math., **90** (1988), 69–86.
- [8] C. CASTAING, M. VALADIER, *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Mathematics **580**, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [9] H. COVITZ, S. B. NADLER JR., *Multivalued contraction mappings in generalized metric spaces*, Israel J. Math., **8** (1970), 5–11.
- [10] K. DEIMLING, *Multivalued Differential Equations*, Walter De Gruyter, Berlin-New York, 1992.
- [11] M. F. DANCA, *Synchronization of piecewise continuous systems of fractional order*, Nonlinear Dynam., **78** (2014), 2065–2084.
- [12] Y. DING, Z. WANG, H. YE, *Optimal control of a fractional-order HIV-immune system with memory*, IEEE Trans. Contr. Sys. Techn., **20** (2012), 763–769.
- [13] A. GRANAS, J. DUGUNDJI, *Fixed Point Theory*, Springer-Verlag, New York, 2005.
- [14] M. FAIEGH, S. KUNTANAPREEDA, H. DELAVARI, D. BALEANU, *LMI-based stabilization of a class of fractional-order chaotic systems*, Nonlinear Dynam., **72** (2013), 301–309.
- [15] J. HENDERSON, N. KOSMATOV, *Eigenvalue comparison for fractional boundary value problems with the Caputo derivative*, Fract. Calc. Appl. Anal., **17** (2014), 872–880.
- [16] S. HU, N. PAPAGEORGIOU, *Handbook of Multivalued Analysis, Volume I: Theory*, Kluwer, Dordrecht, 1997.
- [17] M. JAVIDI, B. AHMAD, *Dynamic analysis of time fractional order phytoplankton-toxic phytoplankton-zooplankton system*, Ecological Modelling, **318** (2015), 8–18.
- [18] A. A. KILBAS, H. M. SRIVASTAVA, J. J. TRUJILLO, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, **204**, Elsevier Science B. V., Amsterdam, 2006.
- [19] M. KISIELEWICZ, *Stochastic Differential Inclusions and Applications*, Springer Optimization and Its Applications, **80**, Springer, New York, 2013.

- [20] M. KORDA, D. HENRION, C. N. JONES, *Convex computation of the maximum controlled invariant set for polynomial control systems*, SIAM J. Control Optim., **52** (2014), 2944–2969.
- [21] B. M. B. KRUSHNA, K. R. PRASAD, *On existence of solutions to the Caputo type fractional order three-point boundary value problems*, Int. J. Anal. Appl., **12** (2016), 60–68.
- [22] B. M. B. KRUSHNA, *Eigenvalues for iterative systems of Riemann–Liouville type p -Laplacian fractional-order boundary-value problems in Banach spaces*, Comp. Appl. Math., **39**, **81** (2020), 1–15.
- [23] A. LASOTA, Z. OPIAL, *An application of the Kakutani–Ky Fan theorem in the theory of ordinary differential equations*, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys., **13** (1965), 781–786.
- [24] M. MONTEIRO, D. P. MANUEL, *Differential Inclusions in Nonsmooth Mechanical Problems: Shocks and Dry Friction*, Progress in Nonlinear Differential Equations and their Applications, **9**, Birkhauser Verlag, Basel, 1993.
- [25] I. PETRAS, R. L. MAGIN, *Simulation of drug uptake in a two compartmental fractional model for a biological system*, Commun. Nonlinear Sci. Numer. Simul., **16** (2011), 4588–4595.
- [26] I. PODLUBNY, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [27] K. R. PRASAD, B. M. B. KRUSHNA, *Multiple positive solutions for a coupled system of Riemann–Liouville fractional order two-point boundary value problems*, Nonlinear Stud., **20** (2013), 501–511.
- [28] K. R. PRASAD, B. M. B. KRUSHNA, *Eigenvalues for iterative systems of Sturm–Liouville fractional order two-point boundary value problems*, Fract. Calc. Appl. Anal., **17** (2014), 638–653.
- [29] V. RECUPERO, *A continuity method for sweeping processes*, J. Differential Equations, **251** (2011), 2125–2142.
- [30] P. RICHARD, M. NICODEMI, R. DELANNAY, P. RIBIERE, D. BIDEAU, *Slow relaxation and compaction of granular system*, Nature Mater., **4** (2005), 121–128.
- [31] B. SENOL, C. YEROGLU, *Frequency boundary of fractional order systems with nonlinear uncertainties*, J. Franklin Inst., **350** (2013), 1908–1925.
- [32] I. M. SOKOLOV, J. KLAFTER, A. BLUMEN, *Fractional Kinetics*, Phys. Today, **55** (2002), 48–54.
- [33] J. R. WANG, Y. ZHANG, *Analysis of fractional order differential coupled systems*, Math. Methods Appl. Sci., **38** (2015), 3322–3338.
- [34] F. ZHANG, G. CHEN, C. LI, J. KURTHS, *Chaos synchronization in fractional differential systems*, Phil. Trans. R. Soc. A., **371** (2013), 20120155.
- [35] Y. ZHOU, B. AHMAD, A. ALSAEDI, *Existence of nonoscillatory solutions for fractional neutral differential equations*, Appl. Math. Lett., **72** (2017), 70–74.

(Received September 29, 2020)

B. M. B. Krushna

Department of Mathematics
MVGR College of Engineering (A)
Visianagaram, 535 005, India
e-mail: muraleebalu@yahoo.com

K. R. Prasad

Department of Applied Mathematics
Andhra University
Visakhapatnam, 530 003, India
e-mail: rajendra92@rediffmail.com

P. Veeraiyah

Department of Applied Mathematics
Andhra University
Visakhapatnam, 530 003, India