

SPACE-TIME ANALYTIC SMOOTHING EFFECT FOR THE NONLINEAR SCHRÖDINGER EQUATIONS WITH NONLINEARITY OF EXPONENTIAL TYPE

GAKU HOSHINO

(Communicated by J. Ildefonso Díaz)

Abstract. In this paper, we consider the global Cauchy problem for the nonlinear Schrödinger equations with nonlinearity of exponential type in higher space dimensions $n \geq 2$. In particular, we study the global existence of the solutions to the Cauchy problem with small data in the framework of intersection of Sobolev and weighted Lebesgue space: $H^{n/2} \cap \mathcal{F}H^{n/2}$. More precisely, we show that if data decay exponentially in $H^{n/2} \cap \mathcal{F}H^{n/2}$ then for any time $t \neq 0$, solutions are real-analytic in both space and time variables and have analytic continuation.

1. Introduction

We study the Cauchy problem for the nonlinear Schrödinger equations in $n \geq 2$ space dimensions as follows:

$$\begin{cases} i\partial_t u + \frac{1}{2}\Delta u = f(u), \\ u(0, x) = \phi(x), \quad x \in \mathbb{R}^n \end{cases} \quad (1.1)$$

where $u : \mathbb{R} \times \mathbb{R}^n \ni (t, x) \mapsto u(t, x) \in \mathbb{C}$, $i = \sqrt{-1}$, $\partial_t = \partial/\partial t$ and $\Delta = \sum_{k=1}^n \partial^2/\partial x_k^2$.

In this study we treat the following nonlinearity of exponential type:

$$f(u) = \left(e^{\lambda|u|^2} - 1 \right) u \quad (1.2)$$

where $\lambda \in \mathbb{C}$. The nonlinear Schrödinger equation with the nonlinearity of exponential type appears in Physics to study laser beams in plasma (see [20] for instance). The Cauchy problem for (1.1)–(1.2) has been studied in [22] with sufficiently small Cauchy data in the Sobolev space $H^{n/2}$ (see Remark 1 in [22]). The Sobolev space $H^{n/2}$ is the critical Sobolev space of (1.1)–(1.2). In this previous study, the critical Sobolev inequality (see [19, 23] for instance)

$$\|u\|_{L^q} \leq Cq^{1/2} \left\| (-\Delta)^{n/4} u \right\|_{L^2}^{1-2/q} \|u\|_{L^2}^{2/q}$$

Mathematics subject classification (2020): 35Q55.

Keywords and phrases: Nonlinear Schrödinger equations, nonlinearity of exponential type, space-time analytic smoothing effect.

This work was supported by JSPS KAKENHI Grant Number 19K14570.

for $2 < q < \infty$, has been applied to estimate the nonlinear term with an exponential growth.

Our main purpose of this study is to show the space-time analytic smoothing effect for (1.1)–(1.2) in $n \geq 2$ with sufficiently small Cauchy data in weighted Sobolev space $H^{n/2} \cap \mathcal{F}H^{n/2}$.

The analytic smoothing effect in space variables for (1.1)–(1.2) in space dimensions $n = 2$ with sufficiently small data in H^1 has been studied in [12] by author with Ozawa. As far as author knows there are no results on space-time analytic smoothing effect for (1.1)–(1.2). Therefore this study is an extension of the previous study [12].

The analyticity and analytic smoothing effect for nonlinear Schrödinger equations has been studied by many authors (see [2, 5, 9, 10, 11, 12, 13, 14, 21, 24] and references therein). The existence of scattering operator for (1.1)–(1.2) in $H^{n/2}$ setting with $n \geq 2$ has been studied in [25]. The scattering for (1.1)–(1.2) in $n = 2$ is studied in [15, 26].

The space-time analytic smoothing effect for local solutions to the nonlinear Schrödinger equations is firstly studied by Hayashi and Kato in [5], by applying the Galilei generator

$$J(t) = x + it\nabla$$

and the pseudo-conformal generator

$$K(t) = |x|^2 + it(2t\partial_t + x \cdot \nabla + \nabla \cdot x).$$

In this previous study, Hayashi and Kato have introduced the following types of function spaces

$$X_T = \{u; \|u\|_{X_T} < \infty\}, \|u\|_{X_T} = \sum_{k=0}^{\infty} \sum_{\alpha \in (\mathbb{N} \cup \{0\})^n} \frac{a^k b^{|\alpha|}}{k! |\alpha|!} \|K^k J^\alpha u\|_{L^\infty([0,T]; L^2)} \quad (1.3)$$

to show the analyticity of solutions in both space and time variables, where $a, b > 0$.

The space-time analytic smoothing effect for global solutions to the nonlinear Schrödinger equations with pseudo-conformal nonlinearity has been studied in [14] by author with Ozawa and the space-time analytic smoothing effect for local solutions to the nonlinear Schrödinger equations with non pseudo-conformal nonlinearity has been studied in [10] by author, where the nonlinearity

$$F(u) = \lambda |u|^{4/n} u$$

is called pseudo-conformal nonlinearity because the nonlinear Schrödinger equations:

$$\begin{cases} i\partial_t u + \frac{1}{2} \Delta u = \lambda |u|^{4/n} u, \\ u(0) = \phi \end{cases} \quad (1.4)$$

are invariant under the pseudo-conformal transform Ψ_θ :

$$(\Psi_\theta u)(t, x) = (1 - \theta t)^{-n/2} e^{-i \frac{\theta |x|^2}{2(1-\theta t)}} u\left(\frac{t}{1 - \theta t}, \frac{x}{1 - \theta t}\right).$$

That is if u satisfy (1.4) then also $\Psi_\theta u$ satisfy (1.4) with the Cauchy data $e^{-i\theta|x|^2/2}\phi$. The commutation relation between K and $i\partial_t + (1/2)\Delta$ is

$$\left(i\partial_t + \frac{1}{2}\Delta\right)K = (K + 4it)\left(i\partial_t + \frac{1}{2}\Delta\right)$$

and $K + 4it$ behaves like the differential operator for the pseudo-conformal nonlinearity ([14]):

$$(K(t) + 4it)^k(|u|^{4/n}u) = \sum_{l_1+l_2+\dots+l_{1+4/n}=k} \frac{(-1)^{l_2+\dots+l_{1+4/n}} k!}{l_1!l_2!\dots l_{1+4/n}!} \left(\prod_{j=1}^{1+4/n} \left(K^{l_j}(t) u_j \right)^{[(-1)^{j+1}]} \right) \quad (1.5)$$

where $u^{[+1]} = u, u^{[-1]} = \bar{u}$ and $k \geq 1$. In [10], author has shown the generalization of (1.5) for non pseudo-conformally invariant nonlinearity $|u|^{2\sigma}u$, as follows

$$\begin{aligned} & (K(t) + 4it)^k \left(\prod_{j=1}^{2\sigma+1} u_j^{[(-1)^{j+1}]} \right) \\ &= \sum_{k_1+k_2=k} \frac{k!}{k_1!k_2!} \cdot \mathcal{R}_\sigma^{k_2}(t) \\ & \times \left(\sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{(-1)^{l_2+l_4+\dots+l_{2\sigma}} k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \left(\prod_{j=1}^{2\sigma+1} \left(K^{l_j}(t) u_j \right)^{[(-1)^{j+1}]} \right) \right). \end{aligned} \quad (1.6)$$

This formula involves the remainder term:

$$\mathcal{R}_\sigma^{k_2}(t) = \prod_{l=1}^{k_2} (2i(l+1-n\sigma)t) = \prod_{l=1}^{n\sigma-2} (2i(l+1-n\sigma)t).$$

If

$$2\sigma + 1 > 1 + \frac{4}{n}$$

then $\mathcal{R}_\sigma^{k_2}(t)$ is estimated as

$$\begin{aligned} |\mathcal{R}_\sigma^{k_2}(t)| &\leq \prod_{l=1}^{n\sigma-2} 2|t||l+1-n\sigma| \\ &\leq 2^{n\sigma-2}|t|^{n\sigma-2}(2n\sigma-1)^{n\sigma-2} \end{aligned}$$

for $|t| \geq 1$. Therefore the increasing term $|t|^{n\sigma-2}$ appears in the estimate of the nonlinear term

$$f(u) = (e^{\lambda|u|^2} - 1)u = \sum_{\sigma=1}^{\infty} \frac{\lambda^\sigma}{\sigma!} |u|^{2\sigma} u$$

in the function space like (1.3) by applying the formula (1.6). This is a difficulty in considering this issue and this point is different from the previous study [12]. To overcome this difficulty we introduce the operator ([3, 7, 8]):

$$|J|^{n/2}(t) = e^{i(t/2)\Delta} |x|^{n/2} e^{-i(t/2)\Delta}$$

because $|J|^{n/2}(t)$ provides the time decay estimate

$$\|u(t)\|_{L^q} \leq C q^{1/2} |t|^{-n(\frac{1}{2}-\frac{1}{q})} \| |J|^{n/2} u(t) \|_{L^2}^{1-2/q} \|u(t)\|_{L^2}^{2/q}$$

for $2 < q < \infty$. From this estimate we get the decreasing term $|t|^{-n\sigma+2}$ and hence we are able to control the increasing term $|t|^{n\sigma-2}$ in the process of nonlinear estimate.

To state our main results precisely, we introduce the notation. We denote $L^p = L^p(\mathbb{R}^n)$ by the usual Lebesgue space with $1 \leq p \leq \infty$. The Fourier transform $\mathcal{F} : \varphi \mapsto \hat{\varphi}$ is defined by

$$\hat{\varphi}(\xi) = (2\pi)^{-n/2} \int_{\mathbb{R}^2} e^{-i\xi \cdot x} \varphi(x) dx, \quad \xi \in \mathbb{R}^n$$

and the inverse Fourier transform $\mathcal{F}^{-1} : \psi \mapsto \psi^\vee$ is defined by

$$\psi^\vee(x) = (2\pi)^{-n/2} \int_{\mathbb{R}^2} e^{ix \cdot \xi} \psi(\xi) d\xi, \quad x \in \mathbb{R}^n.$$

The free Schrödinger propagator is defined by

$$U(t)\varphi = \left(e^{-i(t/2)|\xi|^2} \hat{\varphi} \right)^\vee, \quad t \in \mathbb{R}.$$

The Sobolev space H^s with $s > 0$ is defined by

$$H_p^s = \left\{ \varphi \in \mathcal{S}' ; \|\varphi\|_{H_p^s} = \|\varphi\|_{L^p} + \|(-\Delta)^{s/2}\varphi\|_{L^p} < \infty \right\},$$

where $(-\Delta)^{s/2}\varphi = (|\xi|^s \hat{\varphi})^\vee$. We denote H_2^s by H^s for simplicity. The weighted Sobolev space $\mathcal{F}H^s$ is defined by

$$\mathcal{F}H^s = \left\{ \varphi \in \mathcal{S}' ; \|\varphi\|_{\mathcal{F}H^s} = \|\varphi\|_{L^2} + \||x|^s \varphi\|_{L^2} < \infty \right\}.$$

We say that the pair (r, p) is an admissible pair if

$$\frac{2}{r} = n \left(\frac{1}{2} - \frac{1}{p} \right)$$

with $2 \leq p < \infty$ if $n = 2$ and $2 \leq p \leq 2n/(n-2)$ if $n \geq 3$. Let $I \subset \mathbb{R}$. We define the following function spaces

$$X^s(I) = L^\infty(I; H^s) \cap L^{2+\varepsilon}(I; H_{2+4/\varepsilon}^s)$$

for $n = 2$, where constant $0 < \varepsilon < 10^{-1}$, and

$$X^s(I) = L^\infty(I; H^s) \cap L^2\left(I; H_{2n/(n-2)}^s\right)$$

for $n \geq 3$. Let $t \in \mathbb{R}$. We introduce the generator of Galilei transform

$$J(t)\varphi = (x + it\nabla)\varphi = U(t)xU(-t)\varphi.$$

J has another representation ([6]):

$$J(t)\varphi = M(t)it\nabla(M(-t)\varphi) \quad (1.7)$$

where $M(t) = e^{i|x|^2/2t}$ for $t \neq 0$. We also define the operator $|J|^\gamma$ by

$$|J|^\gamma(t)\varphi = U(t)|x|^\gamma U(-t)\varphi, t \in \mathbb{R}$$

and which has another representation (see [3, 7, 8]):

$$|J|^\gamma(t)\varphi = M(t)|t|^\gamma(-\Delta)^{\gamma/2}(M(-t)\varphi)$$

for $t \neq 0$. The generalized Sobolev space $A_p^\gamma(t)$ with $\gamma > 0$ is defined by

$$A_p^\gamma(t) = \left\{ \varphi \in \mathcal{S}'; \|\varphi\|_{A_p^\gamma(t)} = \|\varphi\|_{L^p} + \left\| |J|^\gamma(t)\varphi \right\|_{L^p} < \infty \right\}, t \in \mathbb{R}.$$

Note that $A_2^\gamma(0) = \mathcal{F}H^\gamma$. We define the following function spaces

$$X_\gamma(I) = L^\infty(I; A_2^\gamma) \cap L^{2+\varepsilon}\left(I; A_{2+4/\varepsilon}^\gamma\right)$$

for $n = 2$, where ε is the constant appears in definition of $X^s(I)$ above, and

$$X_\gamma(I) = L^\infty(I; A_2^\gamma) \cap L^2\left(I; A_{2n/(n-2)}^\gamma\right)$$

for $n \geq 3$. Let $t \in \mathbb{R}$. The generator of pseudo-conformal transform K is defined by

$$K(t)\varphi = \left(|x|^2 + it(2t\partial_t + x \cdot \nabla + \nabla \cdot x)\right)\varphi = U(t)(|x|^2 + 2it^2\partial_t)(U(-t)\varphi).$$

K also has another representation

$$K(t)\varphi = M(t)it(2t\partial_t + x \cdot \nabla + \nabla \cdot x)(M(-t)\varphi)$$

for $t \neq 0$. By the definition above, we see that

$$K(t)J(t)\varphi = J(t)K(t)\varphi$$

for all $t \in \mathbb{R}$.

Let $a = (a_0, a_1, a_2, \dots, a_n) \in (0, \infty)^{1+n}$. We define the following function space of generalized analytic function:

$$G_\gamma^{s,a}(I) = \left\{ u \in X^s(I) \cap X_\gamma(I); \|u\|_{G_\gamma^{s,a}(I)} < \infty \right\},$$

$$\|u\|_{G_\gamma^{s,a}(I)} = \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \|K^k J^\alpha u\|_{X^s(I) \cap X_\gamma(I)}$$

where $a^{(k,\alpha)} = a_0^k a_1^{\alpha_1} a_2^{\alpha_2} \cdots a_n^{\alpha_n}$, $(k,\alpha)! = k! \alpha!$. If $s = \gamma = n/2$, we write

$$G^a(I) = G_{n/2}^{n/2,a}(I)$$

for simplicity. We now state our main result.

THEOREM 1. *Let $n \geq 2$. If $\phi \in H^{n/2} \cap \mathcal{F}H^{n/2}$ satisfying*

$$\sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \| |x|^{2k} x^\alpha \phi \|_{H^{n/2} \cap \mathcal{F}H^{n/2}} \leq \rho$$

for sufficiently small $\rho > 0$ with some $a \in (0, \infty)^{1+n}$. Then there exists a unique global solution $u \in G^a(\mathbb{R})$ to (1.1)–(1.2).

REMARK 1. By [5], we see that the solution $u \in G^a(\mathbb{R})$ has the following property. For any bounded domain $\Gamma \subset (\mathbb{R} \setminus \{0\}) \times \mathbb{R}^n$ there exists $c_j = c_j(a, \Gamma) > 0$, $0 \leq j \leq n$ such that $u(t), t \neq 0$ is real analytic in both space and time variables on Γ and has an analytic continuation to $\tilde{\Gamma}$ where

$$\tilde{\Gamma} = \left\{ (t + i\tau, x + iy) \in \mathbb{C}^{1+n}; \right. \\ \left. (t, x) \in \Gamma, 0 \leq |\tau| < c_0 |t|^2, 0 \leq |y_j| < c_j |t|, 1 \leq j \leq n \right\}.$$

2. Preliminaries

LEMMA 1. ([1, 18, 27]) *Let $n \geq 2$. The following inequalities hold:*

$$\|U(\cdot)\phi\|_{G_\gamma^{s,a}(\mathbb{R})} \leq C \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \| |x|^{2k} x^\alpha \phi \|_{H^s \cap \mathcal{F}H^\gamma}$$

and

$$\left\| \int_0^{(\cdot)} U(\cdot - s) F(s) ds \right\|_{G_\gamma^{s,a}(\mathbb{R})} \leq C \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \| K^k J^\alpha F \|_{L^{r'}(\mathbb{R}; H_{p'}^s \cap A_{p'}^\gamma)}$$

where (r, p) is the admissible pair and q' is the Hölder conjugate of $1 \leq q \leq \infty$.

The following lemma has an important role to control the nonlinearity of exponential type.

LEMMA 2. ([3, 7, 19, 23]) *For any $2 < q < \infty$. We have the following estimates*

$$(1) \quad \|\varphi\|_{L^q} \leq Cq^{1/2} \left\| |\nabla|^{n/2} \varphi \right\|_{L^2}^{1-2/q} \|\varphi\|_{L^2}^{2/q} \quad (2.1)$$

where the constant $C > 0$.

$$(2) \quad \|\varphi\|_{L^q} \leq Cq^{1/2}|t|^{-n\left(\frac{1}{2}-\frac{1}{q}\right)} \left\| |J|^{n/2}(t)\varphi \right\|_{L^2}^{1-2/q} \|\varphi\|_{L^2}^{2/q}, \quad t \neq 0 \quad (2.2)$$

where the constant $C > 0$.

We need the next lemma to calculate the operation of $K + 4it$ to the nonlinear term.

LEMMA 3. ([10]) *Let $t \in \mathbb{R}$ and $\sigma \geq 1$. We have the following equality*

$$(K(t) + 4it)^k \left(\prod_{j=1}^{2\sigma+1} v_j^{[\varepsilon_j]} \right) = \sum_{k_1+k_2=k} \frac{k!}{k_1!k_2!} \mathcal{R}_\sigma^{k_2}(t) \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} v_j^{[\varepsilon_j]} \right] (t)$$

for all $k \geq 1$. Here $v^{[+1]} = v$, $v^{[-1]} = \bar{v}$, $\varepsilon_j = (-1)^{j+1}$,

$$\begin{aligned} \mathcal{L}^m \left[\prod_{j=1}^{2\sigma+1} v_j^{[\varepsilon_j]} \right] (t) &= \sum_{l_1+l_2+\dots+l_{2\sigma+1}=m} \frac{(-1)^{l_2+l_4+\dots+l_{2\sigma}} m!}{l_1!l_2!\dots l_{2\sigma+1}!} \left(\prod_{j=1}^{2\sigma+1} (K^{l_j}(t)v_j)^{[\varepsilon_j]} \right), \\ \mathcal{R}_\sigma^m(t) &= \prod_{l=1}^m (2i(l+1-n\sigma)t) \end{aligned}$$

for $m \geq 1$ and $\mathcal{L}^0[F] = F$, $\mathcal{R}_\sigma^0 = 1$.

Proof. See the proof of Lemma 4 in [10]. \square

3. Proof of Theorem 1

Our proof is based on the standard contraction argument. We define the metric space (B_R^a, d) by

$$\begin{aligned} B_R^a &= \{u \in G^a(\mathbb{R}); \|u\|_{G^a(\mathbb{R})} \leq R\}, \\ d(u, v) &= \|u - v\|_{G^a(\mathbb{R})}. \end{aligned}$$

We see that (B_R^a, d) is the complete metric space. We introduce the operator $\Phi : u \mapsto \Phi u$ defined by

$$\Phi u(t) = U(t)\phi - i \int_0^t U(t-s) \left((e^{\lambda|u|^2} - 1)u \right)(s) ds, \quad t \in \mathbb{R}.$$

Let $(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}$. Then we see that

$$K^k J^\alpha \Phi u(t) = U(t) |x|^{2k} x^\alpha \phi - i \int_0^t U(t-s) (K(s) + 4is)^k J^\alpha(s) \left((e^{\lambda|u|^2} - 1) u \right)(s) ds$$

for $t \in \mathbb{R}$. To prove the map Φ is a contraction mapping in (B_R^a, d) for sufficiently small $R > 0$, we need to estimate the norm $\|\Phi u\|_{G^a(\mathbb{R})}$. We estimate

$$\begin{aligned} \|\Phi u\|_{G^a(\mathbb{R})} &\leq C \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| |x|^{2k} x^\alpha \phi \right\|_{H^{n/2} \cap \mathcal{F} H^{n/2}} \\ &\quad + C \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| (K + 4it)^k J^\alpha \left((e^{\lambda|u|^2} - 1) u \right) \right\|_{L^{v'} \left(\mathbb{R}; H_{\rho'}^{n/2} \cap A_{\rho'}^{n/2} \right)} \\ &\leq C \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| |x|^{2k} x^\alpha \phi \right\|_{H^{n/2} \cap \mathcal{F} H^{n/2}} \\ &\quad + C \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| (K + 4it)^k J^\alpha \left((e^{\lambda|u|^2} - 1) u \right) \right\|_{L^{v'} \left([-1, 1]; H_{\rho'}^{n/2} \cap A_{\rho'}^{n/2} \right)} \\ &\quad + C \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| (K + 4it)^k J^\alpha \left((e^{\lambda|u|^2} - 1) u \right) \right\|_{L^{v'} \left(\mathbb{R} \setminus [-1, 1]; H_{\rho'}^{n/2} \cap A_{\rho'}^{n/2} \right)} \end{aligned}$$

where $(v, \rho) = (3, 6)$ for $n = 2$ and $(v, \rho) = (2, 2n/(n-2))$ for $n \geq 3$. The nonlinear term is written as

$$(e^{\lambda|u|^2} - 1) u = \sum_{\sigma=1}^{\infty} \frac{\lambda^\sigma}{\sigma!} |u|^{2\sigma} u = \sum_{\sigma=1}^{\infty} \frac{\lambda^\sigma}{\sigma!} \left(\prod_{j=1}^{2\sigma+1} u^{[\varepsilon_j]} \right)$$

where $u^{[+1]} = u$, $u^{[-1]} = \bar{u}$, $\varepsilon_j = (-1)^{j+1}$. Then we see that

$$J^\alpha(|u|^{2\sigma} u) = \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \frac{(-1)^{|\beta_2| + |\beta_4| + \dots + |\beta_{2\sigma}|} \alpha!}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left(J^{\beta_j} u \right)^{[\varepsilon_j]}$$

and

$$\begin{aligned} &(K + 4it)^k J^\alpha(|u|^{2\sigma} u) \\ &= \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \frac{(-1)^{|\beta_2| + |\beta_4| + \dots + |\beta_{2\sigma}|} \alpha!}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} (K + 4it)^k \left(\prod_{j=1}^{2\sigma+1} \left(J^{\beta_j} u \right)^{[\varepsilon_j]} \right) \\ &= \sum_{k_1 + k_2 = k} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \frac{k!}{k_1! k_2!} \frac{(-1)^{|\beta_2| + |\beta_4| + \dots + |\beta_{2\sigma}|} \alpha!}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} \left(J^{\beta_j} u \right)^{[\varepsilon_j]} \right]. \end{aligned}$$

We see that

$$\mathcal{R}_\sigma^{k_2}(t) = \prod_{l=1}^{k_2} (2i(l+1-n\sigma)t) = 0$$

for $k_2 \geq n\sigma - 1$ with $\sigma \in \mathbb{Z}_{\geq 1} \cap [2/n, \infty)$, and

$$\begin{aligned} |\mathcal{R}_\sigma^{k_2}(t)| &= \left| \prod_{l=1}^{k_2} (2i(l+1-n\sigma)t) \right| \\ &= 2^{k_2} |t|^{k_2} \prod_{l=1}^{k_2} |l+1-n\sigma| \\ &\leq 2^{k_2} |t|^{k_2} (k_2 + 1 + n\sigma)^{k_2} \\ &\leq 2^{k_2} |t|^{k_2} (2n\sigma - 1)^{k_2} \\ &\leq 2^{n\sigma-2} |t|^{n\sigma-2} (2n\sigma - 1)^{n\sigma-2} \end{aligned} \quad (3.1)$$

for $1 \leq k_2 \leq n\sigma - 2$ with $\sigma \in \mathbb{Z}_{\geq 1} \cap [3/n, \infty)$. We remark that $\prod_Q f_j = 1$ for $Q = \emptyset$.

The remainder term \mathcal{R}^{k_2} involves the term $|t|^{n\sigma-2}$ which increases for $|t| > 0$. For this reason we introduce the function space $A_2^{n/2}$ because we are able to gain the term $|t|^{-n\sigma+2}$ by applying the critical Sobolev inequality (2.2) in Lemma 2, above.

We introduce the subsets $\Omega_\sigma \subset \mathbb{Z}$ and $\Lambda_j \subset \mathbb{Z}$ defined by

$$\Omega_\sigma = \{1, 2, 3, \dots, 2\sigma + 1\}$$

and

$$\Lambda_j = \begin{cases} \{2\} & (j = 1) \\ \{j-1\} & (j \geq 2) \end{cases}$$

respectively. We define the characteristic function on K_σ by

$$\chi_{K_\sigma}(k) = \begin{cases} 1 & (k \in K_\sigma) \\ 0 & (k \in \mathbb{Z} \setminus K_\sigma) \end{cases}$$

with the subset

$$K_\sigma = \{k \in \mathbb{Z}; 1 \leq k_2 \leq n\sigma - 2\} \subset \mathbb{Z}.$$

Next, we consider the nonlinear estimate. The discussion of nonlinear estimate is divided in to two cases, Case 1: $n = 2$ and Case 2: $n \geq 3$.

Case 1: $n = 2$

Let $n = 2$. By the Kato–Ponce inequality we have ([4, 16, 17])

$$\left\| \prod_{j=1}^{2\sigma+1} v_j \right\|_{H_{6/5}^1} \leq C \sum_{j=1}^{2\sigma+1} \|v_j\|_{H_6^1} \left\| \prod_{\mu \neq j} v_\mu \right\|_{L^{3/2}}$$

and

$$\left\| \prod_{j=1}^{2\sigma+1} v_j \right\|_{A_{6/5}^1(t)} \leq C \sum_{j=1}^{2\sigma+1} \|v_j\|_{A_6^1(t)} \left\| \prod_{\mu \neq j} v_\mu \right\|_{L^{3/2}}.$$

From this inequalities and the critical Sobolev inequality (2.1) in Lemma 2, we estimate

$$\begin{aligned}
& \left\| \left(\mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right)(t) \right\|_{H_{6/5}^1 \cap A_{6/5}^1(t)} \\
& \leq \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1! l_2! \cdots l_{2\sigma+1}!} \left\| \prod_{j=1}^{2\sigma+1} (K^{l_j} J^{\beta_j} u(t))^{[\varepsilon_j]} \right\|_{H_{6/5}^1 \cap A_{6/5}^1(t)} \\
& \leq C \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1! l_2! \cdots l_{2\sigma+1}!} \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{H_6^1 \cap A_6^1(t)} \left\| \prod_{\mu \neq j} K^{l_\mu} J^{\beta_\mu} u \right\|_{L^{3/2}} \right) \\
& \leq C \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1! l_2! \cdots l_{2\sigma+1}!} \\
& \quad \times \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u(t) \right\|_{H_6^1 \cap A_6^1(t)} \prod_{\mu \in \Lambda_j} \left\| K^{l_\mu} J^{\beta_\mu} u(t) \right\|_{L^6} \prod_{v \in \Omega_\sigma \setminus (\Lambda_j \cup \{j\})} \left\| K^{l_v} J^{\beta_v} u(t) \right\|_{L^{2(2\sigma-1)}} \right) \\
& \leq C (2(2\sigma-1))^{\sigma-1/2} \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1! l_2! \cdots l_{2\sigma+1}!} \\
& \quad \times \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u(t) \right\|_{H_6^1 \cap A_6^1(t)} \prod_{\mu \in \Lambda_j} \left\| K^{l_\mu} J^{\beta_\mu} u(t) \right\|_{L^6} \prod_{v \in \Omega_\sigma \setminus (\Lambda_j \cup \{j\})} \left\| K^{l_v} J^{\beta_v} u(t) \right\|_{H^1} \right)
\end{aligned}$$

for $|t| < 1$, and by using the critical Sobolev inequality (2.2) in Lemma 2, we estimate

$$\begin{aligned}
& \left\| \left(\mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right)(t) \right\|_{H_{6/5}^1 \cap A_{6/5}^1(t)} \\
& \leq C (2(2\sigma-1))^{\sigma-1/2} |t|^{-2\sigma+2} \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1! l_2! \cdots l_{2\sigma+1}!} \\
& \quad \times \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u(t) \right\|_{A_6^1(t) \cap H_6^1} \prod_{\mu \in \Lambda_j} \left\| K^{l_\mu} J^{\beta_\mu} u(t) \right\|_{L^6} \prod_{v \in \Omega_\sigma \setminus (\Lambda_j \cup \{j\})} \left\| K^{l_v} J^{\beta_v} u(t) \right\|_{A_2^1(t)} \right)
\end{aligned}$$

for $|t| \geq 1$. Therefore by the above estimates with (3.1) and inequality

$$\|\psi\|_{L^3(I; L^6)} \leq \|\psi\|_{L^{2+\varepsilon}(I; L^{2+4/\varepsilon})}^{(2+\varepsilon)/3} \|\psi\|_{L^\infty(I; L^2)}^{1-(2+\varepsilon)/3},$$

we have

$$\begin{aligned}
& \left\| \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}([-1, 1]; H_{6/5}^1 \cap A_{6/5}^1)} \\
& = \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}([-1, 1]; H_{6/5}^1 \cap A_{6/5}^1)}
\end{aligned}$$

$$\begin{aligned}
&\leq C(2(2\sigma-1))^{\sigma-1/2}(4\sigma-1)^{k_2}2^{k_2}\chi_{K_\sigma}(k_2) \\
&\quad \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\cdots l_{2\sigma+1}!} \chi_{K_\sigma}(k_2) \prod_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u\|_{X^{1/2}([-1,1]) \cap X_{1/2}([-1,1])} \\
&\leq C(2(2\sigma-1))^{\sigma-1/2}(4\sigma-1)^{k_2}2^{k_2}\chi_{K_\sigma}(k_2) \\
&\quad \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\cdots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u\|_{X^1([-1,1]) \cap X_1([-1,1])}
\end{aligned}$$

and

$$\begin{aligned}
&\left\| \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}(\mathbb{R} \setminus [-1,1]; H_{6/5}^1 \cap A_{6/5}^1)} \\
&= \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}(\mathbb{R} \setminus [-1,1]; H_{6/5}^1 \cap A_{6/5}^1)} \\
&\leq C(2(2\sigma-1))^{\sigma-1/2}(4\sigma-1)^{k_2}2^{k_2}\chi_{K_\sigma}(k_2) \\
&\quad \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\cdots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u\|_{X^1(\mathbb{R} \setminus [-1,1]) \cap X_1(\mathbb{R} \setminus [-1,1])} \\
&\leq C(2(2\sigma-1))^{\sigma-1/2}(4\sigma-1)^{k_2}2^{k_2}\chi_{K_\sigma}(k_2) \\
&\quad \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\cdots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u\|_{X^1(\mathbb{R} \setminus [-1,1]) \cap X_1(\mathbb{R} \setminus [-1,1])}
\end{aligned}$$

where we have used the estimates

$$|\mathcal{R}_\sigma^{k_2}(t)| \leq 2^{k_2}(4\sigma-1)^{k_2} \quad \text{for } |t| < 1$$

and

$$|\mathcal{R}_\sigma^{k_2}(t)| \leq 2^{k_2}(4\sigma-1)^{k_2}|t|^{2\sigma-2} \quad \text{for } |t| \geq 1.$$

Thus we have

$$\begin{aligned}
&\sum_{\substack{(k,\alpha) \in \mathbb{Z}^{1+2} \\ \geq 0}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \left\| K^k J^\alpha (|u|^{2\sigma} u) \right\|_{L^{3/2}(\mathbb{R}; H_{6/5}^1 \cap A_{6/5}^1)} \\
&\leq \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+2}} \sum_{\substack{k_1+k_2=k \\ k_2 \geq 1}} \sum_{\beta_1+\beta_2+\dots+\beta_{2\sigma+1}=\alpha} \left\{ \frac{a_0^k}{k_1!k_2!} \frac{a_1^{\alpha_1}}{\beta_1!\beta_2!\cdots\beta_{2\sigma+1}!} \right. \\
&\quad \times \left. \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}(\mathbb{R}; H_{6/5}^1 \cap A_{6/5}^1)} \right\}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+2}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \left\{ \frac{a_0^k}{k!} \frac{a_1^\alpha}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \right. \\
& \quad \times \left. \left\| \mathcal{L}^k \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^{3/2}(\mathbb{R}; H_{6/5}^1 \cap A_{6/5}^1)} \right\} \\
& \leq C(2(2\sigma - 1))^{\sigma - 1/2} \\
& \quad \times \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+2}} \sum_{\substack{k_1 + k_2 = k \\ k_2 \geq 1}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \sum_{l_1 + l_2 + \dots + l_{2\sigma+1} = k_1} \left\{ \frac{(2a_0(4\sigma - 1))^{k_2}}{k_2!} \right. \\
& \quad \times \chi_{K_\sigma}(k_2) \frac{a_1^\alpha}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \frac{a_0^{k_1}}{l_1! l_2! \dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^1(\mathbb{R}) \cap X_1(\mathbb{R})} \left. \right\} \\
& \quad + C(2(2\sigma - 1))^{\sigma - 1/2} \\
& \quad \times \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+2}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \sum_{l_1 + l_2 + \dots + l_{2\sigma+1} = k} \left\{ \frac{a_1^\alpha}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \frac{a_0^k}{l_1! l_2! \dots l_{2\sigma+1}!} \right. \\
& \quad \times \left. \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^1(\mathbb{R}) \cap X_1(\mathbb{R})} \right\} \\
& \leq (2(2\sigma - 1))^{\sigma - 1/2} \left(\sum_{k_2=1}^{\infty} \frac{(2a_0(4\sigma - 1))^{k_2}}{k_2!} \chi_{K_\sigma}(k_2) + 1 \right) \|u\|_{G^a(\mathbb{R})}^{2\sigma+1} \\
& \leq C(2(2\sigma - 1))^{\sigma - 1/2} e^{2a_0(4\sigma - 1)} \|u\|_{G^a(\mathbb{R})}^{2\sigma+1} \\
& \leq C(2(2\sigma - 1))^{\sigma - 1/2} e^{2a_0(4\sigma - 1)} R^{2\sigma+1} \\
& = C(4\sigma - 2)^{\sigma - 1/2} e^{2a_0(4\sigma - 1)} R^{2(\sigma - 1)} R^3
\end{aligned}$$

and

$$\sum_{\sigma=1}^{\infty} \frac{|\lambda|^\sigma}{\sigma!} \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+2}} \frac{a^{(k,\alpha)}}{(k, \alpha)!} \left\| K^k J^\alpha (|u|^{2\sigma} u) \right\|_{L^{3/2}(\mathbb{R}; H_{6/5}^1 \cap A_{6/5}^1)} \leq CF(R) R^3$$

where

$$F(R) = \sum_{\sigma=1}^{\infty} \frac{|\lambda|^\sigma}{\sigma!} (4\sigma - 2)^{\sigma - 1/2} e^{2a_0(4\sigma - 1)} R^{2(\sigma - 1)}$$

converges for

$$0 < R < (4e^{4a_0+1} |\lambda|)^{-1/2}$$

by the d'Alembert ratio test. Similarly we obtain

$$d(\Phi u, \Phi v) \leq CF(R) R^2 d(u, v).$$

Therefore we have

$$\|\Phi u\|_{G^a(\mathbb{R})} \leq C\rho + CF(R)R^3, \quad d(\Phi u, \Phi v) \leq CF(R)R^2d(u, v).$$

Then we obtain the unique global solution

$$u = \Phi u$$

as a fixed point of $\Phi : B_R^a \rightarrow B_R^a$ with R and ρ which satisfy

$$CF(R)R^2 \leq \frac{1}{2}, \quad 0 < R < (4e^{4a_0+1}|\lambda|)^{-1/2}$$

and

$$\rho = \frac{R}{2C}.$$

Case 2: $n \geq 3$

Let $n \geq 3$. By the Kato–Ponce inequality we have ([4, 16, 17])

$$\left\| \prod_{j=1}^{2\sigma+1} v_j \right\|_{H_{2n/(n+2)}^{n/2}} \leq C \sum_{j=1}^{2\sigma+1} \|v_j\|_{H_{2n/(n-2)}^{n/2}} \left\| \prod_{\mu \neq j} v_j \right\|_{L^{n/2}}$$

and

$$\left\| \prod_{j=1}^{2\sigma+1} v_j \right\|_{A_{2n/(n+2)}^{n/2}(t)} \leq C \sum_{j=1}^{2\sigma+1} \|v_j\|_{A_{2n/(n-2)}^{n/2}(t)} \left\| \prod_{\mu \neq j} v_j \right\|_{L^{n/2}}.$$

From this inequalities and the critical Sobolev inequality (2.1) in Lemma 2, we estimate

$$\begin{aligned} & \left\| \left(\mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right)(t) \right\|_{H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2}(t)} \\ & \leq \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \left\| \prod_{j=1}^{2\sigma+1} (K^{l_j} J^{\beta_j} u(t))^{[\varepsilon_j]} \right\|_{H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2}(t)} \\ & \leq C \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \\ & \quad \times \left(\sum_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u\|_{H_{2n/(n-2)}^{n/2} \cap A_{2n/(n-2)}^{n/2}(t)} \left\| \prod_{\mu \neq j} K^{l_\mu} J^{\beta_\mu} u \right\|_{L^{n/2}} \right) \\ & \leq C \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \\ & \quad \times \left(\sum_{j=1}^{2\sigma+1} \|K^{l_j} J^{\beta_j} u(t)\|_{H_{2n/(n-2)}^{n/2} \cap A_{2n/(n-2)}^{n/2}(t)} \prod_{\mu \neq j} \|K^{l_\mu} J^{\beta_\mu} u(t)\|_{L^{n/2}} \right) \end{aligned}$$

$$\leq C(n\sigma)^\sigma \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \\ \times \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u(t) \right\|_{H_{2n/(n-2)}^{n/2} \cap A_{2n/(n-2)}^{n/2}(t)} \prod_{\mu \neq j} \left\| K^{l_\mu} J^{\beta_\mu} u(t) \right\|_{H^{n/2}} \right)$$

for $|t| \leq 1$, and by using the critical Sobolev inequality (2) in Lemma 2, we estimate

$$\left\| \left(\mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right)(t) \right\|_{H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2}(t)} \\ \leq C(n\sigma)^\sigma |t|^{-n\sigma+2} \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \\ \times \left(\sum_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u(t) \right\|_{H_{2n/(n-2)}^{n/2} \cap A_{2n/(n-2)}^{n/2}(t)} \prod_{\mu \neq j} \left\| K^{l_\mu} J^{\beta_\mu} u(t) \right\|_{A_2^{n/2}(t)} \right)$$

for $|t| \geq 1$. Therefore by the above estimates with (3.1), we have

$$\left\| \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2([-1,1]; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \\ = \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2([-1,1]; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \\ \leq C(n\sigma)^\sigma (2n\sigma - 1)^{k_2} 2^{k_2} \chi_{K_\sigma}(k_2) \\ \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \chi_{K_\sigma}(k_2) \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}([-1,1]) \cap X_{n/2}([-1,1])} \\ \leq C(n\sigma)^\sigma (2n\sigma - 1)^{k_2} 2^{k_2} \chi_{K_\sigma}(k_2) \\ \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}([-1,1]) \cap X_{n/2}([-1,1])}$$

and

$$\left\| \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2(\mathbb{R} \setminus [-1,1]; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \\ = \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2(\mathbb{R} \setminus [-1,1]; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \\ \leq C(n\sigma)^\sigma (2n\sigma - 1)^{k_2} 2^{k_2} \chi_{K_\sigma}(k_2) \\ \times \sum_{l_1+l_2+\dots+l_{2\sigma+1}=k_1} \frac{k_1!}{l_1!l_2!\dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}(\mathbb{R} \setminus [-1,1]) \cap X_{n/2}(\mathbb{R} \setminus [-1,1])}$$

$$\leq C(n\sigma)^\sigma (2n\sigma - 1)^{k_2} 2^{k_2} \chi_{K_\sigma}(k_2) \\ \times \sum_{l_1 + l_2 + \dots + l_{2\sigma+1} = k_1} \frac{k_1!}{l_1! l_2! \dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}(\mathbb{R} \setminus [-1, 1]) \cap X_{n/2}(\mathbb{R} \setminus [-1, 1])}$$

where we have used the estimates

$$\left| \mathcal{R}_\sigma^{k_2}(t) \right| \leq 2^{k_2} (2n\sigma - 1)^{k_2} \quad \text{for } |t| < 1$$

and

$$\left| \mathcal{R}_\sigma^{k_2}(t) \right| \leq 2^{k_2} (2n\sigma - 1)^{k_2} |t|^{n\sigma-2} \quad \text{for } |t| \geq 1.$$

Thus we have

$$\begin{aligned} & \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k, \alpha)}}{(k, \alpha)!} \left\| K^k J^\alpha (|u|^{2\sigma} u) \right\|_{L^2(\mathbb{R}; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \\ & \leq \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \sum_{\substack{k_1 + k_2 = k \\ k_2 \geq 1}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \left\{ \frac{a_0^k}{k_1! k_2!} \frac{a_1^{\alpha_1}}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \right. \\ & \quad \times \left. \left\| \chi_{K_\sigma}(k_2) \mathcal{R}_\sigma^{k_2} \mathcal{L}^{k_1} \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2(\mathbb{R}; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \right\} \\ & \quad + \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \left\{ \frac{a_0^k}{k!} \frac{a_1^{\alpha}}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \right. \\ & \quad \times \left. \left\| \mathcal{L}^k \left[\prod_{j=1}^{2\sigma+1} (J^{\beta_j} u)^{[\varepsilon_j]} \right] \right\|_{L^2(\mathbb{R}; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \right\} \\ & \leq C(n\sigma)^\sigma \\ & \quad \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \sum_{\substack{k_1 + k_2 = k \\ k_2 \geq 1}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \sum_{l_1 + l_2 + \dots + l_{2\sigma+1} = k_1} \left\{ \frac{(2a_0(2n\sigma - 1))^{k_2}}{k_2!} \chi_{K_\sigma}(k_2) \right. \\ & \quad \times \frac{a_1^{\alpha}}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \frac{a_0^{k_1}}{l_1! l_2! \dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}(\mathbb{R}) \cap X_{n/2}(\mathbb{R})} \left. \right\} \\ & \quad + C(n\sigma)^\sigma \sum_{(k, \alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \sum_{\beta_1 + \beta_2 + \dots + \beta_{2\sigma+1} = \alpha} \sum_{l_1 + l_2 + \dots + l_{2\sigma+1} = k} \left\{ \frac{a_1^{\alpha}}{\beta_1! \beta_2! \dots \beta_{2\sigma+1}!} \right. \\ & \quad \times \frac{a_0^k}{l_1! l_2! \dots l_{2\sigma+1}!} \prod_{j=1}^{2\sigma+1} \left\| K^{l_j} J^{\beta_j} u \right\|_{X^{n/2}(\mathbb{R}) \cap X_{n/2}(\mathbb{R})} \left. \right\} \\ & \leq (n\sigma)^\sigma \left(\sum_{k_2=1}^{\infty} \frac{(2a_0(2n\sigma - 1))^{k_2}}{k_2!} \chi_{K_\sigma}(k_2) + 1 \right) \|u\|_{G^a(\mathbb{R})}^{2\sigma+1} \end{aligned}$$

$$\begin{aligned}
&\leq C(n\sigma)^\sigma e^{2a_0(2n\sigma-1)} \|u\|_{G^a(\mathbb{R})}^{2\sigma+1} \\
&\leq C(n\sigma)^\sigma e^{2a_0(2n\sigma-1)} R^{2\sigma+1} \\
&= C(n\sigma)^\sigma e^{2a_0(2n\sigma-1)} R^{2(\sigma-1)} R^3
\end{aligned}$$

and

$$\sum_{\sigma=1}^{\infty} \frac{|\lambda|^\sigma}{\sigma!} \sum_{(k,\alpha) \in \mathbb{Z}_{\geq 0}^{1+n}} \frac{a^{(k,\alpha)}}{(k,\alpha)!} \left\| K^k J^\alpha (|u|^{2\sigma} u) \right\|_{L^2(\mathbb{R}; H_{2n/(n+2)}^{n/2} \cap A_{2n/(n+2)}^{n/2})} \leq CF(R) R^3$$

where

$$F(R) = \sum_{\sigma=1}^{\infty} \frac{|\lambda|^\sigma}{\sigma!} (n\sigma)^\sigma e^{2a_0(2n\sigma-1)} R^{2(\sigma-1)}$$

converges for

$$0 < R < (ne^{4a_0 n+1} |\lambda|)^{-1/2}$$

by the d'Alembert ratio test. Similarly we obtain

$$d(\Phi u, \Phi v) \leq CF(R) R^2 d(u, v).$$

Therefore we have

$$\|\Phi u\|_{G^a(\mathbb{R})} \leq C\rho + CF(R) R^3, \quad d(\Phi u, \Phi v) \leq CF(R) R^2 d(u, v).$$

Then we obtain the unique global solution

$$u = \Phi u$$

as a fixed point of $\Phi : B_R^a \rightarrow B_R^a$ with R and ρ satisfying

$$CF(R) R^2 \leq \frac{1}{2}, \quad 0 < R < (ne^{4a_0 n+1} |\lambda|)^{-1/2}$$

and

$$\rho = \frac{R}{2C}.$$

This completes the proof of Theorem 1. \square

Acknowledgements. This work was supported by JSPS KAKENHI Grant Number 19K14570.

REFERENCES

- [1] T. CAZENAVE, *Semilinear Schrödinger equations*, Courant Lect. Notes Math., vol. 10, Amer. Math. Soc., 2003.
- [2] A. DEBOUARD, *Analytic solution to non elliptic nonlinear Schrödinger equations*, J. Differ. Equ., **104**, (1993), 196–213.
- [3] J. GINIBRE, T. OZAWA AND G. VELO, *On the existence of the wave operators for a class nonlinear Schrödinger equations*, Ann. Inst. Henri Poincaré Phys. Théor., **60**, (1994), 211–239.
- [4] L. GRAFAKOS AND S. OH, *The Kato–Ponce inequality*, Commun. Partial Differ. Equ. **39**, (2014), 1128–1157.
- [5] N. HAYASHI AND K. KATO, *Analyticity in time and smoothing effect of solutions to nonlinear Schrödinger equations*, Comm. Math. Phys., **184**, (1997), 273–300.
- [6] N. HAYASHI, K. NAKAMITSU AND M. TSUTSUMI, *On solutions of the initial value problem for the nonlinear Schrödinger equations in one space dimension*, Math. Z., **192**, (1986), 637–650.
- [7] N. HAYASHI AND P. I. NAUMKIN, *Asymptotics for large time of solutions to the nonlinear Schrödinger and Hartree equations*, Amer. J. Math., **120**, (1998), 369–389.
- [8] N. HAYASHI AND T. OZAWA, *Scattering theory in the weighted $L^2(\mathbb{R}^n)$ space for some Schrödinger equations*, Ann. Inst. H. Poincaré Phys. Théor., **48**, (1988), 17–37.
- [9] N. HAYASHI AND S. SAITO, *Analyticity and smoothing effect for the Schrödinger equation*, Ann. Inst. H. Poincaré Phys. Théor., **52**, (1990), 163–173.
- [10] G. HOSHINO, *Space-time analytic smoothing effect for a system of nonlinear Schrödinger equations with non pseudo-conformally invariant interactions*, Comm. Partial Differential Equations, **42**, (2017), 802–819.
- [11] G. HOSHINO, *Space-time Gevrey smoothing effect for the dissipative nonlinear Schrödinger equations*, Nonlinear Differ. Equ. Appl., **27**, (2020), no. 32.
- [12] G. HOSHINO AND T. OZAWA, *Analytic smoothing effect for nonlinear Schrödinger equation in two space dimensions*, Osaka J. Math., **51**, (2014), 609–618.
- [13] G. HOSHINO AND T. OZAWA, *Analytic smoothing effect for nonlinear Schrödinger equation with quintic nonlinearity*, J. Math. Anal. Appl., **419**, (2014), 285–297.
- [14] G. HOSHINO AND T. OZAWA, *Space-time analytic smoothing effect for the pseudo-conformally invariant Schrödinger equations*, Nonlinear Differ. Equ. Appl., **23**, (2016), no. 3.
- [15] S. IBRAHIM, M. MAJDOUB, N. MASMOUDI AND K. NAKANISHI, *Scattering for the two-dimensional NLS with exponential nonlinearity*, Nonlinearity, **25**, (2012), 1843–1849.
- [16] T. KATO, *On nonlinear Schrödinger equations. II. H^s -solutions and unconditional well-posedness*, J. Anal. Math., **67**, (1995), 281–306.
- [17] T. KATO AND G. PONCE, *Commutator estimates and the Euler and NavierStokes equations*, Comm. Pure App. Math., **41**, (1988), 891–907.
- [18] M. KEEL AND T. TAO, *Endpoint Strichartz inequalities*, Amer. J. Math., **120**, (1998), 955–980.
- [19] H. KOZONO, T. OGAWA AND H. SOHR, *Asymptotic behavior in L^r for weak solutions of the Navier–Stokes equations in exterior domains*, Manuscripta Math. **74**, (1992), 253–273.
- [20] J. F. LAM, B. LIPPmann AND F. TAPPERT, *Self-trapped laser beams in plasma*, The Physics of Fluids, **20**, (1977), 1176–1179.
- [21] K. NAKAMITSU, *Analytic finite energy solutions of the nonlinear Schrödinger equation*, Comm. Math. Phys., **260**, (2005), 117–130.
- [22] M. NAKAMURA AND T. OZAWA, *Nonlinear Schrödinger equations in the Sobolev space of critical order*, J. Funct. Anal., **155**, (1998), 364–380.
- [23] T. OZAWA, *On critical cases of Sobolev’s inequalities*, J. Funct. Anal., **127**, (1995), 259–269.
- [24] T. OZAWA AND K. YAMAUCHI, *Analytic smoothing effect for global solutions to nonlinear Schrödinger equation*, J. Math. Anal. Appl., **364**, (2010), 492–497.
- [25] B. WANG, *The smoothness of scattering operator for sinh-Gordon and nonlinear Schrödinger equations*, Acta Math. Sinica, English Series, **18**, (2002), 549–564.

- [26] S. WANG, *Energy scattering for Schrödinger equation with exponential nonlinearity in two dimensions*, J. Funct. Space Appl., **2013**, 968603, 13 pp.
- [27] K. YAJIMA, *Existence of solutions for Schrödinger evolution equations*, Comm. Math. Phys., **110**, (1987), 415–426.

(Received June 3, 2022)

Gaku Hoshino
Tokyo Denki University
Faculty of Science and Engineering
Division of Science, Hatoyama, Saitama, 350-0394, Japan
e-mail: g.hoshino@mail.dendai.ac.jp