

GENERALIZED FRACTIONAL OSTROWSKI TYPE INEQUALITIES VIA $\phi - \lambda$ -CONVEX FUNCTION

ALI HASSAN* AND ASIF RAZA KHAN

(Communicated by S. S. Dragomir)

Abstract. In this paper, we would like to state well-known Ostrowski inequality via generalized Montgomery identity for the $\phi - \lambda$ -convex function. Also, we would like to state the generalization of the classical Ostrowski inequality via generalized fractional integrals, which is obtained for functions whose first derivative in absolute values is $\phi - \lambda$ -convex. Moreover we establish some Ostrowski type inequalities via generalized fractional integrals and their particular cases for the class of functions whose derivatives in absolute values at certain powers are $\phi - \lambda$ -convex by using different techniques including Hölder's inequality and power mean inequality. Also, standard results would be capture as special cases. Moreover, some applications in terms of special means would also be given.

1. Introduction

In recent years, the generalization of classical convex function have emerged resulting in applications in the field of Mathematics. From literature, we recall some definitions for different types of convex (concave).

DEFINITION 1. [2] The $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is said to be convex (concave), if

$$\eta(tx + (1-t)y) \leqslant (\geqslant) t\eta(x) + (1-t)\eta(y),$$

$\forall x, y \in I, t \in [0, 1]$.

DEFINITION 2. [2] The $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is said to be *MT*-convex (concave), if $\eta(x) \geqslant 0$ and

$$\eta(tx + (1-t)y) \leqslant (\geqslant) \frac{\sqrt{t}}{2\sqrt{1-t}}\eta(x) + \frac{\sqrt{1-t}}{2\sqrt{t}}\eta(y),$$

$\forall x, y \in I, t \in [0, 1]$.

Mathematics subject classification (2020): 26A33, 26A51, 26D15, 26D99, 47A30, 33B10.

Keywords and phrases: Ostrowski inequality, convex, power mean inequality, Hölder's inequality.

* Corresponding author.

DEFINITION 3. [15] The $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is a P -convex (concave), if $\eta(x) \geq 0$ and $\forall x, y \in I$ and $t \in [0, 1]$ we have

$$\eta(tx + (1-t)y) \leqslant (\geqslant) \eta(x) + \eta(y).$$

DEFINITION 4. [17] The $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is a Godunova-Levin convex (concave), if $\eta(x) \geq 0$ and $\forall x, y \in I$ and $t \in (0, 1)$ we have

$$\eta(tx + (1-t)y) \leqslant (\geqslant) \frac{1}{t}\eta(x) + \frac{1}{1-t}\eta(y).$$

DEFINITION 5. [3] Let $s \in [0, 1]$. The $\eta : I \subset [0, \infty) \rightarrow \mathbb{R}$ is said to be s -convex (concave) in the 2^{nd} kind, if

$$\eta(tx + (1-t)y) \leqslant (\geqslant) t^s\eta(x) + (1-t)^s\eta(y),$$

$$\forall x, y \in I, t \in [0, 1].$$

DEFINITION 6. [9] The $\eta : I \subset \mathbb{R} \rightarrow [0, \infty)$ is of Godunova-Levin s -convex (concave), with $s \in [0, 1]$, if

$$\eta(tx + (1-t)y) \leqslant (\geqslant) \frac{1}{t^s}\eta(x) + \frac{1}{(1-t)^s}\eta(y),$$

$$\forall t \in (0, 1) \text{ and } x, y \in I.$$

DEFINITION 7. [23] Let $h : J \subseteq \mathbb{R} \rightarrow [0, \infty)$ with $h \neq 0$. The $\eta : I \subseteq \mathbb{R} \rightarrow [0, \infty)$ is an h -convex (concave) if $\forall x, y \in I$, we have

$$\eta(tx + (1-t)y) \leqslant (\geqslant) h(t)\eta(x) + h(1-t)\eta(y),$$

$$\forall t \in [0, 1].$$

DEFINITION 8. [10] Let $\phi : (0, 1) \rightarrow (0, \infty)$. The $\eta : I \rightarrow [0, \infty)$ is a ϕ -convex (concave) on the interval I if for all $x, y \in I$ we have

$$\eta(tx + (1-t)y) \leqslant (\geqslant) t\phi(t)\eta(x) + (1-t)\phi(1-t)\eta(y),$$

$$\forall t \in (0, 1).$$

In almost every field of science, inequalities play an important role. Although it is very vast discipline but our focus is mainly on Ostrowski type inequalities. In 1938, Ostrowski established the following interesting integral inequality for differentiable mappings with bounded derivatives. This inequality is well known in the literature as Ostrowski inequality.

THEOREM 1. [20] Let $\varphi : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be differentiable on (ρ_a, ρ_b) with the property that $|\varphi'(t)| \leq M$ for all $t \in (\rho_a, \rho_b)$. Then

$$\left| \varphi(x) - \frac{1}{\rho_b - \rho_a} \int_{\rho_a}^{\rho_b} \varphi(t) dt \right| \leq (\rho_b - \rho_a) M \left[\frac{1}{4} + \left(\frac{x - \frac{\rho_a + \rho_b}{2}}{\rho_b - \rho_a} \right)^2 \right], \quad (1)$$

$\forall x \in (\rho_a, \rho_b)$.

Ostrowski inequality has applications in numerical integration, probability and optimization theory, statistics, information and integral operator theory. Until now, a large number of research papers and books have been written on generalizations of Ostrowski inequalities and their numerous applications in [6]–[14].

DEFINITION 9. The Riemann-Liouville integral operator of order $\zeta > 0$ with $\rho_a \geq 0$ is defined as

$$\begin{aligned} J_{\rho_a}^{\zeta} \varphi(x) &= \frac{1}{\Gamma(\zeta)} \int_{\rho_a}^x (x-t)^{\zeta-1} \varphi(t) dt, \\ J_{\rho_a}^0 \varphi(x) &= \varphi(x). \end{aligned} \quad (2)$$

In case of $\zeta = 1$, the fractional integral reduces to the classical integral.

DEFINITION 10. [21] The Riemann-Liouville integrals $I_{\rho_a^+}^{\zeta} \varphi$ and $I_{\rho_b^-}^{\zeta} \varphi$ of $\varphi \in L_1([\rho_a, \rho_b])$ having order $\zeta > 0$ with $\rho_a \geq 0$, $\rho_a < \rho_b$ are defined by

$$I_{\rho_a^+}^{\zeta} \varphi(x) = \frac{1}{\Gamma(\zeta)} \int_{\rho_a}^x (x-t)^{\zeta-1} \varphi(t) dt, \quad x > \rho_a$$

and

$$I_{\rho_b^-}^{\zeta} \varphi(x) = \frac{1}{\Gamma(\zeta)} \int_x^{\rho_b} (t-x)^{\zeta-1} \varphi(t) dt, \quad x < \rho_b,$$

respectively. Here $\Gamma(\zeta) = \int_0^\infty e^{-u} u^{\zeta-1} du$ is the Gamma function and $I_{\rho_a^+}^0 \varphi(x) = I_{\rho_b^-}^0 \varphi(x) = \varphi(x)$.

DEFINITION 11. [21] Let $g : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be an increasing and positive function on $(\rho_a, \rho_b]$, having a continuous derivative $g'(x)$ on (ρ_a, ρ_b) . The fractional integrals $I_{\rho_a^+, g}^{\zeta} \varphi$ and $I_{\rho_b^-, g}^{\zeta} \varphi$ of φ with respect to the function g on $[\rho_a, \rho_b]$ of order $\zeta > 0$ are defined by

$$I_{\rho_a^+, g}^{\zeta} \varphi(x) = \frac{1}{\Gamma(\zeta)} \int_{\rho_a}^x (g(x) - g(t))^{\zeta-1} g'(t) \varphi(t) dt, \quad x > \rho_a$$

and

$$I_{\rho_b^-, g}^{\zeta} \varphi(x) = \frac{1}{\Gamma(\zeta)} \int_x^{\rho_b} (g(t) - g(x))^{\zeta-1} g'(t) \varphi(t) dt, \quad x < \rho_b,$$

respectively.

REMARK 1. If we replace $g(x) = x$ the above fractional integrals reduce to the Riemann-Liouville fractional integrals.

THEOREM 2. [21] Let $\varphi : I \rightarrow \mathbb{R}$ be differentiable mapping on I^0 , with $\rho_a, \rho_b \in I$, $\rho_a < \rho_b$ $\varphi' \in L_1[\rho_a, \rho_b]$ and for $\zeta > 1$, Montgomery identity for fractional integrals holds:

$$\varphi(x) = \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) - J_{\rho_a}^{\zeta-1}(P_1(x, \rho_b) \varphi(\rho_b)) + J_{\rho_a}^\zeta(P_1(x, \rho_b) \varphi'(\rho_b)),$$

where $P_1(x, t)$ is the fractional Peano Kernel defined by:

$$P_1(x, t) = \begin{cases} \frac{t - \rho_a}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} \Gamma(\zeta), & \text{if } t \in [\rho_a, x], \\ \frac{t - \rho_b}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} \Gamma(\zeta), & \text{if } t \in (x, \rho_b]. \end{cases}$$

THEOREM 3. [21] Let $\varphi : I \rightarrow \mathbb{R}$ be differentiable mapping on I^0 , with $\rho_a, \rho_b \in I$, $\rho_a < \rho_b$ $\varphi' \in L_1[\rho_a, \rho_b]$ and for $\zeta > 1$, Generalized Montgomery identity for fractional integrals holds:

$$(1 - \varepsilon)\varphi(x) = \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) - J_{\rho_a}^{\zeta-1}(P_2(x, \rho_b) \varphi(\rho_b)) - \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) + J_{\rho_a}^\zeta(P_2(x, \rho_b) \varphi'(\rho_b)), \quad (3)$$

where $P_2(x, t)$ is the fractional Peano Kernel defined by:

$$P_2(x, t) = \begin{cases} \frac{t - \mu}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} \Gamma(\zeta), & \text{if } t \in [\rho_a, x], \\ \frac{t - \nu}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} \Gamma(\zeta), & \text{if } t \in (x, \rho_b]. \end{cases}$$

$$\forall x \in [\mu, \nu] \text{ for } \mu = \rho_a + \varepsilon \frac{\rho_b - \rho_a}{2} \text{ and } \nu = \rho_b - \varepsilon \frac{\rho_b - \rho_a}{2}.$$

Throughout this paper, we will assume that $g : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ is an increasing and positive function on $[\rho_a, \rho_b]$, having a continuous derivative $g'(x)$ on (ρ_a, ρ_b) . In order to prove our results, we need the following Lemma.

LEMMA 1. [18] Let $\varphi : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be a differentiable mapping on (ρ_a, ρ_b) with $a < b$. If $\varphi' : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be integrable on $[\rho_a, \rho_b]$. Then the identity for fractional

integrals holds with respect to another function

$$\begin{aligned} & \varphi(x) - \Gamma(\zeta + 1) \left[\frac{I_{x^+, g}^\zeta \varphi(\rho_b)}{2(g(\rho_b) - g(x))^\zeta} + \frac{I_{x^-, g}^\zeta \varphi(\rho_a)}{2(g(x) - g(\rho_a))^\zeta} \right] \\ &= \frac{x - \rho_a}{2(g(x) - g(\rho_a))^\zeta} \int_0^1 (g(tx + (1-t)\rho_a) - g(\rho_a))^\zeta \varphi'(tx + (1-t)\rho_a) dt \\ &\quad - \frac{\rho_b - x}{2(g(\rho_b) - g(x))^\zeta} \int_0^1 (g(\rho_b) - g(tx + (1-t)\rho_b))^\zeta \varphi'(tx + (1-t)\rho_b) dt. \end{aligned}$$

Throughout this paper, we denote

$$\begin{aligned} {}_{\bar{g}}^{\zeta} K_{\rho_a}^{\rho_b}(x) &= \left[\frac{(x - \rho_a)^{\zeta+1}}{2(g(x) - g(\rho_a))^\zeta} + \frac{(\rho_b - x)^{\zeta+1}}{2(g(\rho_b) - g(x))^\zeta} \right], \\ {}_{\phi, g}^{\zeta} \theta_{\rho_a}^{\rho_b}(x) &= \varphi(x) - \Gamma(\zeta + 1) \left[\frac{I_{x^+, g}^\zeta \varphi(\rho_b)}{2(g(\rho_b) - g(x))^\zeta} + \frac{I_{x^-, g}^\zeta \varphi(\rho_a)}{2(g(x) - g(\rho_a))^\zeta} \right]. \end{aligned}$$

We also make use of the Euler's beta function, which is for $x, y > 0$ defined as

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

The main aim of our study is to present Ostrowski inequality for fractional integrals with respect to another function, which is the generalization of the classical Ostrowski inequality (1) via $\phi - \lambda$ -convex. Moreover we establish some Ostrowski type inequalities for the class of functions whose derivatives in absolute values at certain powers are $\phi - \lambda$ -convex by using different techniques including Hölder's inequality [25] and power mean inequality [24]. Also we give the special cases of our results and applications of midpoint inequalities in special means. In the last section gives us conclusion with some remarks and future ideas to generalize the results.

2. Generalized Fractional Ostrowski inequality via $\phi - \lambda$ -convex

In this section, we are introducing very first time the concept of $\phi - \lambda$ -convex function, which contain many classes of convex in literature.

DEFINITION 12. Let $\lambda \in (0, 1]$ and $\phi : (0, 1) \rightarrow (0, \infty)$, The $\eta : I \rightarrow [0, \infty)$ is a $\phi - \lambda$ -convex (concave) on the interval I if $\forall x, y \in I$ we have

$$\eta(tx + (1-t)y) \leq (\geq) t^\lambda \phi(t) \eta(x) + (1-t)^\lambda \phi(1-t) \eta(y), \quad (4)$$

$$\forall t \in (0, 1).$$

REMARK 2. In Definition 12,

1. If $\lambda = 1$ in (4), we get ϕ -convex (concave).
2. If $\lambda = 1$, $l(t) = t$, and by taking $h = l\phi$ in (4), we get h -convex (concave).
3. If $\lambda = 1$, $\phi(t) = \frac{1}{t^{s+1}}$ with $s \in [0, 1]$ in (4), then we get the class of GL s -convex (concave).
4. If $\lambda = 1$, $\phi(t) = \frac{1}{t^2}$ in (4), then we get the concept of GL convex (concave).
5. If $\lambda = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in (4), then we get the concept of s -convex (concave) in 2^{nd} kind.
6. If $\lambda = 1$, $\phi(t) = \frac{1}{t}$ in (4), then we get the concept of P -convex (concave).
7. If $\lambda = 1$, $\phi(t) = 1$ in (4), then we get the concept of ordinary convex (concave).
8. If $\lambda = 1$, $\phi(t) = \frac{1}{2\sqrt{t(1-t)}}$ in (4), then we get the concept of MT -convex (concave).

THEOREM 4. *If Lemma 1 hold. Additionally, assume that $\lambda \in (0, 1]$, $|\varphi'|$ is $\phi - \lambda$ -convex function on $[\rho_a, \rho_b]$ with $\phi(t) \neq \frac{1}{t^2}$ and $|\varphi'(x)| \leq M$, $|g'(x)| \leq L$, $x \in [\rho_a, \rho_b]$. Then for each $x \in (\rho_a, \rho_b)$ the following inequality holds:*

$$\left| \zeta_{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\int_0^1 \left[t^{\zeta+\lambda} \phi(t) + t^\zeta (1-t)^\lambda \phi(1-t) \right] dt \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x). \quad (5)$$

Proof. From the Lemma 1 we have

$$\begin{aligned} \left| \zeta_{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| &\leq \frac{x - \rho_a}{2(g(x) - g(\rho_a))^\zeta} \int_0^1 \frac{|\varphi'(tx + (1-t)\rho_a)|}{(g(tx + (1-t)\rho_a) - g(\rho_a))^{-\zeta}} dt \\ &\quad + \frac{\rho_b - x}{2(g(\rho_b) - g(x))^\zeta} \int_0^1 \frac{|\varphi'(tx + (1-t)\rho_b)|}{(g(\rho_b) - g(tx + (1-t)\rho_b))^{-\zeta}} dt. \end{aligned} \quad (6)$$

Since g is differentiable and $|g'(x)| \leq L$ on $[\rho_a, \rho_b]$, we get that g is Lipschizian function, i.e.

$$g(tx + (1-t)\rho_a) - g(\rho_a) \leq Lt(x - \rho_a) \quad (7)$$

$$g(\rho_b) - g(tx + (1-t)\rho_b) \leq Lt(\rho_b - x). \quad (8)$$

Using inequalities (7) and (8) in (6), we get

$$\begin{aligned} \left| \zeta_{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| &\leq L^\zeta \frac{(x - \rho_a)^{\zeta+1}}{2(g(x) - g(\rho_a))^\zeta} \int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_a)| dt \\ &\quad + L^\zeta \frac{(\rho_b - x)^{\zeta+1}}{2(g(\rho_b) - g(x))^\zeta} \int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_b)| dt. \end{aligned} \quad (9)$$

Since $|\varphi'|$ is $\phi - \lambda$ -convex on $[\rho_a, \rho_b]$ and $|\varphi'(x)| \leq M$, we have

$$\begin{aligned} & \int_0^1 t^\zeta \left[t^\lambda \phi(t) |\varphi'(x)| + (1-t)^\lambda \phi(1-t) |\varphi'(\rho_a)| \right] dt \\ & \leq M \int_0^1 t^\zeta \left[t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t) \right] dt. \end{aligned} \quad (10)$$

and similarly

$$\begin{aligned} & \int_0^1 t^\zeta \left[t^\lambda \phi(t) |\varphi'(x)| + (1-t)^\lambda \phi(1-t) |\varphi'(\rho_b)| \right] dt \\ & \leq M \int_0^1 t^\zeta \left[t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t) \right] dt. \end{aligned} \quad (11)$$

By using inequalities (10) and (11) in (9), we get

$$\left| \frac{\zeta}{\phi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\int_0^1 t^\zeta \left[t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t) \right] dt \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x). \quad \square$$

COROLLARY 1. In Theorem 4,

1. If $\lambda = 1$ in (5), then Fractional Ostrowski type inequality for ϕ -convex:

$$\left| \frac{\zeta}{\phi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\int_0^1 t^\zeta [t\phi(t) + (1-t)\phi(1-t)] dt \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x).$$

2. If $\lambda = 1$, $l(t) = t$ and $h = l\phi$ in (5), then Fractional Ostrowski type inequality for h -convex:

$$\left| \frac{\zeta}{\phi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\int_0^1 t^\zeta [h(t) + h(1-t)] dt \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x).$$

3. If $\lambda = 1$, $\phi(t) = \frac{1}{t^{s+1}}$ with $s \in [0, 1)$ in (5), then Ostrowski inequality for Godunova-Levin s -convex:

$$\left| \frac{\zeta}{\phi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\frac{1}{1 + \zeta - s} + \frac{\Gamma(1 + \zeta) \Gamma(1 - s)}{\Gamma(2 + \zeta - s)} \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x).$$

4. If $\lambda = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in (5), then Ostrowski inequality for s -convex in 2nd kind:

$$\left| \frac{\zeta}{\phi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\frac{1}{1 + \zeta + s} + \frac{\Gamma(1 + \zeta) \Gamma(1 + s)}{\Gamma(2 + \zeta + s)} \right) \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x).$$

5. If $\lambda = 1$, $g(x) = x$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (5), then inequality (2.6) of Theorem 7 in [22].

6. If $\lambda = 1$, $g(x) = x$, $\zeta = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (5), then inequality (2.1) of Theorem 2 in [1].

7. If $\lambda = 1$, $\phi(t) = \frac{1}{t}$ in inequality (5), then Ostrowski inequality for P -convex via fractional integrals:

$$\left| \int_{\rho_a}^{\rho_b} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{2ML^\zeta}{1+\zeta} \int_{\rho_a}^{\rho_b} K_{\rho_a}^{\rho_b}(x)$$

8. If $\lambda = \phi(t) = 1$ in inequality (5), then Ostrowski inequality for convex via fractional integrals:

$$\left| \int_{\rho_a}^{\rho_b} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{1+\zeta} \int_{\rho_a}^{\rho_b} K_{\rho_a}^{\rho_b}(x)$$

9. If $g(x) = x$, $\lambda = \phi(t) = 1$ in inequality (5), then Corollary 1 in [22].

10. If $g(x) = x$, $\lambda = \phi(t) = \zeta = 1$ in inequality (5), then one has inequality (1.3) of Theorem 3 in [22].

11. If $\lambda = 1$, $\phi(t) = \frac{1}{2\sqrt{t(1-t)}}$ in (5), then Fractional Ostrowski type inequality MT -convex:

$$\left| \int_{\rho_a}^{\rho_b} \theta_{\rho_a}^{\rho_b}(x) \right| \leq ML^\zeta \left(\frac{\sqrt{\pi} \Gamma[\frac{1}{2} + \zeta]}{2 \Gamma[1 + \zeta]} \right) \int_{\rho_a}^{\rho_b} K_{\rho_a}^{\rho_b}(x).$$

THEOREM 5. If Lemma 1 hold. Additionally, assume that $\lambda \in (0, 1]$, $|\varphi'|^q$ is $\phi - \lambda$ -convex function on $[\rho_a, \rho_b]$, $q \geq 1$ with $\phi(t) \neq \frac{1}{t^2}$ and $|\varphi'(x)| \leq M$, $|g'(x)| \leq L$, $x \in [\rho_a, \rho_b]$. Then $\forall x \in (\rho_a, \rho_b)$ the following inequality holds:

$$\left| \int_{\rho_a}^{\rho_b} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(\zeta+1)^{1-\frac{1}{q}}} \int_0^1 \left[t^{\zeta+\lambda} \phi(t) + t^\zeta (1-t)^\lambda \phi(1-t) \right] dt^{\frac{1}{q}}. \quad (12)$$

Proof. From the inequality (9) and using power mean inequality [24], we have

$$\begin{aligned} \left| \int_{\rho_a}^{\rho_b} \theta_{\rho_a}^{\rho_b}(x) \right| &\leq L^\zeta \frac{(x - \rho_a)^{\zeta+1}}{2(g(x) - g(\rho_a))^\zeta} \int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_a)|^q dt \\ &+ L^\zeta \frac{(\rho_b - x)^{\zeta+1}}{2(g(\rho_b) - g(x))^\zeta} \int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_b)|^q dt \\ &\leq L^\zeta \frac{(x - \rho_a)^{\zeta+1}}{2(g(x) - g(\rho_a))^\zeta} \left(\int_0^1 t^\zeta dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_a)|^q dt \right)^{\frac{1}{q}} \\ &+ L^\zeta \frac{(\rho_b - x)^{\zeta+1}}{2(g(\rho_b) - g(x))^\zeta} \left(\int_0^1 t^\zeta dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_b)|^q dt \right)^{\frac{1}{q}}. \end{aligned} \quad (13)$$

Since $|\varphi'|^q$ is $\phi - \lambda$ -convex on $[\rho_a, \rho_b]$. and $|\varphi'(x)| \leq M$, we get

$$\int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_a)|^q dt \leq M^q \int_0^1 t^\zeta [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \quad (14)$$

and

$$\int_0^1 t^\zeta |\varphi'(tx + (1-t)\rho_b)|^q dt \leq M^q \int_0^1 t^\zeta [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt. \quad (15)$$

Using the inequalities (13)–(15), we get

$$\left| \frac{\zeta}{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(\zeta+1)^{1-\frac{1}{q}}} \left(\int_0^1 t^\zeta [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x). \quad \square$$

COROLLARY 2. In Theorem 5,

1. If $q = 1$, one has the Theorem 4.

2. If $\lambda = 1$ in (12), then Fractional Ostrowski type inequality for ϕ -convex:

$$\left| \frac{\zeta}{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \left(\int_0^1 t^\zeta [t\phi(t) + (1-t)\phi(1-t)] dt \right)^{\frac{1}{q}}.$$

3. If $\lambda = 1$, $l(t) = t$ and $h = l\phi$ in (12), then Fractional Ostrowski type inequality for h -convex:

$$\left| \frac{\zeta}{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \left(\int_0^1 t^\zeta [h(t) + h(1-t)] dt \right)^{\frac{1}{q}}.$$

4. If $\lambda = 1$, $\phi(t) = \frac{1}{t^{s+1}}$ with $s \in [0, 1)$ in (12), then Ostrowski inequality for Godunova-Levin s -convex:

$$\left| \frac{\zeta}{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \left(\frac{1}{1+\zeta-s} + \frac{\Gamma(1+\zeta)\Gamma(1-s)}{\Gamma(2+\zeta-s)} \right)^{\frac{1}{q}}.$$

5. If $\lambda = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (12), then Fractional Ostrowski type inequality for s -convex in 2nd kind:

$$\left| \frac{\zeta}{\varphi, g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta}{(\zeta+1)^{1-\frac{1}{q}}} \left(\frac{1}{1+\zeta+s} + \frac{\Gamma(1+\zeta)\Gamma(1+s)}{\Gamma(2+\zeta+s)} \right)^{\frac{1}{q}}.$$

6. If $\lambda = 1$, $g(x) = x$, and $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (12), then inequality (2.8) of Theorem 9 in [22].

7. If $g(x) = x$, $\lambda = \zeta = 1$ and $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (12), then inequality (2.3) of Theorem 4 in [1].
8. If $\lambda = 1$, $\phi(t) = \frac{1}{t}$ in inequality (12), then Ostrowski inequality for P -convex via fractional integrals:

$$\left| \zeta_{\phi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{2^{\frac{1}{q}} M L^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \zeta_g K_{\rho_a}^{\rho_b}(x).$$

9. If $\lambda = \phi(t) = 1$ in inequality (12), then Ostrowski inequality for convex via fractional integrals:

$$\left| \zeta_{\phi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{M L^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \zeta_g K_{\rho_a}^{\rho_b}(x).$$

10. If $g(x) = x$, $\lambda = \phi(t) = 1$ in inequality (12), then inequality of Corollary 3 in [22].
11. If $g(x) = x$, $\lambda = \phi(t) = \zeta = 1$ in inequality (12), then one has inequality (1.5) of Theorem 5 in [22].
12. If $\lambda = 1$, $\phi(t) = \frac{1}{2\sqrt{t(1-t)}}$ in (12), then Fractional Ostrowski type inequality MT -convex:

$$\left| \zeta_{\phi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{M L^\zeta}{(1+\zeta)^{1-\frac{1}{q}}} \left(\frac{\sqrt{\pi} \Gamma[\frac{1}{2} + \zeta]}{2 \Gamma[1 + \zeta]} \right)^{\frac{1}{q}} \zeta_g K_{\rho_a}^{\rho_b}(x).$$

THEOREM 6. If Lemma 1 hold. Additionally, assume that $\lambda \in (0, 1]$, $|\phi'|^q$ is $\phi - \lambda$ -convex function on $[\rho_a, \rho_b]$, $q > 1$ with $\phi(t) \neq \frac{1}{t^2}$ and $|\phi'(x)| \leq M$, $|g'(x)| \leq L$, $x \in [\rho_a, \rho_b]$. Then for each $x \in (\rho_a, \rho_b)$ the following inequality holds:

$$\left| \zeta_{\phi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{M L^\zeta \zeta_g K_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}} \left(\int_0^1 \left[t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t) \right] dt \right)^{\frac{1}{q}}, \quad (16)$$

where $p^{-1} + q^{-1} = 1$.

Proof. From the inequality (9) and and using Hölder's inequality [25], we have

$$\begin{aligned} \left| \zeta_{\phi,g} \theta_{\rho_a}^{\rho_b}(x) \right| &\leq L^\zeta \frac{(x - \rho_a)^{\zeta+1}}{2(g(x) - g(\rho_a))^\zeta} \left(\int_0^1 t^{\zeta p} dt \right)^{\frac{1}{p}} \left(\int_0^1 |\phi'(tx + (1-t)\rho_a)|^q dt \right)^{\frac{1}{q}} \\ &\quad + L^\zeta \frac{(\rho_b - x)^{\zeta+1}}{2(g(\rho_b) - g(x))^\zeta} \left(\int_0^1 t^{\zeta p} dt \right)^{\frac{1}{p}} \left(\int_0^1 |\phi'(tx + (1-t)\rho_b)|^q dt \right)^{\frac{1}{q}}. \end{aligned} \quad (17)$$

Since $|\varphi'|^q$ is $\phi - \lambda$ -convex and $|\varphi'(x)| \leq M$, we have

$$\int_0^1 |\varphi'(tx + (1-t)\rho_a)|^q dt \leq M^q \int_0^1 [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \quad (18)$$

and

$$\int_0^1 |\varphi'(tx + (1-t)\rho_b)|^q dt \leq M^q \int_0^1 [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt. \quad (19)$$

Using inequalities (17)–(19), we get

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^{\zeta} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}} \left(\int_0^1 [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}}. \quad \square$$

COROLLARY 3. In Theorem 6,

1. If $\lambda = 1$, $l(t) = t$ and $h = l\phi$ in (16), then Fractional Ostrowski type inequality for h -convex:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^{\zeta} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}} \left(\int_0^1 [h(t) + h(1-t)] dt \right)^{\frac{1}{q}}$$

2. If $\lambda = 1$, $\phi(t) = \frac{1}{t^{s+1}}$ with $s \in [0, 1]$ in (16), then Ostrowski inequality for Godunova-Levin s -convex:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^{\zeta} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}} \left(\frac{2}{1-s} \right)^{\frac{1}{q}}.$$

3. If $\lambda = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (16), then Fractional Ostrowski type inequality for s -convex in 2nd kind:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^{\zeta} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}} \left(\frac{2}{1+s} \right)^{\frac{1}{q}}.$$

4. If $\lambda = 1$, $g(x) = x$ and $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (16), then inequality (2.7) of Theorem 8 in [22].

5. If $\lambda = 1$, $g(x) = x$, $\zeta = 1$ and $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in inequality (16), then inequality (2.4) of Theorem 3 in [1].

6. If $\lambda = 1$, $\phi(t) = \frac{1}{t}$ in inequality (16), then Ostrowski inequality for P -convex via fractional integrals:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{2^{\frac{1}{q}} ML^{\zeta} \frac{\zeta}{g} \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}}.$$

7. If $\lambda = \phi(t) = 1$ in inequality (16), then Ostrowski inequality for convex via fractional integrals:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \frac{ML^\zeta \zeta g \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}}.$$

8. If $g(x) = x$, $\lambda = \phi(t) = 1$ in inequality (16), then one has Corollary 2 in [22].

9. If $g(x) = x$, $\lambda = \phi(t) = \zeta = 1$ in inequality (16), then one has inequality (1.4) of Theorem 4 in [22].

10. If $\lambda = 1$, $\phi(t) = \frac{1}{2\sqrt{t(1-t)}}$ in (16), then Fractional Ostrowski type inequality MT-convex:

$$\left| \zeta_{\varphi,g} \theta_{\rho_a}^{\rho_b}(x) \right| \leq \left(\frac{\pi}{2} \right)^{\frac{1}{q}} \frac{ML^\zeta \zeta g \kappa_{\rho_a}^{\rho_b}(x)}{(\zeta p + 1)^{\frac{1}{p}}}.$$

THEOREM 7. Let $\lambda \in (0,1]$, $\varphi : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be differentiable on (ρ_a, ρ_b) , $\varphi' : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be integrable on $[\rho_a, \rho_b]$ and $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$, be a $\phi - \lambda$ -convex (concave), then we have the inequalities

$$\begin{aligned} & \eta \left[(1-\varepsilon)\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leq (\geq) \frac{(x - \rho_a)^{\lambda-1} (\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)^\lambda} \phi \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \left[\int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right] \\ & \quad + \frac{(\rho_b - x)^{\lambda-\zeta}}{(\rho_b - \rho_a)^\lambda} \phi \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \left[\int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right], \end{aligned} \quad (20)$$

$\forall x \in [\mu, v]$ and $\varepsilon \in [0, 1]$.

Proof. Utilizing the generalized Montgomery identity (3) for fractional, we get

$$\begin{aligned} & (1-\varepsilon)\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \\ & + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \\ & = J_{\rho_a}^\zeta (P_2(x, \rho_b) \varphi'(\rho_b)) \\ & = \frac{1}{\Gamma(\zeta)} \int_{\rho_a}^{\rho_b} P_2(x, t) \frac{\varphi'(t)}{(\rho_b - t)^{1-\zeta}} dt \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \left[\frac{(\rho_b - x)^{1-\zeta}}{x - \rho_a} \int_{\rho_a}^x \frac{\{t - \mu\} \varphi'(t)}{(\rho_b - t)^{1-\zeta}} dt \right] \\
&\quad + \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \left[\frac{(\rho_b - x)^{1-\zeta}}{\rho_b - x} \int_x^{\rho_b} \frac{\{t - v\} \varphi'(t)}{(\rho_b - t)^{1-\zeta}} dt \right],
\end{aligned}$$

$\forall x \in [\mu, v]$ and $\varepsilon \in [0, 1]$. Next by using $\eta : I \subset [0, \infty) \rightarrow \mathbb{R}$, the $\phi - \lambda$ -convex (concave), we get

$$\begin{aligned}
&\eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\
&\quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\
&\leqslant (\geqslant) \left[\left(\frac{x - \rho_a}{\rho_b - \rho_a} \right)^\lambda \phi \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \eta \left[\frac{(\rho_b - x)^{1-\zeta}}{x - \rho_a} \int_{\rho_a}^x \frac{\{t - \mu\} \varphi'(t)}{(\rho_b - t)^{1-\zeta}} dt \right] \right. \\
&\quad \left. + \left[\left(\frac{\rho_b - x}{\rho_b - \rho_a} \right)^\lambda \phi \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \eta \left[\frac{(\rho_b - x)^{1-\zeta}}{\rho_b - x} \int_x^{\rho_b} \frac{\{t - v\} \varphi'(t)}{(\rho_b - t)^{1-\zeta}} dt \right] \right],
\right]
\end{aligned}$$

$\forall x \in [\mu, v]$ and $\varepsilon \in [0, 1]$. Applying Jensen's integral inequality [7], we get the inequality (20). \square

REMARK 3. In Theorem 7, If $\varepsilon = 0$, in (20) we get

$$\begin{aligned}
&\eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\
&\leqslant (\geqslant) \frac{(x - \rho_a)^{\lambda-1} (\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)^\lambda} \phi \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \left[\int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right] \\
&\quad + \frac{(\rho_b - x)^{\lambda-\zeta}}{(\rho_b - \rho_a)^\lambda} \phi \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \left[\int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right].
\end{aligned}$$

COROLLARY 4. In Theorem 7,

1. If $\lambda = 1$ in (20), then Fractional Ostrowski type inequality for ϕ -convex (concave):

$$\begin{aligned}
&\eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\
&\quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\
&\leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)} \left[\phi \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right. \\
&\quad \left. + \phi \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \tag{21}
\end{aligned}$$

REMARK 4. If $\varepsilon = 0$ in (21), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)} \left[\phi \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right. \\ & \quad \left. + \phi \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned}$$

2. If $\lambda = 1$, $l(t) = t$ and $h = l\phi$ in (20), then Fractional Ostrowski type inequality for h -convex (concave):

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) h \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \left[\frac{(\rho_b - x)^{1-\zeta}}{x - \rho_a} \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right] \\ & \quad + h \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \left[\frac{1}{(\rho_b - x)^{\zeta}} \int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned} \quad (22)$$

REMARK 5. If $\varepsilon = 0$ in (22), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) h \left(\frac{x - \rho_a}{\rho_b - \rho_a} \right) \left[\frac{(\rho_b - x)^{1-\zeta}}{x - \rho_a} \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right] \\ & \quad + h \left(\frac{\rho_b - x}{\rho_b - \rho_a} \right) \left[\frac{1}{(\rho_b - x)^{\zeta}} \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned}$$

3. If $\lambda = 1$, $\phi(t) = \frac{1}{t^{s+1}}$ with $s \in [0, 1]$ in (20), then Ostrowski inequality for Godunova-Levin s -convex (concave):

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - \rho_a)^s (\rho_b - x)^{1-\zeta}}{(x - \rho_a)^{1+s}} \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - \rho_a)^s}{(\rho_b - x)^{\zeta+s}} \int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned} \quad (23)$$

REMARK 6. If $\varepsilon = 0$ in (23), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - \rho_a)^s (\rho_b - x)^{1-\zeta}}{(x - \rho_a)^{1+s}} \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - \rho_a)^s}{(\rho_b - x)^{\zeta+s}} \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned}$$

4. If $\lambda = 1$, $\phi(t) = \frac{1}{t^2}$ in (20), then Fractional Ostrowski type inequality for Godunova-Levin convex (concave):

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon (\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - \rho_a)(\rho_b - x)^{1-\zeta}}{(x - \rho_a)^2} \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - \rho_a)}{(\rho_b - x)^{\zeta+1}} \int_x^{\rho_b} \eta \left[\frac{(t - \nu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned} \tag{24}$$

REMARK 7. If $\varepsilon = 0$ in (24), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - \rho_a)(\rho_b - x)^{1-\zeta}}{(x - \rho_a)^2} \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - \rho_a)}{(\rho_b - x)^{\zeta+1}} \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned}$$

5. If $\lambda = 1$, $\phi(t) = t^{s-1}$ with $s \in [0, 1]$ in (20), then Fractional Ostrowski type inequality for s -convex (concave) in 2nd kind:

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^{\zeta} \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon (\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(x - \rho_a)^{s-1} (\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)^s} \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - x)^{s-\zeta}}{(\rho_b - \rho_a)^s} \int_x^{\rho_b} \eta \left[\frac{(t - \nu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned} \tag{25}$$

REMARK 8. If $\varepsilon = 0$ in (25), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(x - \rho_a)^{s-1} (\rho_b - x)^{1-\zeta}}{(\rho_b - \rho_a)^s} \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \\ & \quad + \frac{(\rho_b - x)^{s-\zeta}}{(\rho_b - \rho_a)^s} \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned}$$

6. If $\lambda = 1$, $\phi(t) = \frac{1}{t}$ in (20), then Fractional Ostrowski type inequality for P -convex (concave):

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon (\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{(x - \rho_a)} \int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt + \frac{1}{(\rho_b - x)^\zeta} \int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned} \tag{26}$$

REMARK 9. If $\varepsilon = 0$ in (26), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{(x - \rho_a)} \int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt + \frac{1}{(\rho_b - x)^\zeta} \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt. \end{aligned}$$

7. If $\lambda = \phi(t) = 1$ in (20), then Fractional Ostrowski type inequality for convex (concave):

$$\begin{aligned} & \eta \left[(1 - \varepsilon) \varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon (\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{\rho_b - \rho_a} \left[\int_{\rho_a}^x \eta \left[\frac{(t - \mu) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt + \int_x^{\rho_b} \eta \left[\frac{(t - v) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned} \tag{27}$$

REMARK 10. If $\varepsilon = 0$ in (27), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{1-\zeta}}{\rho_b - \rho_a} \left[\int_{\rho_a}^x \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt + \int_x^{\rho_b} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned}$$

8. If $\lambda = 1$, $\phi(t) = \frac{1}{2\sqrt{t(1-t)}}$ in (20), then Fractional Ostrowski type inequality MT -convex (concave):

$$\begin{aligned} & \eta \left[(1-\varepsilon)\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) \right. \\ & \quad \left. + J_{\rho_a}^{\zeta-1} (P_2(x, \rho_b) \varphi(\rho_b)) + \frac{\varepsilon(\rho_b - x)^{1-\zeta}}{2(\rho_b - \rho_a)^{1-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{\frac{1}{2}-\zeta}}{2\sqrt{(x-\rho_a)}} \left[\int_{\rho_a}^x \eta \left[\frac{(t-\mu)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt + \int_x^{\rho_b} \eta \left[\frac{(t-v)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt \right]. \end{aligned} \quad (28)$$

REMARK 11. If $\varepsilon = 0$ in (28), we get

$$\begin{aligned} & \eta \left[\varphi(x) - \frac{\Gamma(\zeta)}{\rho_b - \rho_a} (\rho_b - x)^{1-\zeta} J_{\rho_a}^\zeta \varphi(\rho_b) + J_{\rho_a}^{\zeta-1} (P_1(x, \rho_b) \varphi(\rho_b)) \right] \\ & \leqslant (\geqslant) \frac{(\rho_b - x)^{\frac{1}{2}-\zeta}}{2\sqrt{(x-\rho_a)}} \left[\int_{\rho_a}^x \eta \left[\frac{(t-\rho_a)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt + \int_x^{\rho_b} \eta \left[\frac{(t-\rho_b)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt \right]. \end{aligned}$$

3. Applications of midpoint inequalities

If we replace φ by $-\varphi$ and $x = \frac{\rho_a+\rho_b}{2}$ in Theorem 7, we get

THEOREM 8. Let $\varphi : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be differentiable on (ρ_a, ρ_b) , $\varphi' : [\rho_a, \rho_b] \rightarrow \mathbb{R}$ be integrable on $[\rho_a, \rho_b]$ and $\eta : I \subset \mathbb{R} \rightarrow \mathbb{R}$, be a $\phi - \lambda$ -convex (concave), then

$$\begin{aligned} & \eta \left[\frac{\Gamma(\zeta) (\frac{\rho_b-\rho_a}{2})^{1-\zeta}}{\rho_b - \rho_a} J_a^\zeta \varphi(\rho_b) - \varphi \left(\frac{\rho_a + \rho_b}{2} \right) \right. \\ & \quad \left. - J_a^{\zeta-1} \left(P_2 \left(\frac{\rho_a + \rho_b}{2}, b \right) \varphi(\rho_b) \right) - \frac{\varepsilon}{2^{2-\zeta}} J_{\rho_a}^0 \varphi(\rho_a) \right] \\ & \leqslant (\geqslant) \frac{2^{\zeta-\lambda} \phi(\frac{1}{2})}{(\rho_b - \rho_a)^\zeta} \left[\int_{\frac{\rho_a+\rho_b}{2}}^{\rho_a} \eta \left[\frac{(t-\mu)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt + \int_{\rho_b}^{\frac{\rho_a+\rho_b}{2}} \eta \left[\frac{(t-v)\varphi'(t)}{(\rho_b-t)^{1-\zeta}} \right] dt \right]. \end{aligned} \quad (29)$$

$\forall \varepsilon \in [0, 1]$.

REMARK 12. In Theorem 8,

1. If $\varepsilon = 0$, in (29) we get

$$\begin{aligned} & \eta \left[\frac{\Gamma(\zeta) \left(\frac{\rho_b - \rho_a}{2} \right)^{1-\zeta}}{\rho_b - \rho_a} J_a^\zeta \varphi(\rho_b) - \varphi \left(\frac{\rho_a + \rho_b}{2} \right) - J_a^{\zeta-1} \left(P_1 \left(\frac{\rho_a + \rho_b}{2}, b \right) \varphi(\rho_b) \right) \right] \\ & \leqslant (\geqslant) \frac{2^{\zeta-\lambda} \phi \left(\frac{1}{2} \right)}{(\rho_b - \rho_a)^\zeta} \left[\int_{\frac{\rho_a+\rho_b}{2}}^{\rho_a} \eta \left[\frac{(t - \rho_a) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt + \int_{\rho_b}^{\frac{\rho_a+\rho_b}{2}} \eta \left[\frac{(t - \rho_b) \varphi'(t)}{(\rho_b - t)^{1-\zeta}} \right] dt \right]. \end{aligned}$$

2. If $\zeta = 1$ in (29) we get

$$\begin{aligned} & \eta \left[(\varepsilon - 1) \varphi \left(\frac{\rho_a + \rho_b}{2} \right) - \varepsilon \frac{\varphi(a) + \varphi(\rho_b)}{2} + \frac{1}{\rho_b - \rho_a} \int_{\rho_a}^{\rho_b} \varphi(t) dt \right] \\ & \leqslant (\geqslant) \frac{2^{1-\lambda} \phi \left(\frac{1}{2} \right)}{\rho_b - \rho_a} \left[\int_{\rho_a}^{\frac{\rho_a+\rho_b}{2}} \eta [(\mu - t) \varphi'(t)] dt + \int_{\frac{\rho_a+\rho_b}{2}}^{\rho_b} \eta [(\nu - t) \varphi'(t)] dt \right]. \end{aligned}$$

3. If $\varepsilon = 0$, $\zeta = 1$ in (29) we get

$$\begin{aligned} & \eta \left[\frac{1}{\rho_b - \rho_a} \int_{\rho_a}^{\rho_b} \varphi(t) dt - \varphi \left(\frac{\rho_a + \rho_b}{2} \right) \right] \\ & \leqslant (\geqslant) \frac{2^{1-\lambda} \phi \left(\frac{1}{2} \right)}{\rho_b - \rho_a} \left[\int_{\rho_a}^{\frac{\rho_a+\rho_b}{2}} \eta [(\rho_a - t) \varphi'(t)] dt + \int_{\frac{\rho_a+\rho_b}{2}}^{\rho_b} \eta [(\rho_b - t) \varphi'(t)] dt \right]. \end{aligned}$$

REMARK 13. Assume that $\eta : I \subset [0, \infty) \rightarrow \mathbb{R}$ be an ϕ -convex (concave):

1. If $\zeta = 1$, $\varphi(t) = \frac{1}{t}$ in inequality (30) where $t \in [\rho_a, \rho_b] \subset (0, \infty)$, then we have

$$\begin{aligned} & \eta \left[\frac{A(\rho_a, \rho_b) + (\varepsilon - 1)L(\rho_a, \rho_b)}{A(\rho_a, \rho_b)L(\rho_a, \rho_b)} - \varepsilon \frac{A(\rho_a, \rho_b)}{G^2(\rho_a, \rho_b)} \right] \\ & \leqslant (\geqslant) \frac{2^{1-\lambda} \phi \left(\frac{1}{2} \right)}{\rho_b - \rho_a} \left[\int_{\rho_a}^{\frac{\rho_a+\rho_b}{2}} \eta \left[\frac{t - \mu}{t^2} \right] dt + \int_{\frac{\rho_a+\rho_b}{2}}^{\rho_b} \eta \left[\frac{t - \nu}{t^2} \right] dt \right]. \end{aligned}$$

2. If $\zeta = 1$, $\varphi(t) = -\ln t$ in inequality (30), where $t \in [\rho_a, \rho_b] \subset (0, \infty)$, then we have

$$\begin{aligned} & \eta \left[\ln \left(\frac{\exp[\varepsilon A(\ln \rho_a, \ln \rho_b)] A^{(1-\varepsilon)}(\rho_a, \rho_b)}{I(\rho_a, \rho_b)} \right) \right] \\ & \leqslant (\geqslant) \frac{2^{1-\lambda} \phi \left(\frac{1}{2} \right)}{\rho_b - \rho_a} \left[\int_{\rho_a}^{\frac{\rho_a+\rho_b}{2}} \eta \left[\frac{t - \mu}{t} \right] dt + \int_{\frac{\rho_a+\rho_b}{2}}^{\rho_b} \eta \left[\frac{t - \nu}{t} \right] dt \right]. \end{aligned}$$

3. If $\zeta = 1$, $\varphi(t) = t^p$, $p \in \mathbb{R} - \{0, -1\}$ in inequality (30), where $t \in [\rho_a, \rho_b] \subset (0, \infty)$, then we have

$$\begin{aligned} & \eta [L_p^p(\rho_a, \rho_b) - (\varepsilon - 1)A^p(\rho_a, \rho_b) - \varepsilon A(a^p, b^p)] \\ & \leqslant (\geqslant) \frac{2^{1-\lambda} \phi(\frac{1}{2})}{\rho_b - \rho_a} \left[\int_{\rho_a}^{\frac{\rho_a + \rho_b}{2}} \eta \left[\frac{p(\mu-t)}{t^{1-p}} \right] dt + \int_{\frac{\rho_a + \rho_b}{2}}^{\rho_b} \eta \left[\frac{p(v-t)}{t^{1-p}} \right] dt \right]. \end{aligned}$$

REMARK 14. In Theorem 5,

1. Let $g(x) = x$, $\zeta = 1$, $x = \frac{\rho_a + \rho_b}{2}$, $0 < \rho_a < \rho_b$, $q \geqslant 1$ and $\varphi : \mathbb{R} \rightarrow \mathbb{R}^+$, $\varphi(x) = x^n$ in (12). Then

$$|A(\rho_a, \rho_b) - L_n^n(\rho_a, \rho_b)| \leqslant \frac{M(\rho_b - \rho_a)}{(2)^{2-\frac{1}{q}}} \left(\int_0^1 [t^{1+\lambda} \phi(t) + t(1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}}.$$

2. Let $g(x) = x$, $\zeta = 1$, $x = \frac{\rho_a + \rho_b}{2}$, $0 < \rho_a < \rho_b$, $q \geqslant 1$ and $\varphi : (0, 1] \rightarrow \mathbb{R}$, $\varphi(x) = -\ln x$ in (12). Then

$$|\ln I(\rho_a, \rho_b) - \ln A(\rho_a, \rho_b)| \leqslant \frac{M(\rho_b - \rho_a)}{(2)^{2-\frac{1}{q}}} \left(\int_0^1 [t^{1+\lambda} \phi(t) + t(1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}}.$$

REMARK 15. In Theorem 6,

1. Let $g(x) = x$, $\zeta = 1$, $x = \frac{\rho_a + \rho_b}{2}$, $0 < \rho_a < \rho_b$, $q \geqslant 1$ and $\varphi : \mathbb{R} \rightarrow \mathbb{R}^+$, $\varphi(x) = x^n$ in (16). Then

$$|A(\rho_a, \rho_b) - L_n^n(\rho_a, \rho_b)| \leqslant \frac{M(\rho_b - \rho_a)}{2(p+1)^{\frac{1}{p}}} \left(\int_0^1 [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}}.$$

2. Let $g(x) = x$, $\zeta = 1$, $x = \frac{\rho_a + \rho_b}{2}$, $0 < \rho_a < \rho_b$, $q \geqslant 1$ and $\varphi : (0, 1] \rightarrow \mathbb{R}$, $\varphi(x) = -\ln x$ in (16). Then

$$|\ln I(\rho_a, \rho_b) - \ln A(\rho_a, \rho_b)| \leqslant \frac{M(\rho_b - \rho_a)}{2(p+1)^{\frac{1}{p}}} \left(\int_0^1 [t^\lambda \phi(t) + (1-t)^\lambda \phi(1-t)] dt \right)^{\frac{1}{q}}.$$

4. Conclusion

Ostrowski inequality is one of the most celebrated inequalities, we can find its various generalizations and variants in literature. In this paper, the generalization of Fractional Ostrowski inequality via generalized Montgomery identity [14] with $\phi - \lambda$ -convex. This class of functions contains many important classes including class of ϕ -convex [10], h -convex [23], Godunova-Levin s -convex [9], s -convex in the 2^{nd} kind

[3] and hence contains class of convex and MT -convex [2]. It also contains class of P -convex [15] and class of Godunova-Levin functions [17]. We have stated our main result in section 2, which is the generalization of Ostrowski inequality via generalized Montgomery identity by generalized fractional integrals for $\phi - \lambda$ -convex. Further, we used different techniques including Hölder's inequality [25] and power mean inequality [24] for generalization of Ostrowski inequality. In second last section we have given some applications in terms of special means including arithmetic, geometric, harmonic, logarithmic, identric and p -logarithmic means by using the midpoint inequalities.

REFERENCES

- [1] M. ALOMARI, M. DARUS, S. S. DRAGOMIR AND P. CERONE, *Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense*, Appl. Math. Lett., **23**, (2010), 1071–1076.
- [2] E. F. BECKENBACH, *Convex*, Bull. Amer. Math. Soc., **54**, (1948), 439–460.
- [3] W. W. BRECKNER, *Stetigkeitsaussagen Fur Eine Klasse Verallgemeinerter Konvexer Funktionen in Topologischen Linearen Raumern* (German), Publ. Inst. Math., **23** 37, (1978), 13–20.
- [4] E. SET, *New inequalities of Ostrowski type for mappings whose derivatives are s -convex in the second sense via fractional integrals*, Comput. Math. Appl., **63** (2012), 1147–1154.
- [5] M. J. V. CORTEZ, J. E. HERNÁNDEZ, *Ostrowski and Jensen-type inequalities via (s,m) -convex in the second sense*, Bol. Soc. Mat. Mex., **26**, (2020), 287–302.
- [6] S. S. DRAGOMIR, *A Companion of Ostrowski's Inequality for Functions of Bounded Variation and Applications*, Int. J. Nonlinear Anal. Appl., **5** (2014), 89–97.
- [7] S. S. DRAGOMIR, *The Generalization of Ostrowski Inequality via Montgomery identity*, Acta Math. Univ. Comenianae, LXXXIV, **1**, (2015), 63–78.
- [8] S. S. DRAGOMIR, *On the Ostrowski's Integral Inequality for Mappings with Bounded Variation and Applications*, Math. Inequal. Appl., **4** (1), (2001), 59–66.
- [9] S. S. DRAGOMIR, *Integral inequalities of Jensen type for λ -convex*, In Proceedings of RGMIA, Res. Rep. Coll., **1** (17), (2014).
- [10] S. S. DRAGOMIR, *Inequalities of Jensen Type for ϕ -convex*, Fasciculi Mathematici, **5**, (2015), 35–52.
- [11] S. S. DRAGOMIR, *Refinements of the Generalised Trapezoid and Ostrowski Inequalities for Functions of Bounded Variation*, Arch. Math., **91** (5), (2008), 450–460.
- [12] S. S. DRAGOMIR AND N. S. BARNETT, *An Ostrowski Type Inequality for Mappings whose Second Derivatives are Bounded and Applications*, J. Indian Math. Soc. (N.S.), **66** (4) (1999), 237–245.
- [13] S. S. DRAGOMIR, P. CERONE, N. S. BARNETT AND J. ROUMELIOTIS, *An Inequality of the Ostrowski Type for Double Integrals and Applications for Cubature Formulae*, Tamsui Oxf. J. Math. Sci., **16** (1), (2000), 1–16.
- [14] S. S. DRAGOMIR, P. CERONE AND J. ROUMELIOTIS, *A new Generalization of Ostrowski Integral Inequality for Mappings whose Derivatives are Bounded and Applications in Numerical Integration and for Special Means*, Appl. Math. Lett., **13** (2000), 19–25.
- [15] S. S. DRAGOMIR, J. PEĆARIĆ AND L. PERSSON, *Some inequalities of Hadamard type*, Soochow J. Math., **21** (3), (1995), 335–341.
- [16] A. EKİNCİ, *Klasik Eşitsizlikler Yoluyla Konveks Fonksiyonlar için Integral Eşitsizlikler*, Ph. D. Thesis, Thesis ID: 361162 in tez2.yok.gov.tr Atatürk University, 2014.
- [17] E. K. GODUNOVA, V. I. LEVIN, *Inequalities for functions of a broad class that contains convex, monotone and some other forms of functions*, Numerical Mathematics and Mathematical Physics, (Russian), **166**, (1985), 138–142.
- [18] M. MATLOKA, *On Ostrowski type inequalities via fractional integrals of The with respect to another function*, Journal pf Nonlinear Science and Applications, **13**, (2020), 100–106
- [19] M. A. NOOR AND M. U. AWAN, *Some integral inequalities for two kinds of convexities via fractional integrals*, TJMM, **5**, (2013), 129–136.
- [20] A. M. OSTROWSKI, *Über die absolutabweichung einer differentierbaren funktion von ihrem integralmittelwert*, Comment. Math. Helv., **10**, (1938), 226–227.

- [21] S. G. SAMKO, A. A. KILBAS, O. I. MARICHEV, *Fractional Integrals and Derivatives, Theory and Applications* Gordon and Breach New York, **1**, (1993).
- [22] E. SET, *New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals*, Comput. Math. Appl., **63**, (2012), 1147–1154.
- [23] S. VAROŠANEC, *On h-convexity*, J. Math. Anal. Appl., **326** (1), (2007), 303–311.
- [24] Z. G. XIAO, AND A. H. ZHANG, *Mixed power mean inequalities*, Research Communication on Inequalities, **8** (1), (2002), 15–17.
- [25] X. YANG, *A note on Hölder inequality*, Appl. Math. Comput., **134**, (2003), 319–322.

(Received August 26, 2021)

Ali Hassan

*Department of Mathematics
Shah Abdul Latif University
Khairpur-66020, Pakistan
e-mail: alihassan.iiui.math@gmail.com*

Asif Raza Khan

*Department of Mathematics
University of Karachi
University Road, Karachi-75270, Pakistan
e-mail: asifrk@uok.edu.pk*