

OPERATORS OF FRACTIONAL CALCULUS AND ASSOCIATED INTEGRAL TRANSFORMS OF THE (p, q)-EXTENDED HURWITZ-LERCH ZETA FUNCTION

PURNIMA CHOPRA*, SHILPA KUMAWAT AND HEMLATA SAXENA

(Communicated by H. M. Srivastava)

Abstract. In this paper, our aim is to establish certain fractional integral and derivative formulas of the generalized (p, q)-extended Hurwitz-Lerch zeta function by using generalized Marichev-Saigo-Maeda fractional operators which involve, in their kernel, Appell's two-variable hypergeometric function $F_3(\cdot)$. These results are expressed in terms of the Hadamard product (or the convolution) of two analytic functions in terms of (p, q)-extended Hurwitz-Lerch zeta function and Fox-Wright hypergeometric function ${}_r\Psi_s(\cdot)$. We then obtain their composition formulas by using fractional integral and derivative formulas and certain Integral transforms associated with Beta, Laplace and Whittaker transforms involving generalized (p, q)-extended Hurwitz-Lerch Zeta function.

1. Introduction

Operators of fractional calculus (such as the Riemann-Liouville, Weyl, Liouville-Caputo, and other operators of fractional integration and fractional derivative) have been developed and investigated widely and extensively, because mainly of their importance and potential for applications (see for details, [15], [24] and [38]). We first recall a general pair of fractional integral operators popularly known as Marichev-Saigo-Maeda which involve, in their kernel, Appell's two-variable hypergeometric function $F_3(\cdot)$ of the third kind, which is defined by (see, for details, [17, 26, 27]):

DEFINITION 1. Let $\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta \in \mathbb{C}$ and $x > 0$, then for $\text{Re}(\eta) > 0$,

$$\begin{aligned} \left(I_{0,x}^{\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta} f \right) (x) &= \frac{x^{-\sigma_1}}{\Gamma(\eta)} \int_0^x (x-t)^{\eta-1} t^{-\sigma'_1} \\ &\quad \times F_3 \left(\sigma_1, \sigma'_1, \nu_1, \nu'_1; \eta; 1 - \frac{t}{x}, 1 - \frac{x}{t} \right) f(t) dt. \end{aligned} \quad (1)$$

Mathematics subject classification (2020): 26A33, 33B15, 33C05, 33C99, 44A20.

Keywords and phrases: Hurwitz-Lerch zeta function, (p, q)-extended Hurwitz-Lerch zeta function, fractional calculus operators.

* Corresponding author.

and

$$\begin{aligned} \left(I_{x,\infty}^{\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta} f \right) (x) &= \frac{x^{-\sigma'_1}}{\Gamma(\eta)} \int_x^\infty (t-x)^{\eta-1} t^{-\sigma_1} \\ &\quad \times F_3 \left(\sigma_1, \sigma'_1, \nu_1, \nu'_1; \eta; 1 - \frac{x}{t}, 1 - \frac{t}{x} \right) f(t) dt. \end{aligned} \quad (2)$$

Here $F_3(\cdot)$ denotes the Appell's hypergeometric function of two variables [37].

DEFINITION 2. Let $\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta \in \mathbb{C}$ and $x > 0$, then for $\text{Re}(\eta) > 0$,

$$\begin{aligned} \left(D_{0,x}^{\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta} f \right) (x) &= \left(I_{0+}^{-\sigma'_1, -\sigma_1, -\nu'_1, -\nu_1, -\eta} f \right) (x) \\ &= \left(\frac{d}{dx} \right)^n \left(I_{0+}^{-\sigma'_1, -\sigma_1, -\nu'_1+n, -\nu_1, -\eta+n} f \right) (x) \quad (n = [\text{Re}(\eta)] + 1) \\ &= \frac{1}{\Gamma(n-\eta)} \left(\frac{d}{dx} \right)^n x^{\sigma'_1} \int_0^x (x-t)^{n-\eta-1} t^{\sigma} \\ &\quad \times F_3 \left(-\sigma'_1, -\sigma_1, n - \nu'_1, -\nu_1; n - \eta; 1 - \frac{t}{x}, 1 - \frac{x}{t} \right) f(t) dt. \end{aligned} \quad (3)$$

and

$$\begin{aligned} \left(D_{x,\infty}^{\sigma_1, \sigma'_1, \nu_1, \nu'_1, \eta} f \right) (x) &= \left(I_-^{-\sigma'_1, -\sigma_1, -\nu'_1, -\nu_1, -\eta} f \right) (x) \\ &= \left(-\frac{d}{dx} \right)^n \left(I_-^{-\sigma'_1, -\sigma_1, -\nu'_1, -\nu_1, -\eta+n} f \right) (x) \quad (n = [\text{Re}(\eta)] + 1) \\ &= \frac{1}{\Gamma(n-\eta)} \left(-\frac{d}{dx} \right)^n x^{\sigma'_1} \int_x^\infty (t-x)^{n-\eta-1} t^{\sigma'} \\ &\quad \times F_3 \left(-\sigma'_1, -\sigma_1, \nu'_1, n - \nu_1; n - \eta; 1 - \frac{x}{t}, 1 - \frac{t}{x} \right) f(t) dt. \end{aligned} \quad (4)$$

These operators includes Saigo hypergeometric fractional calculus operators, Riemann-Liouville and Erdélyi-Kober fractional calculus operators as special cases for various choice of parameters (see for details, [15], [24] and [38]). In recent papers, various authors established several interesting generalized fractional formulas involving (p, q) -extended Bessel function and modified Bessel function, (p, q) -extended Sturve function and modified Sturve function and (p, q) -extended Mathieu series (see, for details, [9, 13]) and other recent developments on several general families of integral transformations and fractional integrations and differentiations of certain higher transcendental functions [28, 31, 33, 34].

The more generalized form so-called (p, q) -extended Hurwitz-Lerch zeta function has been considered very recently by Luo *et al.* [12] in the following form

$$\Phi_{\lambda, \mu; \nu}(z, s, a; p, q) := \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} \frac{B(\mu+n, \nu-\mu; p, q)}{B(\mu, \nu-\mu)} \frac{z^n}{(n+a)^s} \quad (5)$$

$$(\min\{\Re(p), \Re(q)\} \geq 0; p, q, \lambda, \mu, s \in \mathbb{C}; \nu, a \in \mathbb{C} \setminus \mathbb{Z}_0^-; |z| < 1).$$

where $B(x, y; p, q)$ is the (p, q) -extended Beta function introduced by Choi *et al.* [3]

$$B(x, y; p, q) = B_{p,q}(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} e^{-\frac{p}{t} - \frac{q}{1-t}} dt, \quad (6)$$

when $\min\{Re(x), Re(y)\} > 0; \min\{Re(p), Re(q)\} \geq 0$. They also introduced (p, q) -extended of hypergeometric function as

$$F_{p,q}(a, b; c; z) = \sum_{n \geq 0} (a)_n \frac{B(b+n, c-b; p, q)}{B(b, c-b)} \frac{z^n}{n!}, \quad (7)$$

$$(p, q \geq 0; |z| < 1; \Re(c) > \Re(b) > 0).$$

Related properties, various integral representations, differentiation formulae, Mellin transform, recurrence relations, summations are also given in [3]. For $p = q$, reduces immediately to the following p -Gauss hypergeometric function $F_p(a, b; c; z)$ studied by Chaudhry *et al.* [2]:

$$F_p(a, b; c; z) := \sum_{n=0}^{\infty} (a)_n \frac{B(b+n, c-b; p)}{B(b, c-b)} \frac{z^n}{n!}$$

$$(p \geq 0; |z| < 1; \Re(c) > \Re(b) > 0).$$

This (p, q) -extended Hurwitz-Lerch zeta function includes another forms of extended Hurwitz-Lerch zeta function as special cases (see, for details, [9, 12, 13]). A detailed study and survey on some recent developments on higher transcendental functions, incomplete Hurwitz-Lerch zeta function, general families of double and multiple Hurwitz-Lerch zeta function and other related functions of analytic number theory with applications can be seen in recent papers [4, 5, 6, 7, 8, 19, 21, 22, 23, 29, 30, 32, 35, 36].

In our present investigation, we require the definition of the Hadamard product (or the convolution) of two analytic functions [9]. If the R_f and R_g be the radii of convergence of the two power series

$$f(z) := \sum_{n=0}^{\infty} a_n z^n \quad (|z| < R_f) \quad \text{and} \quad g(z) := \sum_{n=0}^{\infty} b_n z^n \quad (|z| < R_g),$$

respectively. Then the Hadamard product is the new emerged series defined by

$$(f * g)(z) := \sum_{n=0}^{\infty} a_n b_n z^n = (g * f)(z) \quad (|z| < R) \quad (8)$$

where

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n b_n}{a_{n+1} b_{n+1}} \right| = \left(\lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| \right) \cdot \left(\lim_{n \rightarrow \infty} \left| \frac{b_n}{b_{n+1}} \right| \right) = R_f \cdot R_g,$$

so that, in general, we have $R \geq R_f \cdot R_g$.

In this present note, we aim to develop the compositions of the generalized fractional integral and differential operators (1), (2), (3) and (4) for the (p, q) -extended Hurwitz-Lerch zeta function (5) by using the Hadamard product (8) in terms of (p, q) -extended Hurwitz-Lerch zeta function and Wright hypergeometric function.

2. Fractional formulas of the (p, q) -extended Hurwitz-Lerch zeta function

The Wright hypergeometric function ${}_r\Psi_s(z)$ ($r, s \in \mathbb{N}_0$) having numerator and denominator parameters r and s , respectively, defined for $\alpha_1, \dots, \alpha_r \in \mathbb{C}$ and $\beta_1, \dots, \beta_s \in \mathbb{C} \setminus \mathbb{Z}_0^-$ by (see, for example, [14, 16, 24, 37]):

$${}_r\Psi_s \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_r, A_r) \\ (\beta_1, B_1), \dots, (\beta_s, B_s) \end{matrix}; z \right] = \sum_{n=0}^{\infty} \frac{\Gamma(\alpha_1 + A_1 n) \cdots \Gamma(\alpha_r + A_r n)}{\Gamma(\beta_1 + B_1 n) \cdots \Gamma(\beta_s + B_s n)} \frac{z^n}{n!} \quad (9)$$

$$\left(A_j \in \mathbb{R}^+ (j = 1, \dots, r); B_j \in \mathbb{R}^+ (j = 1, \dots, s); 1 + \sum_{j=1}^s B_j - \sum_{j=1}^r A_j \geq 0 \right),$$

with

$$|z| < \nabla := \left(\prod_{j=1}^r A_j^{-A_j} \right) \cdot \left(\prod_{j=1}^s B_j^{B_j} \right).$$

Also, if we take $A_j = B_k = 1$ ($j = 1, \dots, r; k = 1, \dots, s$) in (9), reduces to the generalized hypergeometric function ${}_rF_s$ ($r, s \in \mathbb{N}_0$) (see, e.g., [37]):

$${}_rF_s \left[\begin{matrix} \alpha_1, \dots, \alpha_r \\ \beta_1, \dots, \beta_s \end{matrix}; z \right] = \frac{\Gamma(\beta_1) \cdots \Gamma(\beta_s)}{\Gamma(\alpha_1) \cdots \Gamma(\alpha_r)} {}_r\Psi_s \left[\begin{matrix} (\alpha_1, 1), \dots, (\alpha_r, 1) \\ (\beta_1, 1), \dots, (\beta_s, 1) \end{matrix}; z \right]. \quad (10)$$

The following image formulas or power function are useful in our investigation [1].

LEMMA 1. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho \in \mathbb{C}$ and $x > 0$. Then the following relation exists

(a) If $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max \{0, \operatorname{Re}(\sigma_1 + \sigma'_1 + v_1 - \eta), \operatorname{Re}(\sigma'_1 - v'_1)\}$, then

$$\begin{aligned} & \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta, t^{\rho-1}} \right) (x) \\ &= \frac{\Gamma(\rho) \Gamma(\rho + \eta - \sigma_1 - \sigma'_1 - v_1) \Gamma(\rho + v'_1 - \sigma'_1)}{\Gamma(\rho + v'_1) \Gamma(\rho + \eta - \sigma_1 - \sigma'_1) \Gamma(\rho + \eta - \sigma'_1 - v_1)} x^{\rho + \eta - \sigma_1 - \sigma'_1 - 1} \end{aligned} \quad (11)$$

(b) If $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min \{\operatorname{Re}(-v_1), \operatorname{Re}(\sigma_1 + \sigma'_1 - \eta), \operatorname{Re}(\sigma_1 + v'_1 - \eta)\}$, then

$$\begin{aligned} & \left(I_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta, t^{\rho-1}} \right) (x) \\ &= \frac{\Gamma(1 - \rho - v_1) \Gamma(1 - \rho - \eta + \sigma_1 + \sigma'_1) \Gamma(1 - \rho - \eta + \sigma_1 + v'_1)}{\Gamma(1 - \rho) \Gamma(1 - \rho - \eta + \sigma_1 + \sigma'_1 + v'_1) \Gamma(1 - \rho + \sigma_1 - v_1)} \\ & \quad \times x^{\rho + \eta - \sigma_1 - \sigma'_1 - 1}. \end{aligned} \quad (12)$$

LEMMA 2. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho \in \mathbb{C}$ and $x > 0$. Then the following relation exists

(a) If $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\eta - \sigma_1 - \sigma'_1 + v'_1), \operatorname{Re}(v_1 - \sigma_1)\}$, then

$$\begin{aligned} & \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} t^{\rho-1} \right) (x) \\ &= \frac{\Gamma(\rho) \Gamma(\rho - \eta + \sigma_1 + \sigma'_1 + v'_1) \Gamma(\rho - v_1 + \sigma_1)}{\Gamma(\rho - v_1) \Gamma(\rho - \eta + \sigma_1 + \sigma'_1) \Gamma(\rho - \eta + \sigma_1 + v'_1)} x^{\rho - \eta + \sigma_1 + \sigma'_1 - 1} \end{aligned} \quad (13)$$

(b) If $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(v'_1), \operatorname{Re}(\eta - \sigma_1 - \sigma'_1), \operatorname{Re}(\eta - \sigma'_1 - v_1)\}$, then

$$\begin{aligned} & \left(D_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} t^{\rho-1} \right) (x) \\ &= \frac{\Gamma(1 - \rho - v'_1) \Gamma(1 - \rho + \eta - \sigma_1 - \sigma'_1) \Gamma(1 - \rho + \eta - \sigma'_1 - v_1)}{\Gamma(1 - \rho) \Gamma(1 - \rho + \eta - \sigma_1 - \sigma'_1 - v) \Gamma(1 - \rho - \sigma'_1 - v'_1)} \\ & \quad \times x^{\rho - \eta + \sigma_1 + \sigma'_1 - 1}. \end{aligned} \quad (14)$$

We begin the main results exposition with presenting the composition formulas of generalized fractional operators (1), (2), (3) and (4) involving the (p, q) -extended Hurwitz-Lerch zeta function by using the Hadamard product (8) in terms of (p, q) -extended Hurwitz-Lerch zeta function (5) and Fox-Wright function (9).

THEOREM 1. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\sigma_1 + \sigma'_1 + v_1 - \eta), \operatorname{Re}(\sigma'_1 - v'_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following formula for fractional integration holds true:

$$\begin{aligned} & \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}(t^\gamma, s, a; p, q) \right\} \right) (x) \\ &= x^{\rho + \eta - \sigma_1 - \sigma'_1 - 1} \Phi_{\lambda, \mu; v}(x^\gamma, s, a; p, q) \\ & \quad * {}_4\Psi_3 \left[\begin{matrix} (1, 1), (\rho, \gamma), (\rho + \eta - \sigma_1 - \sigma'_1 - v_1, \gamma), (\rho + v'_1 - \sigma'_1, \gamma) \\ (\rho + v'_1, \gamma), (\rho + \eta - \sigma_1 - \sigma'_1, \gamma), (\rho + \eta - \sigma'_1 - v_1, \gamma) \end{matrix}; x^\gamma \right]. \end{aligned}$$

Proof. Applying the definitions (5), (1) and then we change the order of integration and using the relation (11), we find for $x > 0$

$$\begin{aligned} & \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}(t^\gamma, s, a; p, q) \right\} \right) (x) \\ &= \sum_{k=0}^{\infty} \frac{(\lambda)_k \mathbf{B}(\mu + k, v - \mu; p, q)}{(k + a)^s \mathbf{B}(\mu, v - \mu) k!} \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho + \gamma k - 1} \right\} \right) (x) \\ &= x^{\rho + \eta - \sigma_1 - \sigma'_1 - 1} \sum_{k=0}^{\infty} \frac{(\lambda)_k \mathbf{B}(\mu + k, v - \mu; p, q)}{(k + a)^s \mathbf{B}(\mu, v - \mu) k!} \\ & \quad \times \frac{\Gamma(\rho + \gamma k) \Gamma(\rho + \eta - \sigma_1 - \sigma'_1 - v_1 + \gamma k) \Gamma(\rho + v'_1 - \sigma'_1 + \gamma k)}{\Gamma(\rho + v'_1 + \gamma k) \Gamma(\rho + \eta - \sigma_1 - \sigma'_1 + \gamma k) \Gamma(\rho + \eta - \sigma'_1 - v_1 + \gamma k)} x^{\gamma k}. \end{aligned} \quad (15)$$

Finally by using the Hadamard product (8) in (15), which in view of (5) and (9), yields the desired formula. \square

THEOREM 2. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(-v_1), \operatorname{Re}(\sigma_1 + \sigma'_1 - \eta), \operatorname{Re}(\sigma_1 + v'_1 - \eta)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following formula for fractional integration holds true:

$$\begin{aligned} & \left(I_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\frac{1}{t^\gamma}, s, a; p, q \right) \right\} \right) (x) \\ &= x^{\rho+\eta-\sigma_1-\sigma'_1-1} \Phi_{\lambda, \mu; v} \left(\frac{1}{x^\gamma}, s, a; p, q \right) \\ & * {}_4\Psi_3 \left[\begin{matrix} (1, 1), (1-\rho-v_1, \gamma), (1-\rho-\eta+\sigma_1+\sigma'_1, \gamma), (1-\rho-\eta+\sigma_1+v'_1, \gamma); \\ (1-\rho, \gamma), (1-\rho-\eta+\sigma_1+\sigma'_1+v'_1, \gamma), (1-\rho+\sigma_1-v_1, \gamma); \end{matrix} \frac{1}{x^\gamma} \right]. \end{aligned}$$

THEOREM 3. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\eta - \sigma_1 - \sigma'_1 - v'_1), \operatorname{Re}(v_1 - \sigma_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following formula for fractional differentiation holds true:

$$\begin{aligned} & \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} (t^\gamma, s, a; p, q) \right\} \right) (x) \\ &= x^{\rho-\eta+\sigma_1+\sigma'_1-1} \Phi_{\lambda, \mu; v} (x^\gamma, s, a; p, q) \\ & * {}_4\Psi_3 \left[\begin{matrix} (1, 1), (\rho, \gamma), (\rho-\eta+\sigma_1+\sigma'_1+v'_1, \gamma), (\rho-v_1+\sigma_1, \gamma); \\ (\rho-v_1, \gamma), (\rho-\eta+\sigma_1+\sigma'_1, \gamma), (\rho-\eta+\sigma_1+v'_1, \gamma); \end{matrix} x^\gamma \right]. \end{aligned}$$

Proof. Applying the definitions (5), (3) and then we change the order of integration and using the relation (13), we find for $x > 0$

$$\begin{aligned} & \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} (x^\gamma, s, a; p, q) \right\} \right) (x) \\ &= \sum_{k=0}^{\infty} \frac{(\lambda)_k \mathbf{B}(\mu+k, v-\mu; p, q)}{(k+a)^s \mathbf{B}(\mu, v-\mu) k!} \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho+\gamma k-1} \right\} \right) (x) \\ &= x^{\rho-\eta+\sigma_1+\sigma'_1-1} \sum_{k=0}^{\infty} \frac{(\lambda)_k \mathbf{B}(\mu+k, v-\mu; p, q)}{(k+a)^s \mathbf{B}(\mu, v-\mu) k!} \\ & \quad \times \frac{\Gamma(\rho+\gamma k) \Gamma(\rho-\eta+\sigma_1+\sigma'_1+v'_1+\gamma k) \Gamma(\rho-v_1+\sigma_1+\gamma k)}{\Gamma(\rho-v_1+\gamma k) \Gamma(\rho-\eta+\sigma_1+\sigma'_1+\gamma k) \Gamma(\rho-\eta+\sigma_1+v'_1+\gamma k)} x^{\gamma k}. \end{aligned} \tag{16}$$

Finally by using the Hadamard product (8) in (16), which in view of (5) and (9), yields the desired formula (3). \square

THEOREM 4. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(v'_1), \operatorname{Re}(\eta - \sigma_1 - \sigma'_1), \operatorname{Re}(\eta - \sigma'_1 - v_1)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following fractional differentiation formula holds true:

$$\begin{aligned} & \left(D_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\frac{1}{t^\gamma}, s, a; p, q \right) \right\} \right) (x) \\ &= x^{\rho-\eta+\sigma_1+\sigma'_1-1} \Phi_{\lambda, \mu; v} \left(\frac{1}{x^\gamma}, s, a; p, q \right) \\ & * {}_4\Psi_3 \left[\begin{matrix} (1, 1), (1-\rho-v'_1, \gamma), (1-\rho+\eta-\sigma_1-\sigma'_1, \gamma), (1-\rho+\eta-\sigma'_1-v_1, \gamma) \\ (1-\rho, \gamma), (1-\rho+\eta-\sigma_1-\sigma'_1-v_1, \gamma), (1-\rho-\sigma'_1-v'_1, \gamma) \end{matrix}; \frac{1}{x^\gamma} \right]. \end{aligned}$$

The above Theorem 1 to Theorem 4 includes various special cases for which particular choice of parameters reduces to Saigo hypergeometric fractional calculus operators, Riemann-Liouville and Erdélyi-Kober fractional calculus operators. These special cases can be easily evaluated and are left for interesting readers.

3. Certain integral transforms

With the help of the results established in the previous section, in this section, we shall present certain very interesting results in the form of several Theorems associated with Beta, Laplace and Whittaker transforms. For this, first we would like to define these transforms.

DEFINITION 3. The Euler-Beta transform [25] of the function $f(z)$ is defined, as usual, by

$$\mathcal{B}\{f(z); a, b\} = \int_0^1 z^{a-1} (1-z)^{b-1} f(z) dz. \quad (17)$$

DEFINITION 4. The Laplace transform (see, e.g., [25]) of the function $f(z)$ is defined, as usual, by

$$L\{f(z); t\} = \int_0^\infty e^{-tz} f(z) dz. \quad (\operatorname{Re}(t) > 0) \quad (18)$$

The following integral involving Whittaker function (see Mathai *et al.* [16, p. 79]):

$$\int_0^\infty t^{\rho-1} e^{-\frac{1}{2}at} W_{\kappa, v}(at) dt = a^{-\rho} \frac{\Gamma(\frac{1}{2} \pm v + \rho)}{\Gamma(1 - \kappa + \rho)} \quad (\operatorname{Re}(a) > 0, \operatorname{Re}(\rho \pm v) > -\frac{1}{2}), \quad (19)$$

is useful in this section, where $W_{\kappa, v}$ is the Whittaker function [18, p. 334].

The following interesting results in the form of Theorems will be established in this section. As these results are direct consequences of the definitions (17), (18), (19) and Theorems 1 to 4, therefore they are given here without proof.

THEOREM 5. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\sigma_1 + \sigma'_1 + v_1 - \eta), \operatorname{Re}(\sigma'_1 - v'_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Beta-transform formula holds true:

$$\begin{aligned} B \left\{ \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((tz)^\gamma, s, a; p, q) \right\} \right) (x) : l, m \right\} \\ = x^{\rho+\eta-\sigma_1-\sigma'_1-1} \Gamma(m) \Phi_{\lambda, \mu; v}(x^\gamma, s, a; p, q) \\ * {}_5\Psi_4 \left[\begin{matrix} (1, 1), (l, \gamma), (\rho, \gamma), (\rho + \eta - \sigma_1 - \sigma'_1 - v_1, \gamma), (\rho + v'_1 - \sigma'_1, \gamma); \\ (l + m, \gamma), (\rho + v'_1, \gamma), (\rho + \eta - \sigma_1 - \sigma'_1, \gamma), (\rho + \eta - \sigma'_1 - v_1 + \gamma, \gamma); \end{matrix} x^\gamma \right]. \end{aligned}$$

THEOREM 6. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(-v_1), \operatorname{Re}(\sigma_1 + \sigma'_1 - \eta), \operatorname{Re}(\sigma_1 + v'_1 - \eta)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Beta-transform formula holds true:

$$\begin{aligned} B \left\{ \left(I_{x, \infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{z}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) : l, m \right\} \\ = x^{\rho+\eta-\sigma_1-\sigma'_1-1} \Gamma(m) \Phi_{\lambda, \mu; v} \left(\frac{1}{x^\gamma}, s, a; p, q \right) \\ * {}_5\Psi_4 \left[\begin{matrix} (1, 1), (l, \gamma), (1 - \rho - v_1, \gamma), \\ (l + m, \gamma), (1 - \rho, \gamma), \\ (1 - \rho - \eta + \sigma_1 + \sigma'_1, \gamma), (1 - \rho - \eta + \sigma_1 + v'_1, \gamma); \\ (1 - \rho - \eta + \sigma_1 + \sigma'_1 + v'_1, \gamma), (1 - \rho + \sigma_1 - v_1, \gamma); \end{matrix} \frac{1}{x^\gamma} \right]. \end{aligned}$$

THEOREM 7. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\eta - \sigma_1 - \sigma'_1 - v'_1), \operatorname{Re}(v_1 - \sigma_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Beta-transform formula holds true:

$$\begin{aligned} B \left\{ \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((tz)^\gamma, s, a; p, q) \right\} \right) (x) : l, m \right\} \\ = x^{\rho-\eta+\sigma_1+\sigma'_1-1} \Gamma(m) \Phi_{\lambda, \mu; v}(x^\gamma, s, a; p, q) \\ * {}_5\Psi_4 \left[\begin{matrix} (1, 1), (l, \gamma), (\rho, \gamma), (\rho - \eta + \sigma_1 + \sigma'_1 + v'_1, \gamma), (\rho - v_1 + \sigma_1, \gamma); \\ (l + m, \gamma), (\rho - v_1, \gamma), (\rho - \eta + \sigma_1 + \sigma'_1, \gamma), (\rho - \eta + \sigma_1 + v'_1 + \gamma, \gamma); \end{matrix} x^\gamma \right]. \end{aligned}$$

THEOREM 8. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(v'_1), \operatorname{Re}(\eta - \sigma_1 - \sigma'_1), \operatorname{Re}(\eta - \sigma'_1 - v_1)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Beta-transform

formula holds true:

$$\begin{aligned}
 & B \left\{ \left(D_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{z}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) : l, m \right\} \\
 &= x^{\rho-\eta+\sigma_1+\sigma'_1-1} \Gamma(m) \Phi_{\lambda, \mu; v} \left(\frac{1}{x^\gamma}, s, a; p, q \right) \\
 & \quad * {}_5\Psi_4 \left[\begin{matrix} (1, 1), (l, \gamma), (1-\rho-v'_1, \gamma), \\ (1-\rho, \gamma), (1-\rho+\eta-\sigma_1-\sigma'_1-v_1, \gamma), \\ (1-\rho+\eta-\sigma_1-\sigma'_1, \gamma), (1-\rho+\eta-\sigma'_1-v_1, \gamma); \frac{1}{x^\gamma} \end{matrix} \right].
 \end{aligned}$$

THEOREM 9. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\sigma_1 + \sigma'_1 + v_1 - \eta), \operatorname{Re}(\sigma'_1 - v'_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Laplace-transform formula holds true:

$$\begin{aligned}
 & L \left\{ z^{l-1} \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((tz)^\gamma, s, a; p, q) \right\} \right) (x) \right\} \\
 &= \frac{x^{\rho+\eta-\sigma_1-\sigma'_1-1}}{s^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{x}{s} \right)^\gamma, s, a; p, q \right) \\
 & \quad * {}_5\Psi_3 \left[\begin{matrix} (1, 1), (l, \gamma), (\rho, \gamma), (\rho+\eta-\sigma_1-\sigma'_1-v_1, \gamma), (\rho+v'_1-\sigma'_1, \gamma); \left(\frac{x}{s} \right)^\gamma \\ (\rho+v'_1, \gamma), (\rho+\eta-\sigma_1-\sigma'_1, \gamma), (\rho+\eta-\sigma'_1-v_1, \gamma) \end{matrix} \right].
 \end{aligned}$$

THEOREM 10. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(-v_1), \operatorname{Re}(\sigma_1 + \sigma'_1 - \eta), \operatorname{Re}(\sigma_1 + v'_1 - \eta)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Laplace-transform formula holds true:

$$\begin{aligned}
 & L \left\{ z^{l-1} \left(I_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{z}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) \right\} \\
 &= \frac{x^{\rho+\eta-\sigma_1-\sigma'_1-1}}{s^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{1}{xs} \right)^\gamma, s, a; p, q \right) \\
 & \quad * {}_5\Psi_3 \left[\begin{matrix} (1, 1), (l, \gamma), (1-\rho-v_1, \gamma), (1-\rho-\eta+\sigma_1+\sigma'_1, \gamma), \\ (1-\rho, \gamma), (1-\rho-\eta+\sigma_1+\sigma'_1+v'_1, \gamma), \\ (1-\rho-\eta+\sigma_1+v'_1, \gamma); \left(\frac{1}{xs} \right)^\gamma \end{matrix} \right].
 \end{aligned}$$

THEOREM 11. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\eta - \sigma_1 - \sigma'_1 - v'_1), \operatorname{Re}(v_1 - \sigma_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following formula Laplace-transform

holds true:

$$\begin{aligned} & L \left\{ z^{l-1} \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((tz)^\gamma, s, a; p, q) \right\} \right) (x) : \right\} \\ &= \frac{x^{\rho-\eta+\sigma_1+\sigma'_1-1}}{s^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{x}{s} \right)^\gamma, s, a; p, q \right) \\ & * {}_5\Psi_3 \left[\begin{matrix} (1, 1), (l, \gamma), (\rho, \gamma), (\rho - \eta + \sigma_1 + \sigma'_1 + v'_1, \gamma), (\rho - v_1 + \sigma_1, \gamma); \\ (\rho - v_1, \gamma), (\rho - \eta + \sigma_1 + \sigma'_1, \gamma), (\rho - \eta + \sigma_1 + v'_1, \gamma); \end{matrix} \left(\frac{x}{s} \right)^\gamma \right]. \end{aligned}$$

THEOREM 12. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\text{Re}(\eta) > 0$ and $\text{Re}(\rho) < 1 + \min\{\text{Re}(v'_1), \text{Re}(\eta - \sigma_1 - \sigma'_1), \text{Re}(\eta - \sigma'_1 - v_1)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following Laplace-transform formula holds true:

$$\begin{aligned} & L \left\{ z^{l-1} \left(D_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{z}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) \right\} \\ &= \frac{x^{\rho-\eta+\sigma_1+\sigma'_1-1}}{s^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{1}{xs} \right)^\gamma, s, a; p, q \right) \\ & * {}_5\Psi_3 \left[\begin{matrix} (1, 1), (l, \gamma), (1 - \rho - v'_1, \gamma), \\ (1 - \rho, \gamma), (1 - \rho + \eta - \sigma_1 - \sigma'_1 - v_1, \gamma), \\ (1 - \rho + \eta - \sigma_1 - \sigma'_1, \gamma), (1 - \rho + \eta - \sigma'_1 - v_1, \gamma); \\ (1 - \rho - \sigma'_1 - v'_1, \gamma); \end{matrix} \left(\frac{1}{xs} \right)^\gamma \right]. \end{aligned}$$

THEOREM 13. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\text{Re}(\eta) > 0$ and $\text{Re}(\rho) > \max\{0, \text{Re}(\sigma_1 + \sigma'_1 + v_1 - \eta), \text{Re}(\sigma'_1 - v'_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following integral formula holds true:

$$\begin{aligned} & \int_0^\infty z^{l-1} e^{-\frac{1}{2}\delta z} W_{\tau, \zeta}(\delta z) \left\{ \left(I_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((wtz)^\gamma, s, a; p, q) \right\} \right) (x) \right\} dz \\ &= \frac{x^{\rho+\eta-\sigma_1-\sigma'_1-1}}{\delta^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{wx}{\delta} \right)^\gamma, s, a; p, q \right) \\ & * {}_6\Psi_4 \left[\begin{matrix} (1, 1), (\frac{1}{2} + \zeta + l, \gamma), (\frac{1}{2} - \zeta + l, \gamma), \\ (\frac{1}{2} - \tau + l, \gamma), (\rho + v'_1, \gamma), \\ (\rho, \gamma), (\rho + \eta - \sigma_1 - \sigma'_1 - v_1, \gamma), (\rho + v'_1 - \sigma'_1, \gamma); \\ (\rho + \eta - \sigma_1 - \sigma'_1, \gamma), (\rho + \eta - \sigma'_1 - v_1, \gamma); \end{matrix} \left(\frac{wx}{\delta} \right)^\gamma \right]. \end{aligned}$$

THEOREM 14. Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\text{Re}(\eta) > 0$ and $\text{Re}(\rho) < 1 + \min\{\text{Re}(-v_1), \text{Re}(\sigma_1 + \sigma'_1 - \eta), \text{Re}(\sigma_1 + v'_1 - \eta)\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following integral for-

mula holds true:

$$\begin{aligned} & \int_0^\infty z^{l-1} e^{-\frac{1}{2}\delta z} W_{\tau,\zeta}(\delta z) \left\{ \left(I_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{wz}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) \right\} dz \\ &= \frac{x^{\rho+\eta-\sigma_1-\sigma'_1-1}}{\delta^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{w}{x\delta} \right)^\gamma, s, a; p, q \right) \\ & \quad * {}_6\Psi_4 \left[\begin{matrix} (1, 1), (\frac{1}{2} + \zeta + l, \gamma), (\frac{1}{2} - \zeta + l, \gamma), (1 - \rho - v_1, \gamma), \\ (\frac{1}{2} - \tau + l, \gamma), (1 - \rho, \gamma), \\ (1 - \rho - \eta + \sigma_1 + \sigma'_1, \gamma), (1 - \rho - \eta + \sigma_1 + v'_1, \gamma); \end{matrix} \left(\frac{w}{x\delta} \right)^\gamma \right]. \end{aligned}$$

THEOREM 15. *Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) > \max\{0, \operatorname{Re}(\eta - \sigma_1 - \sigma'_1 - v'_1), \operatorname{Re}(v_1 - \sigma_1)\}$ with $|t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following integral formula holds true:*

$$\begin{aligned} & \int_0^\infty z^{l-1} e^{-\frac{1}{2}\delta z} W_{\tau,\zeta}(\delta z) \left\{ \left(D_{0,x}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v}((wtz)^\gamma), s, a; p, q \right\} \right) (x) \right\} \\ &= \frac{x^{\rho-\eta+\sigma_1+\sigma'_1-1}}{\delta^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{wx}{\delta} \right)^\gamma, s, a; p, q \right) \\ & \quad * {}_6\Psi_4 \left[\begin{matrix} (1, 1), (\frac{1}{2} + \zeta + l, \gamma), (\frac{1}{2} - \zeta + l, \gamma) \\ (\frac{1}{2} - \tau + l, \gamma), (\rho - v_1, \gamma), \\ (\rho + \gamma, \gamma), (\rho - \eta + \sigma_1 + \sigma'_1 + v'_1, \gamma), (\rho - v_1 + \sigma_1, \gamma); \end{matrix} \left(\frac{wx}{\delta} \right)^\gamma \right]. \end{aligned}$$

THEOREM 16. *Let $\sigma_1, \sigma'_1, v_1, v'_1, \eta, \rho, p, q, \lambda, \mu, s \in \mathbb{C}$ with $\gamma \in \mathbb{R}^+$ and $v, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ such that $\operatorname{Re}(\eta) > 0$ and $\operatorname{Re}(\rho) < 1 + \min\{\operatorname{Re}(v'_1), \operatorname{Re}(\eta - \sigma_1 - \sigma'_1, \operatorname{Re}(\eta - \sigma'_1 - v_1))\}$ with $|1/t| < 1$. Then for $\min\{\Re(p), \Re(q)\} \geq 0$, the following integral formula holds true:*

$$\begin{aligned} & \int_0^\infty z^{l-1} e^{-\frac{1}{2}\delta z} W_{\tau,\zeta}(\delta z) \left\{ \left(D_{x,\infty}^{\sigma_1, \sigma'_1, v_1, v'_1, \eta} \left\{ t^{\rho-1} \Phi_{\lambda, \mu; v} \left(\left(\frac{wz}{t} \right)^\gamma, s, a; p, q \right) \right\} \right) (x) \right\} \\ &= \frac{x^{\rho-\eta+\sigma_1+\sigma'_1-1}}{\delta^l} \Phi_{\lambda, \mu; v} \left(\left(\frac{w}{x\delta} \right)^\gamma, s, a; p, q \right) \\ & \quad * {}_6\Psi_4 \left[\begin{matrix} (1, 1), (\frac{1}{2} + \zeta + l, \gamma), (\frac{1}{2} - \zeta + l, \gamma), (1 - \rho - v'_1, \gamma), \\ (\frac{1}{2} - \tau + l, \gamma), (1 - \rho, \gamma), \\ (1 - \rho + \eta - \sigma_1 - \sigma'_1, \gamma), (1 - \rho + \eta - \sigma'_1 - v_1, \gamma); \end{matrix} \left(\frac{w}{x\delta} \right)^\gamma \right]. \end{aligned}$$

The above Theorem 5 to Theorem 16 includes various special cases for which particular choice of parameters reduces to Saigo hypergeometric fractional calculus operators, Riemann-Liouville and Erdélyi-Kober fractional calculus operators. These special cases can be easily evaluated and are left for interesting readers.

4. Concluding remarks and observations

Recently, various developments have been surveyed and studied generalized fractional integrals and differential operators for certain extensions and generalizations of higher transcendental functions, incomplete Hurwitz-Lerch zeta function, general families of double and multiple Hurwitz-Lerch zeta function and other related functions of analytic number theory with applications [28, 31, 33, 34]. In our present investigation, with the help of the concept of the Hadamard product (or the convolution) of two analytic functions, we have obtained the composition formulas of the generalized Marichev-Saigo-Maeda fractional integrals and differential operators (1), (2), (3) and (4) involving the (p, q) -extended Hurwitz-Lerch zeta function $\Phi_{\lambda, \mu; \nu}(z, s, a; p, q)$ in terms of the Hadamard product (8) of the (p, q) -extended Hurwitz-Lerch zeta function $\Phi_{\lambda, \mu; \nu}(z, s, a; p, q)(z, s, a)$ and the Fox–Wright function ${}_r\Psi_s(z)$. Further, we have also deduced the certain image formulas associated with the integral transforms such as Euler-Beta transform, Laplace transform and Whittaker transform. Next, as special cases, we can deduce the certain image formulas for the Saigo's, Riemann-Liouville (R-L) and Erdélyi-Kober (E-K) fractional integral and differential operators as Corollaries, which we have left as exercise for readers. The results obtained in this paper are assumed to be have applications in various field of Physical and Engineering Sciences of extended version of generalized fractional integrals and differential operators and integral transforms related papers [4, 5, 6, 7, 8, 19, 21, 22, 23, 29, 30, 32, 35, 36].

Acknowledgements. The authors thanks to the anonymous referee for valuable comments and useful suggestions to improve the quality of the paper in present form.

REFERENCES

- [1] R. P. AGARWAL, A. KILIÇMAN, R. K. PARMAR AND A. K. RATHIE, *Certain generalized fractional calculus formulas and integral transforms involving (p, q) -Mathieu-type series*, Adv. Differ. Equ. **221**, (2019), 1–11, <https://doi.org/10.1186/s13662-019-2142-0>.
- [2] M. A. CHAUDHRY, A. QADIR, H. M. SRIVASTAVA AND R. B. PARIS, *Extended hypergeometric and confluent hypergeometric functions*, Appl. Math. Comput. **159** (2004), 589–602.
- [3] J. CHOI, A. K. RATHIE, R. K. PARMAR, *Extension of extended beta, hypergeometric and confluent hypergeometric functions*, Honam Math. J. **36** 2 (2014), 339–367.
- [4] J. CHOI AND R. K. PARMAR, *The Incomplete Lauricella and Fourth Appell Functions*, Far East Journal of Mathematical sciences, **96**, (2015), 315–328.
- [5] J. CHOI, RAKESH K. PARMAR, P. CHOPRA, *Extended Mittag-Leffler function and associated fractional calculus operators*, Georgian Mathematical Journal, **27** 2 (2020), 199–209, <https://doi.org/10.1515/gmj-2019-2030>.
- [6] JUNGSEANG CHOI, RAKESH K. PARMAR AND P. CHOPRA, *The Incomplete Srivastava's Triple Hypergeometric Functions γ_B^H and Γ_B^H* , Filomat, **30** (2016), 1779–1787, <https://doi.org/10.2298/FIL1607779C>.
- [7] J. CHOI AND RAKESH K. PARMAR, *The Incomplete Srivastava's Triple Hypergeometric Functions γ_A^H and Γ_A^H* , Miskolc Mathematical Notes, **19** (2018), 191–200, <https://doi.org/10.18514/MMN.2018.1503>.
- [8] J. CHOI AND RAKESH K. PARMAR, *An extension of the generalized Hurwitz-Lerch Zeta function of two variables*, Filomat, **31** 1 (2017), 91–96, <https://doi.org/10.2298/FIL1701091>.
- [9] J. CHOI AND RAKESH K. PARMAR, *Fractional Integration And Differentiation of the (p, q) -extended Bessel function*, Bulletin of the Korean Mathematical Society, **55** 2 (2018), 599–610, <https://doi.org/10.4134/BKMS.b170193>.

- [10] J. CHOI AND RAKESH K. PARMAR, *Fractional calculus of the (p, q) -extended Struve function*, Far East Journal of Mathematical Sciences, **103** 2 (2018), 541–559, <http://dx.doi.org/10.17654/MS103020541>.
- [11] J. CHOI, R. K. PARMAR AND T. K. POGÁNY, *Mathieu-type series built by (p, q) -extended Gaussian hypergeometric function*, Bull. Korean Math. Soc. **54** 3 (2017), 789–797, <https://doi.org/10.4134/BKMS.b160313>.
- [12] M. J. LUO, R. K. PARMAR AND R. K. RAINA, *On extended Hurwitz-Lerch zeta function*, J. Math. Anal. Appl. **448** (2017), 12811–1304.
- [13] D. JANKOV MAŠIREVIĆ, RAKESH K. PARMAR AND T. K. POGÁNY, *(p, q) -extended Bessel and modified Bessel functions of the first kind*, Results in Mathematics, **72** (2017), 617–632, <https://doi.org/10.1007/s00025-016-0649-1>.
- [14] A. A. KILBAS, H. M. SRIVASTAVA, AND J. J. TRUJILLO, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematical Studies, vol. **204**, Elsevier (North-Holland) Science Publishers, Amsterdam, London and New York, 2006.
- [15] V. KIRYAKOVA, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series, 301, Longman Scientific and Technical, Harlow; copublished in the United States with John Wiley and Sons, Inc., New York (1994).
- [16] A. M. MATHAI, R. K. SAXENA AND H. J. HAUBOLD, *The H -Functions: Theory and Applications*, Springer, New York, 201.
- [17] O. I. MARICHEV, *Volterra Equation of Mellin Convolution Type with a Horn Function in the Kernel*, Izv. AN BSSR Ser. Fiz.-Mat. Nauk. **1**, 128–129 (1974) [in Russian].
- [18] F. W. J. OLVER, D. W. LOZIER, R. F. BOISVERT AND C. W. CLARK (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.
- [19] R. K. PARMAR AND R. K. SAXENA, *Incomplete extended Hurwitz-Lerch Zeta functions and associated properties*, Communications of the Korean Mathematical Society, **32** 2 (2017), 287–304.
- [20] RAKESH K. PARMAR, T. K. POGÁNY AND R. K. SAXENA, *On properties and applications of (p, q) -extended τ -hypergeometric functions*, Comptes rendus. Mathématique, **356** 3 (2018), 278–282, <https://doi.org/10.1016/j.crma.2017.12.014>.
- [21] R. K. PARMAR AND T. K. POGÁNY, *Extended Srivastava's triple hypergeometric $H_{A,p,q}$ function and related bounding inequalities*, J. Contemp. Math. Anal. **52** 6 (2017), 261–272, <https://doi.org/10.3103/S1068362317060036>.
- [22] R. K. PARMAR AND R. K. SAXENA, *The Incomplete Generalized τ -Hypergeometric and Second τ -Appell Functions*, J. Korean Math. Soc., **53** (2016), 363–379, <https://doi.org/10.4134/JKMS.2016.53.2.363>.
- [23] R. K. PARMAR AND R. K. SAXENA, *Incomplete Extended Hurwitz-Lerch Zeta Functions and Associated Properties*, Communication of Korean Mathematical Society, **32** (2017), 287–304, <https://doi.org/10.4134/CKMS.c150227>.
- [24] S. G. SAMKO, A. A. KILBAS, AND O. I. MARICHEV, *Fractional Integrals and Derivatives: Theory and Applications*, Translated from the Russian: *Integrals and Derivatives of Fractional Order and Some of Their Applications* (Nauka i Tekhnika, Minsk, 1987); Gordon and Breach Science Publishers: Reading, UK, 1993.
- [25] I. N. SNEDDON, *The use of the Integral Transforms*, Tata McGraw-Hill, New Delhi, 1979.
- [26] M. SAIGO, *On Generalized Fractional Calculus Operators*, in Recent Advances in Applied Mathematics, Proceedings of the International Workshop held at Kuwait University (Kuwait; May 4–7, 1996), Kuwait University, Department of Mathematics and Computer Science, Kuwait, 441–450 (1996).
- [27] M. SAIGO AND N. MAEDA, *More Generalization of Fractional Calculus*, in Transform Methods and Special Functions, Proceedings of the Second International Workshop Dedicated to the 100th Anniversary of the Birth of Nikola Obreschkoff (Varna; August 23–30, 1996) (P. Rusev, I. Dimovski and V. Kiryakova, Editors) (Bulgarian Academy of Sciences, Institute of Mathematics and Informatics, Sofia, 1998) pp. 386–400.
- [28] H. M. SRIVASTAVA, *Some general families of integral transformations and related results*, Appl. Math. Comput. Sci. **6**, (2022), 27–41.
- [29] H. M. SRIVASTAVA, *A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics*, Symmetry, **13** (2021), Article ID 2294, 1–22.

- [30] H. M. SRIVASTAVA, *An introductory overview of fractional-calculus operators based upon the Fox-Wright and related higher transcendental functions*, J. Adv. Engrg. Comput. **5** (2021), 135–166.
- [31] H. M. SRIVASTAVA, *Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations*, J. Nonlinear Convex Anal., **22** (2021), 1501–1520.
- [32] H. M. SRIVASTAVA, *Some general families of the Hurwitz-Lerch Zeta functions and their applications: Recent developments and directions for further researches*, Proc. Inst. Math. Mech. Nat. Acad. Sci. Azerbaijan, **45** (2019), 234–269.
- [33] H. M. SRIVASTAVA, AND P. AGARWAL, *Certain Fractional Integral Operators and the Generalized Incomplete Hypergeometric Functions*, Applications and Applied Mathematics: An International Journal (AAM), **8**, (2013), 333–345.
- [34] H. M. SRIVASTAVA, E. S. A. ABU JARAD, F. JARAD, G. SRIVASTAVA, AND M. H. A. ABU JARAD, *The Marichev-Saigo-Maeda fractional-calculus operators involving the (p, q) -extended Bessel and Bessel-Wright functions*, Fractal Fract. **5** (2021), Article ID 210, 1–15.
- [35] H. M. SRIVASTAVA, R. C. S. CHANDEL, AND H. KUMAR, *Some general Hurwitz-Lerch type zeta functions associated with the Srivastava-Daoust multiple hypergeometric functions*, J. Nonlinear Var. Anal., **6**, (2022), 299–315.
- [36] H. M. SRIVASTAVA, R. ŞAHİN, AND O. YAĞCI, *A family of incomplete Hurwitz-Lerch zeta functions of two variables*, Miskolc Math. Notes, **21** (2020), 401–415.
- [37] H. M. SRIVASTAVA AND P. W. KARLSSON, *Multiple Gaussian Hypergeometric Series*, Halsted Press, (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.
- [38] H. M. SRIVASTAVA AND R. K. SAXENA, *Operators of fractional integration and their applications*, Applied Mathematics and Computation, **118**, (2001), 1–52.

(Received September 3, 2023)

Purnima Chopra
Formerly Department of Mathematics
Marudhar Engineering College
Bikaner-334001, Rajasthan, India
e-mail: purnimachopra05@gmail.com

Shilpa Kumawat
Department of Mathematics
Career Point University
Kota-325003, Rajasthan, India
e-mail: shilpaprajapat786@gmail.com

Hemlata Saxena
Department of Mathematics
Career Point University
Kota-325003, Rajasthan, India
e-mail: saxenadrhemlata@gmail.com