

## MULTIVALENT FUNCTIONS WITH VARYING ARGUMENTS

R. M. EL-ASHWAH, M. K. AOUF, A. A. M. HASSAN AND A. H. HASSAN

**Abstract.** Silverman [4] was defined the class of univalent functions  $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$  for which  $\arg(a_k)$  prescribed in such way that  $f(z)$  is univalent if and only if  $f(z)$  is starlike. In this paper we introduce the subclass of  $p$ -valent functions with varying arguments, especially  $p$ -valent starlike functions and  $p$ -valent convex functions, moreover we give some interesting properties of functions in these classes, including coefficients estimates, distortion theorems and extreme functions.

*Mathematics subject classification (2010):* 30C45.

*Keywords and phrases:* Analytic, varying arguments, extreme points.

### REFERENCES

- [1] M. K. AOUF, *A generalization of multivalent functions with negative coefficients*, J. Korean Math. Soc., 25 (1988), no. 1, 53–66.
- [2] S. OWA, *On new classes of  $p$ -valent function with negative coefficients*, Simon Stevin 59 (1985), no. 4, 385–402.
- [3] D. A. PATIL AND N. K. THAKARE, *On convex hulls and extreme points of  $p$ -valent starlike and convex classes with applicents*, Bull. Math. Soc. Sci. Math. R. S. Roumaine (N. S.), 27 (1983), no. 75, 145–160.
- [4] H. SILVERMAN, *Univalent functions with varying arguments*, Houston J. Math., 17 (1981), 283–287.