

## A LOGARITHMIC MEAN AND INTERSECTIONS OF OSCULATING HYPERPLANES IN $R^n$

ALAN HORWITZ

*Abstract.* We discuss a special case of the means defined in [1]. Let  $C$  be the curve in  $R^n$  with vector equation  $\hat{\alpha}(t) = \langle t, t \log t, \dots, t(\log t)^{n-1} \rangle$ . Let  $0 < a_1 < \dots < a_n$  and let  $O_k$  be the osculating hyperplane to  $C$  at  $a_k$ . Then we show that  $O_1, \dots, O_n$  have a unique point of intersection,  $P = (i_1, \dots, i_n) \in R^n$ , and in particular,  $i_1$  equals the mean

$$M(a_1, \dots, a_n) = (n-1)! \sum_{j=1}^n \frac{a_j}{\prod_{\substack{i=1 \\ i \neq j}}^n (\ln a_j - \ln a_i)},$$

the logarithmic mean of Neuman.

*Mathematics subject classification (2010):* 25E60, 26B99.

*Keywords and phrases:* logarithmic mean; osculating hyperplane; Wronskian.

### REFERENCES

- [1] A. HORWITZ, *Means, Generalized Divided Differences, and Intersections of Osculating Hyperplanes*, J. Math. Anal. Appl. **200** (1996), 126–148.
- [2] JORMA K. MERIKOSKI, *Extending Means Of Two Variables To Several Variables*, JIPAM, Volume 5, Issue 3, Article 65, 2004.
- [3] S. MUSTONEN, *Logarithmic mean for several arguments*, (2002), [<http://www.survo.fi/papers/logmean.pdf>].
- [4] EDWARD NEUMAN, *The Weighted Logarithmic Mean*, J. Math. Anal. Appl. **188** (1994), 885–900.
- [5] A. O. PITTENGER, *The logarithmic mean in  $n$  variables*, Amer. Math. Monthly **92** (1985), 99–104.
- [6] K. STOLARSKY, *Generalizations of the logarithmic mean*, Math. Mag. **48** (1975), 87–92.
- [7] ZHEN-GANG XIAO AND ZHI-HUA ZHANG, *The Inequalities  $G \leq L \leq I \leq A$  in  $n$  Variables*, JIPAM, Volume 4, Issue 2, Article 39, 2003.