

ON SOME GENERALIZATIONS OF ENESTRÖM-KAKEYA THEOREM

NISAR A. RATHER

Abstract. In this paper, we obtain some generalizations of a well-known result of Eneström-Kakeya concerning the bounds for the moduli of the zeros of polynomials with complex coefficients which improve some known results.

1. Introduction

If $P(z) = \sum_{j=0}^n a_j z^j$ is a polynomial of degree n with real coefficients satisfying

$$a_n \geq a_{n-1} \geq \cdots \geq a_1 \geq a_0 > 0$$

then according to a well-known result of Eneström-Kakeya (see [21, 22, 23]) all the zeros of $P(z)$ lie in $|z| \leq 1$.

We may apply this result to $P(tz)$ to obtain following more general result.

THEOREM A. *If $P(z) = \sum_{j=0}^n a_j z^j$ is a polynomial of degree n with real coefficients satisfying*

$$t^n a_n \geq t^{n-1} a_{n-1} \geq \cdots \geq t a_1 \geq a_0 > 0,$$

for some $t > 0$. Then all the zeros of $P(z)$ lie in $|z| \leq t$.

In literature [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25] there exists several extensions of Eneström-Kakeya theorem. An exhaustive survey on the Eneström-Kakeya theorem and some of its generalizations is given in [15] by Gardner and Govil. For the polynomials with complex coefficients, A. Aziz and Q. G. Mohammad [5] used matrix method and proved among others, the following generalization of Theorem A.

THEOREM B. *If $P(z) = \sum_{j=0}^n a_j z^j$ is a polynomial of degree n with complex coefficients such that for $k = 0, 1, 2, \dots, n$ and for some $t > 0$,*

$$t^n |a_n| \leq t^{n-1} |a_{n-1}| \leq \cdots \leq t^k |a_k| \geq t^{k-1} |a_{k-1}| \geq \cdots \geq t |a_1| \geq |a_0|,$$

then $P(z)$ has all its zeros in the circle

$$|z| \leq t \left\{ \frac{2t^k |a_k|}{t^n |a_n|} - 1 \right\} + 2 \sum_{j=0}^n \frac{|a_j - |a_j||}{|a_n| t^{n-j-1}}. \quad (1)$$

A. Aziz and Q. G. Mohammad [5] also proved the following result.

Mathematics subject classification (2010): 26C10, 30C15.

Keywords and phrases: Zeros, polynomials, complex domain.

THEOREM C. Let $P(z) = \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with complex coefficients. If $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$ for $j = 0, 1, 2, \dots, n$. If $t > 0$ can be found such that

$$t^n \alpha_n \leq t^{n-1} \alpha_{n-1} \leq \dots \leq t^{k+1} \alpha_{k+1} \leq t^k \alpha_k, \quad t^k \alpha_k \geq t^{k-1} \alpha_{k-1} \geq \dots \geq t \alpha_1 \geq \alpha_0$$

and

$$t^n \beta_n \leq t^{n-1} \beta_{n-1} \leq \dots \leq t^{m+1} \beta_{m+1} \leq t^m \beta_m, \quad t^m \beta_m \geq t^{m-1} \beta_{m-1} \geq \dots \geq t \beta_1 \geq \beta_0,$$

$0 \leq k, m \leq n$, $a_{-1} = a_{n+1} = 0$, $\alpha_n > 0$, then all the zero of $P(z)$ lie in circle

$$|z| \leq \frac{2}{|a_n| t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{t(\alpha_n + \beta_n)}{|a_n|}. \quad (2)$$

In this paper, we first present the following result which among other things provides a refinement of Theorem B for $0 \leq k \leq n-1$.

THEOREM 1. Let $P(z) = \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with complex coefficients. If $t > 0$ can be found such that

$$t^n |a_n| \leq t^{n-1} |a_{n-1}| \leq \dots \leq t^{k+1} |a_{k+1}| \leq t^k |a_k|, \quad t^k |a_k| \geq t^{k-1} |a_{k-1}| \geq \dots \geq t |a_1| \geq |a_0|,$$

$0 \leq k \leq n-1$, then all the zero of $P(z)$ lie in

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| \leq \frac{2t^{k+1} |a_k|}{t^n |a_n|} + 2 \sum_{v=0}^n \frac{|a_v - a_{v-1}|}{t^{n-v-1} |a_n|} - \frac{|a_{n-1}| + |a_{n-1} - a_{n-2}| + \dots + t |a_n - a_{n-1}|}{|a_n|}. \quad (3)$$

Proof. Consider the polynomial

$$\begin{aligned} F(z) &= (t-z)P(z) = (t-z)(a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0) \\ &= -a_n z^{n+1} + (ta_n - a_{n-1}) z^n + \sum_{v=0}^{n-1} (ta_v - a_{v-1}) z^v \quad (a_{-1} = 0). \end{aligned}$$

Let $|z| > t$, then

$$\begin{aligned} |F(z)| &\geq |a_n| |z|^n \left[\left| z + \frac{a_{n-1}}{a_n} - t \right| - \sum_{v=0}^{n-1} \left| \frac{ta_v - a_{v-1}}{a_n} \right| \frac{1}{|z|^{n-v}} \right] \\ &> |a_n| |z|^n \left[\left| z + \frac{a_{n-1}}{a_n} - t \right| - \frac{1}{|a_n|} \sum_{v=0}^{n-1} \frac{|ta_v - a_{v-1}|}{t^{n-v}} \right]. \end{aligned} \quad (4)$$

Now,

$$\begin{aligned}
\sum_{v=0}^{n-1} \frac{|ta_v - a_{v-1}|}{t^{n-v}} &\leqslant \sum_{v=0}^{n-1} \frac{|t| |a_v| - |a_{v-1}|}{t^{n-v}} + \sum_{v=0}^{n-1} \frac{t |a_v - |a_v|| + |a_{v-1} - |a_{v-1}||}{t^{n-v}} \\
&= \sum_{v=0}^k \frac{|t| |a_v| - |a_{v-1}|}{t^{n-v}} + \sum_{v=k+1}^{n-1} \frac{|t| |a_v| - |a_{v-1}|}{t^{n-v}} \\
&\quad + \sum_{v=0}^{n-1} \frac{t |a_v - |a_v|| + |a_{v-1} - |a_{v-1}||}{t^{n-v}} \\
&\leqslant \frac{2|a_k|}{t^{n-k-1}} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}} - \left(|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t |a_n - |a_n|| \right).
\end{aligned}$$

Using this in (4), we have

$$\begin{aligned}
|F(z)| &\geqslant |a_n| |z|^{n+1} \left[\left| z + \frac{a_{n-1}}{a_n} - t \right| - \frac{2|a_k|}{t^{n-k-1} |a_n|} - 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1} |a_n|} \right. \\
&\quad \left. + \frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t |a_n - |a_n||}{|a_n|} \right] > 0,
\end{aligned}$$

if

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| > \frac{2|a_k|}{t^{n-k-1} |a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1} |a_n|} - \frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t |a_n - |a_n||}{|a_n|}.$$

Therefore, it follows that all the zeros of $F(z)$ whose modulus is greater than t lie in circle

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| \leqslant \frac{2|a_k|}{t^{n-k-1} |a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1} |a_n|} - \frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t |a_n - |a_n||}{|a_n|}. \quad (5)$$

We now show that all those zeros of $F(z)$ whose modulus is less than or equal to t also satisfy (5). Let $|z| \leqslant t$,

$$\begin{aligned}
\left| z + \frac{a_{n-1}}{a_n} - t \right| &\leqslant t + \left| \frac{a_{n-1}}{a_n} - t \right| \leqslant t + \frac{|t| |a_n| - |a_{n-1}|}{|a_n|} + \frac{t |a_n - |a_n|| + |a_{n-1} - |a_{n-1}||}{|a_n|} \\
&= t + \frac{|a_{n-1}| - t |a_n|}{|a_n|} + \frac{|a_n - |a_n|| t + |a_{n-1} - |a_{n-1}||}{|a_n|} \\
&= \frac{2|a_{n-1}|}{|a_n|} + \frac{|a_{n-1} - |a_{n-1}||}{|a_n|} + \frac{t |a_n - |a_n||}{|a_n|} - \frac{|a_{n-1}|}{|a_n|}.
\end{aligned} \quad (6)$$

Now, by hypothesis,

$$\left| \frac{a_{n-1}}{a_n} \right| \leqslant \frac{|a_k|}{|a_n| t^{n-k-1}}, \quad \text{for } 0 \leqslant k \leqslant n-1. \quad (7)$$

Using (7) in (6), we obtain for $0 \leq k \leq n-1$,

$$\begin{aligned} \left| z + \frac{a_{n-1}}{a_n} - t \right| &\leq \frac{2|a_k|}{|a_n|t^{n-k-1}} + \frac{|a_{n-1} - |a_{n-1}||}{|a_n|} + \frac{t|a_n - |a_n||}{|a_n|} - \frac{|a_{n-1}|}{|a_n|} \\ &\leq \frac{2|a_k|}{t^{n-k-1}|a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}|a_n|} - \frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t|a_n - |a_n||}{|a_n|}. \end{aligned}$$

Thus, we have shown that if $|z| \leq t$, then for $0 \leq k \leq n-1$,

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| \leq \frac{2|a_k|}{t^{n-k-1}|a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}|a_n|} - \frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t|a_n - |a_n||}{|a_n|}.$$

This shows that all the zeros of $F(z)$ whose moduli is less than or equal to t also lie in the circle defined by (5). But all the zeros of $P(z)$ are also the zeros of $F(z)$, therefore, we conclude that all the zeros of $P(z)$ lie in the circle defined by (5). This completes the proof of Theorem 1. \square

REMARK 1. In general Theorem 1 gives much better result than Theorem B for $0 \leq k \leq n-1$. To see this, we show that the circle defined by (3) is contained in the circle defined by (1). Let z be any point belonging to the circle defined by (3), then

$$\begin{aligned} \left| z + \frac{a_{n-1}}{a_n} - t \right| &\leq \frac{2|a_k|}{t^{n-k-1}|a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}|a_n|} \\ &\quad - \left(\frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t|a_n - |a_n||}{|a_n|} \right). \end{aligned}$$

This implies

$$\begin{aligned} |z| &= \left| z + \frac{a_{n-1}}{a_n} - t + t - \frac{a_{n-1}}{a_n} \right| \leq \left| z + \frac{a_{n-1}}{a_n} - t \right| + \left| t - \frac{a_{n-1}}{a_n} \right| \\ &\leq \frac{2|a_k|}{t^{n-k-1}|a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}|a_n|} - \left(\frac{|a_{n-1}| + |a_{n-1} - |a_{n-1}|| + t|a_n - |a_n||}{|a_n|} \right) \\ &\quad + \frac{|a_n|t - |a_{n-1}|}{|a_n|} + \frac{|a_n - |a_n||t + |a_{n-1} - |a_{n-1}||}{|a_n|} \\ &= \frac{2|a_k|}{t^{n-k-1}|a_n|} + 2 \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}|a_n|} - \left(\frac{|a_{n-1}| + t|a_n - |a_n||}{|a_n|} \right) \\ &\quad + \frac{|a_{n-1} - |a_n||t}{|a_n|} + \frac{|a_n - |a_n||t}{|a_n|} \\ &\leq \frac{2t|a_k|}{t^{n-k}|a_n|} + \frac{2}{|a_n|} \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}} - t = t \left\{ \frac{2t^k|a_k|}{t^n|a_n|} - 1 \right\} + \frac{2}{|a_n|} \sum_{v=0}^n \frac{|a_v - |a_v||}{t^{n-v-1}}. \end{aligned}$$

This shows that the point z belongs to the circle defined by (1). Hence the circle defined by (3) is contained in the circle defined by (1).

For polynomial $P(z) = \sum_{j=0}^n a_j z^j$ with real and positive coefficients, we obtain the following result.

COROLLARY 1. Let $P(z) = \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with real and positive coefficients. If $t > 0$ can be found such that

$$t^n a_n \leq t^{n-1} a_{n-1} \leq \cdots \leq t^{k+1} a_{k+1} \leq t^k a_k, \quad t^k a_k \geq t^{k-1} a_{k-1} \geq \cdots \geq t a_1 \geq a_0,$$

$0 \leq k \leq n-1$, then all the zero of $P(z)$ lie in

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| \leq \frac{2t^{k+1} a_k}{t^n a_n} - \frac{a_{n-1}}{a_n}.$$

Next, we present the following result which improves the bound of Theorem C for $0 \leq k, m \leq n-1$.

THEOREM 2. Let $P(z) = \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with complex coefficients. If $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$ for $j = 0, 1, 2, \dots, n$. If $t > 0$ can be found such that

$$0 < t^n \alpha_n \leq t^{n-1} \alpha_{n-1} \leq \cdots \leq t^{k+1} \alpha_{k+1} \leq t^k \alpha_k, \quad t^k \alpha_k \geq t^{k-1} \alpha_{k-1} \geq \cdots \geq t \alpha_1 \geq \alpha_0 > 0$$

and

$$0 < t^n \beta_n \leq t^{n-1} \beta_{n-1} \leq \cdots \leq t^{m+1} \beta_{m+1} \leq t^m \beta_m, \\ t^m \beta_m \geq t^{m-1} \beta_{m-1} \geq \cdots \geq t \beta_1 \geq \beta_0 > 0,$$

$0 \leq k, m \leq n-1$, $a_{-1} = a_{n+1} = 0$, then all the zero of $P(z)$ lie in

$$\left| z + \frac{a_{n-1} - t a_n}{a_n} \right| \leq \frac{2}{|a_n| t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}. \quad (8)$$

Proof. Consider the polynomial

$$F(z) = (t-z)P(z) = (t-z)(a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0) \\ = -a_n z^{n+1} + (ta_n - a_{n-1}) z^n + \sum_{v=0}^{n-1} (ta_v - a_{v-1}) z^v \quad (a_{-1} = 0).$$

Let $|z| > t$, then

$$|F(z)| \geq |z|^n \left[|a_n z + a_{n-1} - t a_n| - \sum_{v=0}^{n-1} |ta_v - a_{v-1}| \frac{1}{|z|^{n-v}} \right] \\ > |z|^n \left[|a_n z + a_{n-1} - t a_n| - \frac{1}{t^n} \sum_{v=0}^{n-1} |ta_v - a_{v-1}| t^v \right]. \quad (9)$$

Now by hypothesis

$$\begin{aligned}
 \sum_{v=0}^{n-1} |ta_v - a_{v-1}| t^v &\leq \sum_{v=0}^{n-1} |t\alpha_v - \alpha_{v-1}| t^v + \sum_{v=0}^{n-1} |t\beta_v - \beta_{v-1}| t^v \\
 &= \sum_{v=0}^k |t\alpha_v - \alpha_{v-1}| t^v + \sum_{v=k+1}^{n-1} |t\alpha_v - \alpha_{v-1}| t^v + \sum_{v=0}^m |t\beta_v - \beta_{v-1}| t^v \\
 &\quad + \sum_{v=m+1}^{n-1} |t\beta_v - \beta_{v-1}| t^v \\
 &= 2\alpha_k t^{k+1} + 2\beta_m t^{m+1} - (\alpha_{n-1} + \beta_{n-1}) t^n.
 \end{aligned}$$

Using this in (9), we obtain

$$|F(z)| \geq |z|^{n+1} \left\{ |a_n z + a_{n-1} - ta_n| - 2\alpha_k \frac{t^{k+1}}{t^n} - 2\beta_m \frac{t^{m+1}}{t^n} + (\alpha_{n-1} + \beta_{n-1}) \right\} > 0,$$

if

$$\left| z + \frac{a_{n-1} - ta_n}{a_n} \right| > \frac{2}{|a_n| t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}.$$

Hence all the zeros of $F(z)$ whose modulus is greater than t lie in the circle

$$\left| z + \frac{a_{n-1} - ta_n}{a_n} \right| \leq \frac{2}{|a_n| t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}. \quad (10)$$

Now, if $|z| \leq t$, then we have

$$\begin{aligned}
 |a_n z + a_{n-1} - ta_n| &\leq |a_n| t + |a_{n-1} - ta_n| \leq t \alpha_n + t \beta_n + (\alpha_{n-1} - t \alpha_n) + (\beta_{n-1} - t \beta_n) \\
 &= \alpha_{n-1} + \beta_{n-1},
 \end{aligned}$$

this gives,

$$\left| z + \frac{a_{n-1}}{a_n} - t \right| \leq \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}. \quad (11)$$

By hypothesis for $0 \leq k, m \leq n-1$,

$$t^{n-1} \alpha_{n-1} \leq \alpha_k t^k \quad \text{and} \quad t^{n-1} \beta_{n-1} \leq \beta_m t^m.$$

Therefore,

$$2(t^{n-1} \alpha_{n-1} + t^{n-1} \beta_{n-1}) \leq 2(\alpha_k t^k + \beta_m t^m).$$

Equivalently,

$$\alpha_{n-1} + \beta_{n-1} \leq \frac{2}{t^n} (\alpha_k t^{k+1} + \beta_m t^{m+1}) - (\alpha_{n-1} + \beta_{n-1}). \quad (12)$$

Using (12) in (11), we obtain for $0 \leq k \leq n-1$,

$$\left| z + \frac{a_{n-1} - ta_n}{a_n} \right| \leq \frac{2}{|a_n|t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}.$$

This shows that all the zeros of $F(z)$ whose modulus is less than or equal to t also satisfy the inequality (10). Thus we conclude that all the zeros of $F(z)$ and hence that of $P(z)$ lie in the circle defined by (10). This completes the proof of Theorem 2. \square

REMARK 2. In general Theorem 2 also gives much better result than Theorem B for $0 \leq k \leq n-1$. To see this, we show that the circle defined by (8) is contained in the circle defined by (2). Let z be any point belonging to the circle defined by (8), then

$$\left| z + \frac{a_{n-1} - ta_n}{a_n} \right| \leq \frac{2}{|a_n|t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|}.$$

This implies

$$\begin{aligned} |z| &= \left| z + \frac{a_{n-1} - ta_n}{a_n} - \frac{a_{n-1} - ta_n}{a_n} \right| \leq \left| z + \frac{a_{n-1} - ta_n}{a_n} \right| + \left| \frac{a_{n-1} - ta_n}{a_n} \right| \\ &\leq \frac{2}{|a_n|t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{\alpha_{n-1} + \beta_{n-1}}{|a_n|} + \frac{a_{n-1} - ta_n}{|a_n|} + \frac{\beta_{n-1} - t\beta_n}{|a_n|} \\ &= \frac{2}{|a_n|t^n} \left\{ \alpha_k t^{k+1} + \beta_m t^{m+1} \right\} - \frac{t(\alpha_n + \beta_n)}{|a_n|}, \end{aligned}$$

which shows that the point z belongs to the circle defined by (2). Hence the circle defined by (8) is contained in the circle defined (2).

Acknowledgement. The author is highly grateful to the referee for his useful suggestions.

REFERENCES

- [1] C. AFFANE-AJI, N. AGARWAL AND N. K. GOVIL, *Location of zeros of polynomials*, Mathematical and Computer Modelling, **50** (2009), 306–313.
- [2] C. AFFANE-AJI AND N. K. GOVIL, *On the regions containing all the zeros of a polynomial*, Nonlinear Analysis: Stability, Approximation and Inequalities, Edited by P. G. Georgiev, P. Pardalos and H. M. Srivastava, Springer, Optimization and its Applications, **68** (2012), 39–55.
- [3] N. ANDERSON, E. B. SAFF, R. S. VARGA, *On the Eneström-Kakeya theorem and its sharpness*, Linear Algebra Appl., **28** (1979), 5–16.
- [4] N. ANDERSON, E. B. SAFF, R. S. VARGA, *An extension of the Eneström-Kakeya theorem and its sharpness*, SIAM J. Math. Anal., **12** (1981), 10–22.
- [5] A. AZIZ AND Q. G. MOHAMMAD, *Zero free regions for polynomials and some generalizations of Eneström-Kakeya theorem*, Canad. Math. Bull., **27**(3) (1984), 265–272.
- [6] A. AZIZ AND W. M. SHAH, *On the zeros of polynomial and related analytic functions*, Glas. Mat., **3**(53) (1998), 173–184.
- [7] A. AZIZ AND B. A ZARGAR, *Some extensions of Eneström-Kakeya theorem*, Glas. Mat., **33** (1996), 239–244.

- [8] A. AZIZ AND B. A ZARGAR, *Bounds for the zeros of a polynomial with restricted coefficients*, Appl. Math., **3** (2012), 3–33.
- [9] J. CAO AND R. GARDNER, *Restrictions on the zeros of as a consequence of conditions on the coefficients of even powers and odd powers of the variable*, J. Comput. Appl. Math., **155** (2003), 153–163.
- [10] A. CAUCHY, *Exercices de mathématique*, Oeuvres,(2) Volume **9**, p. 122 (1829).
- [11] G. T. CARGO AND O. SHISHA, *Zeros of polynomials and fractional order differences of their coefficients*, J. Math. Anal. Appl., **7** (1963) 176–182.
- [12] Y. CHOO AND G. K. CHOI, *On the robustness property of Eneström-Kakeya theorem*, Int. J. Math. Anal., **5**(42) (2011), 2089–2096.
- [13] K. K. DEWAN AND M. BIDKHAM, *On the Eneström-Kakeya theorem*, J. Math. Anal. Appl., **180** (1993), 29–36.
- [14] R. B. GARDNER AND N. K. GOVIL, *Some generalizations of Eneström-Kakeya theorem*, Acta Math. Hungar., **74**(1-2) (1997), 125–134.
- [15] R. B. GARDNER AND N. K. GOVIL, *The Eneström-Kakeya theorem and some of its generalizations*, in S. Joshi, M. Dorff and I. Lahiri(Eds), Current Topics in Pure and Computational Complex Analysis, New Delhi; Springer- Verlag (2014), 171–200.
- [16] M. KOVAČEVIĆ AND I. MILOVANOVIĆ, *On a generalization of the Eneström-Kakeya theorem*, Pure Math. and Applic. Ser. A, **3** (1992), 543–547.
- [17] V. K. JAIN, *On the zeros of polynomials*, Proc. Indian Acad. Sci. Math. Sci., **119**(1) (2009), 37–43.
- [18] A. JOYAL, G. LABELLE AND Q. I. RAHMAN, *On the locations of zeros of polynomials*, Canad. Math. Bull., **10**(1) (1967), 53–63.
- [19] P. V. KRISHNAIAH, *On the Eneström-Kakeya theorem*, J. London Math. Soc., **30** (1955) 314–19.
- [20] A. LIN, B. HUANG, J. CAO, R. GARDNER, *Some results on the location of zeros of analytic functions*, Chinese J. Engrg. Math., **21**(5) (2004), 785–791.
- [21] M. MARDEN, *Geometry of polynomials*, Math. Surveys No. 3, Amer. Math. Soc., Providence R. I. 1966.
- [22] G. V. MILOVANOVIĆ, D. S. MITRINOVİĆ AND TH. M. RASSIAS, *Topics in polynomials: extremal properties, inequalities, zeros*, World scientific Publishing Co., Singapore, (1994).
- [23] Q. I. RAHMAN AND G. SCHMEISSER, *Analytic theory of polynomial*, Clarendon Press, Oxford, 2002.
- [24] N. A. RATHER, S. AHMAD AND M. I. MIR, *On the Eneström-Kakeya theorem*, Int. J. Pure Appl. Math., **41** (2007), 807–816.
- [25] N. A. RATHER, S. AHMAD, *A remark on generalizations of Eneström-Kakeya theorem*, Math. Balkanica., **21**, Fasc. 3-4(2007), 319–328.

(Received May 24, 2016)

Nisar A. Rather
 P.G. Department of Mathematics
 Kashmir University
 Hazratbal, Srinagar-190006, India
 e-mail: dr.narather@gmail.com