

CRITERIA FOR NORMAL AND φ -NORMAL FUNCTIONS

BANARSI LAL*, VIRENDER SINGH AND PRIYANKA

Abstract. We establish the following result: Let $f(z)$ be meromorphic in the unit disk \mathbb{D} , and let $S = \{\alpha_1, \alpha_2, \alpha_3\}$ be a set of three distinct complex numbers in \mathbb{C} and $A > 0$. If $f(z) \in S$ implies $|L(f)| < A$, then $f(z)$ is a normal function on \mathbb{D} . The result is further extended to φ -normal functions.

Mathematics subject classification (2020): 30D35, 30D45.

Keywords and phrases: Normal function, meromorphic function, differential inequalities.

REFERENCES

- [1] A. AULASKARI AND J. RATTYA, *Properties of meromorphic φ -normal function*, J. Michigan. Math. **60** (2011), 93–111.
- [2] Q. CHEN AND D. TONG, *Normal functions concerning derivatives and shared sets*, Boletín da la Sociedad Matematica Mexicana **25** (2019), 589–596.
- [3] O. LEHTO, K. L. VIRTANEN, *Boundary behaviour and normal meromorphic functions*, Acta Math. **97** (1957), 47–65.
- [4] T. VAN TAN AND N. VAN THIN, *On Lappan's five-point theorem*, Comput. Methods Funct. Theory **17** (2017), 47–63.