

HARDY AND RELlich INEQUALITIES WITH REMAINDERS

W. DESMOND EVANS AND ROGER T. LEWIS

(communicated by A. Kufner)

Abstract. In this paper our primary concern is with the establishment of weighted Hardy inequalities in $L^p(\Omega)$ and Rellich inequalities in $L^2(\Omega)$ depending upon the distance to the boundary of domains $\Omega \subset \mathbb{R}^n$ with a finite diameter $D(\Omega)$. Improved constants are presented in most cases.

1. Introduction

Recently, considerable attention has been given to extensions of the multi-dimensional Hardy inequality of the form

$$\int_{\Omega} |\nabla u(\mathbf{x})|^2 d\mathbf{x} \geq \mu(\Omega) \int_{\Omega} \frac{|u(\mathbf{x})|^2}{\delta(\mathbf{x})^2} d\mathbf{x} + \lambda(\Omega) \int_{\Omega} |u(\mathbf{x})|^2 d\mathbf{x}, \quad u \in H_0^1(\Omega), \quad (1.1)$$

where Ω is an open connected subset of \mathbb{R}^n and

$$\delta(\mathbf{x}) := \text{dist}(\mathbf{x}, \partial\Omega).$$

It is known that for $\mu(\Omega) = \frac{1}{4}$ there are smooth domains for which $\lambda(\Omega) \leq 0$, and for $\lambda(\Omega) = 0$, there are smooth domains for which $\mu(\Omega) < \frac{1}{4}$ - see M. Marcus, V.J. Mizel, and Y. Pinchover [8] and T. Matskewich and P.E. Sobolevskii [9]. In [2], H. Brezis and M. Marcus showed that for domains of class C^2 inequality (1.1) holds for

$$\mu(\Omega) = \frac{1}{4} \quad \text{and some} \quad \lambda(\Omega) \in (-\infty, \infty)$$

and when Ω is convex

$$\lambda(\Omega) \geq \frac{1}{4D(\Omega)^2} \quad (1.2)$$

in which $D(\Omega)$ is the diameter of Ω .

M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and A. Laptev [6] answered a question posed by H. Brezis and M. Markus in [2] by establishing the improvement to (1.2) that (1.1) holds for a convex domain Ω , with

$$\mu(\Omega) = \frac{1}{4}, \quad \lambda(\Omega) \geq \frac{K(n)}{4|\Omega|^{\frac{2}{n}}}, \quad \text{and} \quad K(n) := n \left[\frac{s_{n-1}}{n} \right]^{2/n} \quad (1.3)$$

Mathematics subject classification (2000): 47A63, 46E35, 26D10.

Key words and phrases: Rellich inequality, Hardy inequality, Remainder terms.

in which $s_{n-1} := |\mathbb{S}^{n-1}|$ and $|\Omega|$ is the volume of Ω .

For a convex domain Ω and $\mu(\Omega) = 1/4$, a lower bound for $\lambda(\Omega)$ in (1.1) in terms of $|\Omega|$ was also obtained by S. Filippas, V. Maz'ya, and A. Tertikas in [5] as a special case of results on L^p Hardy inequalities. They prove that $\lambda(\Omega) \geq 3D_{int}(\Omega)^{-2}$, where $D_{int}(\Omega) = 2 \sup_{x \in \Omega} \delta(x)$, the internal diameter of Ω . Since $3D_{int}(\Omega)^{-2} \geq \frac{3}{4n}K(n)/|\Omega|^{2/n}$, their result is an improvement of (1.3) for $n = 2, 3$, but the estimates don't compare for $n > 3$.

In this paper we show that (1.1) holds for (1.3) replaced by

$$\mu(\Omega) = \frac{1}{4} \quad \text{and} \quad \lambda(\Omega) \geq \frac{3K(n)}{2|\Omega|^{\frac{2}{n}}}$$

as well as proving weighted versions of the Hardy inequality in $L^p(\Omega)$ for $p > 1$.

In the case $p = 2$, the following are special cases of our results. If Ω is convex and $\sigma \in (0, 1]$, then

$$\int_{\Omega} |\nabla u(\mathbf{x})|^2 d\mathbf{x} \geq \frac{2^\sigma n(1-\sigma)^2}{4D(\Omega)^\sigma} \int_{\Omega} \left\{ \frac{B(n, 2-\sigma)}{\delta(\mathbf{x})^{2-\sigma}} + 3 \left(\frac{s_{n-1}}{n|\Omega|} \right)^{\frac{2-\sigma}{n}} \right\} |u(\mathbf{x})|^2 d\mathbf{x} \quad (1.4)$$

for

$$B(n, p) := \frac{\Gamma(\frac{p+1}{2}) \cdot \Gamma(\frac{n}{2})}{\sqrt{\pi} \cdot \Gamma(\frac{n+p}{2})}. \quad (1.5)$$

If $\sigma \in [\frac{2-n}{2}, 0]$ and Ω is convex, then

$$\begin{aligned} \int_{\Omega} \delta(\mathbf{x})^\sigma |\nabla u(\mathbf{x})|^2 d\mathbf{x} &\geq \frac{n(1-\sigma)^2}{4} B(n, 2-\sigma) \int_{\Omega} \delta(\mathbf{x})^{\sigma-2} |u(\mathbf{x})|^2 d\mathbf{x} \\ &+ \frac{C_H(n, \sigma)}{|\Omega|^{\frac{2(1-\sigma)}{n}}} \int_{\Omega} \delta(\mathbf{x})^{|\sigma|} |u(\mathbf{x})|^2 d\mathbf{x}. \end{aligned}$$

for $C_H(n, \sigma)$ given in (3.4).

Similar results for weighted forms of the Hardy inequality in $L^p(\Omega)$ are given in section 4. Finally, we comment that our one-dimensional inequalities in §2 lead to improved constants for the Rellich inequality obtained by G. Barbatis in [1] for $n \geq 4$.

2. One-dimensional inequalities

As is the case in [6], our proofs are based on one-dimensional Hardy-type inequalities coupled with the use of the mean-distance function introduced by Davies to extend to higher dimensions; see [4]. The basic one-dimensional inequality is as follows:

LEMMA 1. *Let $u \in C_0^1(0, 2b)$, $\rho(t) := \min\{t, 2b - t\}$ and let $f \in C^1[0, b]$ be monotonic on $[0, b]$. Then for $p > 1$*

$$\int_0^{2b} |f'(\rho(t))||u(t)|^p dt \leq p^p \int_0^{2b} \frac{|f(\rho(t)) - f(b)|^p}{|f'(\rho(t))|^{p-1}} |u'(t)|^p dt. \quad (2.1)$$

Proof. First let $u := v\chi_{(0,b]}$, the restriction to $(0, b]$ of some $v \in C_0^1(0, 2b)$. For any constant c

$$-\int_0^b [f(t) - c]' |u(t)|^p dt = -[f(t) - c] |u(t)|^p \Big|_0^b + \int_0^b [f(t) - c] \frac{p}{2} [|u(t)|^2]^{\frac{p}{2}-1} [|u(t)|^2]' dt.$$

By choosing $c = f(b)$, we have that

$$-\int_0^b f'(t) |u(t)|^p dt = p \int_0^b [f(t) - f(b)] |u(t)|^{p-2} \Re \epsilon [\overline{u(t)} u'(t)] dt. \tag{2.2}$$

Similarly, for $u = v\chi_{(b,2b]}$, $v \in C_0^1(0, 2b)$, we have

$$-\int_b^{2b} f'(2b - s) |u(s)|^p ds = p \int_b^{2b} [f(2b - s) - f(b)] |u(s)|^{p-2} \Re \epsilon [\overline{u(s)} u'(s)] ds.$$

Therefore, since f is monotonic, for any $u \in C_0^1(0, 2b)$

$$\begin{aligned} \int_0^{2b} |f'(\rho(t))| |u(t)|^p dt &= p \int_0^{2b} [f(\rho(t)) - f(b)] |u(t)|^{p-2} \Re \epsilon [\overline{u(t)} u'(t)] dt \\ &\leq p \int_0^b |f'(\rho(t))|^{\frac{p-1}{p}} |u(t)|^{p-1} \frac{|f(\rho(t)) - f(b)|}{|f'(\rho(t))|^{\frac{p-1}{p}}} |u'(t)| dt \\ &\leq p \left[\int_0^b |f'(\rho(t))| |u(t)|^p dt \right]^{\frac{p-1}{p}} \left[\int_0^b \frac{|f(\rho(t)) - f(b)|^p}{|f'(\rho(t))|^{p-1}} |u'(t)|^p dt \right]^{\frac{1}{p}} \end{aligned}$$

on applying Hölder’s inequality. Inequality (2.1) now follows. □

The next lemma provides the one-dimensional result needed to improve (1.3), which was proved in [6].

LEMMA 2. *Let $\sigma \leq 1$ and define $\mu(t) := 2b - \rho(t)$. For all $u \in C_0^1(0, 2b)$*

$$\int_0^{2b} \rho(t)^\sigma |u'(t)|^2 dt \geq \left(\frac{1 - \sigma}{2} \right)^2 \int_0^{2b} \rho(t)^{\sigma-2} \left[1 + k(\sigma) \left(\frac{2\rho(t)}{\mu(t)} \right)^{1-\sigma} \right]^2 |u(t)|^2 dt, \tag{2.3}$$

for

$$k(\sigma) := \begin{cases} [1 - 2^{\frac{1}{\sigma}-1}]^{-\sigma}, & \sigma < 0, \\ 1, & \sigma \in [0, 1]. \end{cases}$$

Proof. On setting $f(t) = t^{\sigma-1}$ in (2.1) we get

$$|1 - \sigma|^p \int_0^{2b} \rho(t)^{\sigma-2} |u(t)|^p dt \leq p^p \int_0^{2b} \rho(t)^{p+\sigma-2} \left| 1 - \left[\frac{\rho(t)}{b} \right]^{1-\sigma} \right|^p |u'(t)|^p dt. \tag{2.4}$$

With $u \in C_0^1(0, 2b)$, let $p = 2$ and substitute $v(t) = [1 - (\frac{\rho(t)}{b})^{1-\sigma}]u(t)$ in (2.4). We claim that this gives

$$\int_0^{2b} \rho^\sigma(t) |v'(t)|^2 dt \geq \left(\frac{1-\sigma}{2}\right)^2 \int_0^{2b} \rho(t)^{\sigma-2} \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right]^{-2} |v(t)|^2 dt \quad (2.5)$$

for any real number σ . The substitution gives

$$\rho(t)^{\sigma/2} v'(t) = -(1-\sigma)b^{\sigma-1} \rho(t)^{-\sigma/2} \rho'(t)u(t) + \rho(t)^{\sigma/2} \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right] u'(t).$$

Consequently,

$$\begin{aligned} \rho(t)^\sigma |v'(t)|^2 &= (1-\sigma)^2 b^{2\sigma-2} \rho(t)^{-\sigma} |u(t)|^2 + \rho(t)^\sigma \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right]^2 |u'(t)|^2 \\ &\quad - (1-\sigma)b^{\sigma-1} \rho'(t) \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right] [|u|^2]' \end{aligned}$$

which implies that

$$\begin{aligned} \int_0^{2b} \rho(t)^\sigma |v'(t)|^2 dt &= \int_0^{2b} \rho(t)^\sigma \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right]^2 |u'(t)|^2 dt \\ &\quad + \int_0^{2b} (1-\sigma)^2 b^{2\sigma-2} \rho(t)^{-\sigma} |u(t)|^2 dt \\ &\quad + (1-\sigma)b^{\sigma-1} \int_0^{2b} \frac{d}{dt} \left[\rho'(t) \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right] \right] |u|^2 dt \\ &= \int_0^{2b} \rho(t)^\sigma \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right]^2 |u'(t)|^2 dt \end{aligned} \quad (2.6)$$

since $\rho'(t) = 1$ in $(0, b)$ and -1 in $(b, 2b)$. Therefore, (2.5) follows from (2.4).

Since $2b = \mu(t) + \rho(t)$

$$\begin{aligned} \left[1 - \left(\frac{\rho(t)}{b}\right)^{1-\sigma}\right]^{-2} &= \left[1 + \frac{\rho(t)^{1-\sigma}}{b^{1-\sigma} - \rho(t)^{1-\sigma}}\right]^2 \\ &= \left[1 + 2^{1-\sigma} \left(\frac{\rho(t)}{\mu(t)}\right)^{1-\sigma} k_\sigma \left(\frac{\rho(t)}{\mu(t)}\right)\right]^2 \end{aligned} \quad (2.7)$$

for

$$k_\sigma(x) := \frac{1}{(1+x)^{1-\sigma} - (2x)^{1-\sigma}}, \quad x \in [0, 1), \quad \sigma \neq 1.$$

For $\sigma < 1$, $k_\sigma(x) > 0$ in $(0, 1)$, $k_\sigma(0) = 1$ and $k_\sigma(x) \rightarrow \infty$ as $x \rightarrow 1^-$. By examining the derivative of $k_\sigma(x)$

$$k'_\sigma(x) = \frac{-(1 - \sigma)((1 + x)^{-\sigma} - 2^{1-\sigma}x^{-\sigma})}{[(1 + x)^{1-\sigma} - (2x)^{1-\sigma}]^2}$$

we see that

$$\lim_{x \rightarrow 0^+} k'_\sigma(x) = \begin{cases} -(1 - \sigma), & \sigma < 0, \\ 1, & \sigma = 0, \\ \infty, & 0 < \sigma < 1. \end{cases}$$

For $\sigma < 0$, $k_\sigma(x)$ is minimized at

$$x_\sigma := 1/(2^{1-\frac{1}{\sigma}} - 1) < 1.$$

Calculations show that

$$k_\sigma(x_\sigma) = [1 - 2^{\frac{1}{\sigma}-1}]^{-\sigma} =: k(\sigma).$$

For $\sigma \in [0, 1)$, $k'_\sigma(x)$ is never zero in $(0, 1)$ indicating that $k_\sigma(x)$ is minimized at $x = 0$ for $\sigma \in [0, 1)$ and $x \in [0, 1)$. The inequality (2.3) now follows. \square

In order to treat the case in which $p \neq 2$, we make use of the methods of Tidblom [11] and prove a weighted version of Theorem 1.1 in [11].

LEMMA 3. *Let $u \in C_0^1(0, 2b)$, $p \in (1, \infty)$, and $\sigma \leq p - 1$. Then*

$$\int_0^{2b} \rho(t)^\sigma |u'(t)|^p dt \geq \left[\frac{p - \sigma - 1}{p} \right]^p \int_0^{2b} \{ \rho(t)^{\sigma-p} + (p - 1)b^{\sigma-p} \} |u(t)|^p dt.$$

Proof. We may assume that $\sigma \neq p - 1$ since otherwise the conclusion is trivial. According to (2.2) for a monotonic function f and a positive function g ,

$$\begin{aligned} \int_0^b |f'(t)| |u(t)|^p dt &\leq \int_0^b p |f(t) - f(b)| |u(t)|^{p-1} |u'(t)| dt \\ &\leq p \left[\int_0^b g(t) |u'(t)|^p dt \right]^{1/p} \left[\int_0^b \left(\frac{|f(t) - f(b)|^p}{g(t)} \right)^{\frac{1}{p-1}} |u(t)|^p dt \right]^{1-\frac{1}{p}}. \end{aligned}$$

Consequently,

$$p^p \int_0^b g(t) |u'(t)|^p dt \geq \frac{\left(\int_0^b |f'(t)| |u(t)|^p dt \right)^p}{\left(\int_0^b \left(\frac{|f(t) - f(b)|^p}{g(t)} \right)^{1/(p-1)} |u(t)|^p dt \right)^{p-1}}.$$

Now, as in [11], using a corollary to Young's inequality, namely

$$A^p/B^{p-1} \geq pA - (p - 1)B,$$

with $A = \int_0^b |f'(t)| |u(t)|^p dt, B = \int_0^b \left(\frac{|f(t)-f(b)|^p}{g(t)} \right)^{1/p-1} |u(t)|^p dt$, it follows that

$$p^p \int_0^b g(t) |u'(t)|^p dt \geq \int_0^b \left\{ p|f'(t)| - (p-1) \left(\frac{|f(t)-f(b)|^p}{g(t)} \right)^{1/(p-1)} \right\} |u(t)|^p dt.$$

Choose $f(t) = t^{\sigma-p+1}$ and $g(t) = (p-\sigma-1)^{-(p-1)} t^\sigma$. Then

$$\begin{aligned} \left(\frac{|f(t)-f(b)|^p}{g(t)} \right)^{1/(p-1)} &= (p-\sigma-1) \left[\frac{|t^{\sigma-p+1}-b^{\sigma-p+1}|^p}{t^\sigma} \right]^{\frac{1}{p-1}} \\ &= (p-\sigma-1) t^{\sigma-p} \left[\left(1 - \left(\frac{t}{b} \right)^{p-\sigma-1} \right)^p \right]^{\frac{1}{p-1}}. \end{aligned}$$

Consequently, for $t \in (0, b)$

$$\begin{aligned} p|f'(t)| - (p-1) \left(\frac{|f(t)-f(b)|^p}{g(t)} \right)^{1/(p-1)} &= (p-\sigma-1) \left\{ p t^{\sigma-p} - (p-1) t^{\sigma-p} \left[\left(1 - \left(\frac{t}{b} \right)^{p-\sigma-1} \right)^p \right]^{\frac{1}{p-1}} \right\} \\ &= (p-\sigma-1) t^{\sigma-p} \left\{ 1 + (p-1) \left(1 - \left[1 - \left(\frac{t}{b} \right)^{p-\sigma-1} \right]^{\frac{p}{p-1}} \right) \right\} \\ &\geq (p-\sigma-1) t^{\sigma-p} \left\{ 1 + (p-1) \left(\frac{t}{b} \right)^{p-\sigma-1} \right\} \\ &\geq (p-\sigma-1) \left\{ t^{\sigma-p} + (p-1) \left(\frac{1}{b^{p-\sigma}} \right) \right\}. \end{aligned}$$

and the inequality follows. In the inequality above we have used the fact that

$$\left[1 - \left(\frac{t}{b} \right)^{p-\sigma-1} \right]^{\frac{p}{p-1}} \leq 1 - \left(\frac{t}{b} \right)^{p-\sigma-1}.$$

The proof is completed by following the last part of the proof of Lemma 1. □

For a certain range of values taken by $\sigma, \sigma \in [-c_\sigma, 1)$ with $c_\sigma > 0$, the inequality in $L^2(\Omega)$ given by Lemma 2 gives a better bound than Lemma 3 with $p = 2$. In fact for $\sigma < 1$

$$\rho(t)^{\sigma-2} \left[1 + k(\sigma) \left(\frac{2\rho(t)}{\mu(t)} \right)^{1-\sigma} \right]^2 = \rho(t)^{\sigma-2} + \frac{2^{2-\sigma} k(\sigma)}{\rho(t) \mu(t)^{1-\sigma}} + \frac{2^{2-2\sigma} k(\sigma)^2}{\rho(t) \sigma \mu(t)^{2-2\sigma}}$$

with

$$\begin{aligned} &\frac{2^{2-\sigma} k(\sigma)}{\rho(t) \mu(t)^{1-\sigma}} + \frac{2^{2-2\sigma} k(\sigma)^2}{\rho(t) \sigma \mu(t)^{2-2\sigma}} \\ &\geq \begin{cases} \frac{5}{2^\sigma} b^{\sigma-2}, & \sigma \in [0, 1), \\ [2 - \sigma + b^\sigma k(\sigma) \rho(t)^{|\sigma|}] k(\sigma) b^{\sigma-2}, & \sigma < 0. \end{cases} \end{aligned} \tag{2.8}$$

Since $k(\sigma)$ decreases to 0 as $|\sigma| \rightarrow \infty$ and $k(-3) \approx 0.22$, then the left-hand side of (2.8) is greater than $b^{\sigma-2}$ for $\sigma \in [-3, 1)$.

3. A Hardy inequality in $L^2(\Omega)$

We need the following notation (c.f.[6]). For each $\mathbf{x} \in \Omega$ and $v \in \mathbb{S}^{n-1}$,

$$\begin{aligned} \tau_v(\mathbf{x}) &:= \min\{s > 0 : \mathbf{x} + sv \notin \Omega\}; \\ D_v(\mathbf{x}) &:= \tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x}); \\ \rho_v(\mathbf{x}) &:= \min\{\tau_v(\mathbf{x}), \tau_{-v}(\mathbf{x})\}; \\ \mu_v(\mathbf{x}) &:= \max\{\tau_v(\mathbf{x}), \tau_{-v}(\mathbf{x})\} = D_v(\mathbf{x}) - \rho_v(\mathbf{x}); \\ D(\Omega) &:= \sup_{\mathbf{x} \in \Omega, v \in \mathbb{S}^{n-1}} D_v(\mathbf{x}); \\ \Omega_{\mathbf{x}} &:= \{\mathbf{y} \in \Omega : \mathbf{x} + t(\mathbf{y} - \mathbf{x}) \in \Omega, \forall t \in [0, 1]\}. \end{aligned}$$

Note that $D(\Omega)$ is the diameter of Ω and $\Omega_{\mathbf{x}}$ is the part of Ω which can be “seen” from the point $\mathbf{x} \in \Omega$. The volume of $\Omega_{\mathbf{x}}$ is denoted by $|\Omega_{\mathbf{x}}|$.

Let $d\omega(v)$ denote the normalized measure on \mathbb{S}^{n-1} (so that $1 = \int_{\mathbb{S}^{n-1}} d\omega(v)$) and define

$$\rho(\mathbf{x}; s) := \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^s d\omega(v). \tag{3.1}$$

Hence $\rho^{-1/2}(\mathbf{x}; -2) = \rho(\mathbf{x})$ the “mean-distance” function introduced by Davies in [4]. For

$$B(n, p) := \int_{\mathbb{S}^{n-1}} |\cos(\mathbf{e}, v)|^p d\omega(v) = \frac{\Gamma(\frac{p+1}{2}) \cdot \Gamma(\frac{n}{2})}{\sqrt{\pi} \cdot \Gamma(\frac{n+p}{2})}, \quad \mathbf{e} \in \mathbb{R}^n, \tag{3.2}$$

it is known that

$$\rho(\mathbf{x}; -p) := \int_{\mathbb{S}^{n-1}} \frac{1}{\rho_v(\mathbf{x})^p} d\omega(v) \geq \frac{B(n, p)}{\delta(\mathbf{x})^p} \tag{3.3}$$

for convex domains Ω — see Exercise 5.7 in [4], [3], and [11]. Note that $B(n, 2) = n^{-1}$. This fact can be applied to most of the results below when Ω is convex.

For a Hardy inequality in $L^2(\Omega)$ with weights we will need to define

$$C_H(n, \sigma) := n \left(\frac{\delta_{n-1}}{n} \right)^{\frac{2(1-\sigma)}{n}} k(\sigma) [2^{|\sigma|} + 2^{2|\sigma|-1} k(\sigma)] (1 - \sigma)^2 \tag{3.4}$$

for $\sigma \in [\frac{2-n}{2}, 0]$ and $n \geq 2$ where as given in Lemma 2

$$k(\sigma) := \begin{cases} [1 - 2^{\frac{1}{\sigma}-1}]^{-\sigma}, & \sigma < 0, \\ 1, & \sigma \in [0, 1]. \end{cases}$$

Note that $C_H(n, 0) = \frac{3}{2}K(n)$ for $K(n)$ defined in (1.3).

THEOREM 1. *If $\frac{2-n}{2} \leq \sigma \leq 0$, then for any $u \in C_0^1(\Omega)$*

$$\begin{aligned} \int_{\Omega} \delta(\mathbf{x})^\sigma |\nabla u(\mathbf{x})|^2 d\mathbf{x} &\geq \frac{n(1-\sigma)^2}{4} \int_{\Omega} \rho(\mathbf{x}; \sigma - 2) |u(\mathbf{x})|^2 d\mathbf{x} \\ &+ C_H(n, \sigma) \int_{\Omega} \frac{\delta(\mathbf{x})^{|\sigma|}}{|\Omega_{\mathbf{x}}|^{\frac{2(1-\sigma)}{n}}} |u(\mathbf{x})|^2 d\mathbf{x}. \end{aligned} \tag{3.5}$$

If $0 < \sigma \leq 1$, then

$$\int_{\Omega} |\nabla u(\mathbf{x})|^2 d\mathbf{x} \geq \frac{2^\sigma n(1-\sigma)^2}{4D(\Omega)^\sigma} \int_{\Omega} \left\{ \rho(\mathbf{x}; \sigma - 2) + 3 \left(\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right)^{\frac{2-\sigma}{n}} \right\} |u(\mathbf{x})|^2 d\mathbf{x}. \tag{3.6}$$

If Ω is convex, then for any $u \in C_0^1(\Omega)$

$$\begin{aligned} \int_{\Omega} \delta(\mathbf{x})^\sigma |\nabla u(\mathbf{x})|^2 d\mathbf{x} &\geq \frac{n(1-\sigma)^2}{4} B(n, 2-\sigma) \int_{\Omega} \delta(\mathbf{x})^{\sigma-2} |u(\mathbf{x})|^2 d\mathbf{x} \\ &\quad + \frac{C_H(n, \sigma)}{|\Omega|^{\frac{2(1-\sigma)}{n}}} \int_{\Omega} \delta(\mathbf{x})^{|\sigma|} |u(\mathbf{x})|^2 d\mathbf{x}. \end{aligned}$$

when $\sigma \in [\frac{2-n}{2}, 0]$ and

$$\int_{\Omega} |\nabla u(\mathbf{x})|^2 d\mathbf{x} \geq \frac{2^\sigma n(1-\sigma)^2}{4D(\Omega)^\sigma} \int_{\Omega} \left\{ B(n, 2-\sigma) \delta(\mathbf{x})^{\sigma-2} + 3 \left(\frac{s_{n-1}}{n|\Omega|} \right)^{\frac{2-\sigma}{n}} \right\} |u(\mathbf{x})|^2 d\mathbf{x}.$$

when $\sigma \in (0, 1]$.

Proof. Let $\partial_v u$, $v \in \mathbb{S}^{n-1}$, denote the derivative of u in the direction of v , i.e., $\partial_v u = v \cdot (\nabla u)$. It follows from Lemma 2 that for $\sigma \in (-\infty, 1]$

$$\int_{\Omega} \rho_v^\sigma(\mathbf{x}) |\partial_v u|^2 d\mathbf{x} \geq \left(\frac{1-\sigma}{2} \right)^2 \int_{\Omega} \rho_v(\mathbf{x})^{\sigma-2} \left(1 + k(\sigma) \left[\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right]^{(1-\sigma)} \right)^2 |u(\mathbf{x})|^2 d\mathbf{x}. \tag{3.7}$$

Expanding the integrand in (3.7), we have

$$\begin{aligned} &\rho_v(\mathbf{x})^{\sigma-2} \left(1 + k(\sigma) \left[\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right]^{(1-\sigma)} \right)^2 \\ &= \rho_v(\mathbf{x})^{\sigma-2} + 2^{2-\sigma} \frac{k(\sigma)\rho_v(\mathbf{x})^{-\sigma}}{(\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x}))^{1-\sigma}} + 2^{2(1-\sigma)} k(\sigma)^2 \frac{\rho_v(\mathbf{x})^{-\sigma}}{\mu_v(\mathbf{x})^{2(1-\sigma)}}. \end{aligned} \tag{3.8}$$

If $\sigma \leq 0$

$$\begin{aligned} &\rho_v(\mathbf{x})^{\sigma-2} \left[1 + k(\sigma) \left(\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right)^{(1-\sigma)} \right]^2 \\ &\geq \rho_v(\mathbf{x})^{\sigma-2} + 2^{2-\sigma} \frac{k(\sigma)\delta(\mathbf{x})^{|\sigma|}}{(\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x}))^{1-\sigma}} + 2^{2(1-\sigma)} k(\sigma)^2 \frac{\delta(\mathbf{x})^{|\sigma|}}{\tau_v(\mathbf{x})^{2(1-\sigma)} + \tau_{-v}(\mathbf{x})^{2(1-\sigma)}} \end{aligned} \tag{3.9}$$

since $\rho_v(\mathbf{x})^{-\sigma} \geq \delta(\mathbf{x})^{|\sigma|}$ in this case. As in [6], we note that since

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x}))^{1-\sigma} d\omega(v) &\leq \int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x}))^{2(1-\sigma)} d\omega(v) \\ &\leq \left[\int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x}))^n d\omega(v) \right]^{\frac{2(1-\sigma)}{n}} \\ &= \left[\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right]^{\frac{2(1-\sigma)}{n}} \end{aligned}$$

for $\sigma \geq \frac{2-n}{2}$, then

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} \frac{1}{(\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x}))^{1-\sigma}} d\omega(v) &\geq \left[\int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x}))^{1-\sigma} d\omega(v) \right]^{-1} \\ &\geq \left[\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right]^{-\frac{2(1-\sigma)}{n}}. \end{aligned}$$

For the third term in inequality (3.9)

$$\int_{\mathbb{S}^{n-1}} \left(\tau_v(\mathbf{x})^{2(1-\sigma)} + \tau_{-v}(\mathbf{x})^{2(1-\sigma)} \right) d\omega(v) = 2 \int_{\mathbb{S}^{n-1}} \tau_v(\mathbf{x})^{2(1-\sigma)} d\omega(v)$$

implying that for $\sigma \geq \frac{2-n}{2}$

$$\int_{\mathbb{S}^{n-1}} \left(\tau_v(\mathbf{x})^{2(1-\sigma)} + \tau_{-v}(\mathbf{x})^{2(1-\sigma)} \right)^{-1} d\omega(v) \geq \frac{1}{2} \left[\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right]^{-\frac{2(1-\sigma)}{n}}.$$

Consequently, for $\frac{2-n}{2} \leq \sigma \leq 0$ we have that

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^{\sigma-2} \left[1 + k(\sigma) \left(\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right)^{(1-\sigma)} \right]^2 d\omega(v) \\ \geq \rho(\mathbf{x}; \sigma - 2) + C_H(n, \sigma) \delta(\mathbf{x})^{|\sigma|} / \left[n |\Omega_{\mathbf{x}}|^{\frac{2(1-\sigma)}{n}} \right]. \end{aligned} \tag{3.10}$$

Upon combining this fact with (3.7) we have

$$\begin{aligned} \left(\frac{1-\sigma}{2} \right)^2 \int_{\Omega} \left\{ \rho(\mathbf{x}; \sigma - 2) + \frac{C_H(n, \sigma) \delta(\mathbf{x})^{|\sigma|}}{n |\Omega_{\mathbf{x}}|^{2(1-\sigma)/n}} \right\} |u(\mathbf{x})|^2 d\mathbf{x} \\ \leq \int_{\Omega} \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^{\sigma} |\partial_v u(\mathbf{x})|^2 d\omega(v) d\mathbf{x} \\ = \int_{\Omega} \delta(\mathbf{x})^{\sigma} \int_{\mathbb{S}^{n-1}} |\cos(v, \nabla u(\mathbf{x}))|^2 d\omega(v) |\nabla u(\mathbf{x})|^2 d\mathbf{x} \end{aligned} \tag{3.11}$$

for $\sigma \leq 0$. Since

$$\int_{\mathbb{S}^{n-1}} |\cos(v, \alpha)|^2 d\omega(v) = \frac{1}{n} \tag{3.12}$$

for any fixed $\alpha \in \mathbb{S}^{n-1}$ (see Tidblom [11], p.2270), inequality (3.5) follows.

For $0 < \sigma \leq 1$, we consider first the third term on the right-hand side of (3.8). We have

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^\sigma \mu_v(\mathbf{x})^{2(1-\sigma)} d\omega(v) \\ & \leq \int_{\mathbb{S}^{n-1}} 2^{-\sigma} (\tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x}))^\sigma (\tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x}))^{2(1-\sigma)} d\omega(v) \\ & = 2^{-\sigma} \|\tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x})\|_{L^{2-\sigma}(\mathbb{S}^{n-1})}^{2-\sigma} \\ & \leq 2^{-\sigma} \left[\|\tau_v(\mathbf{x})\|_{L^{2-\sigma}(\mathbb{S}^{n-1})} + \|\tau_{-v}(\mathbf{x})\|_{L^{2-\sigma}(\mathbb{S}^{n-1})} \right]^{2-\sigma} \\ & = 2^{2(1-\sigma)} \int_{\mathbb{S}^{n-1}} \tau_v(\mathbf{x})^{2-\sigma} d\omega(v) \\ & \leq 2^{2(1-\sigma)} \left[\int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x}))^n d\omega(v) \right]^{\frac{2-\sigma}{n}} \\ & = 2^{2(1-\sigma)} \left[\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right]^{\frac{2-\sigma}{n}} \end{aligned}$$

for $n \geq 2$ by the Minkowski and Hölder inequalities. Therefore, the term

$$\int_{\mathbb{S}^{n-1}} \frac{\rho_v(\mathbf{x})^{-\sigma} d\omega(v)}{\mu_v(\mathbf{x})^{2(1-\sigma)}} \geq 2^{2(\sigma-1)} \left(\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right)^{\frac{2-\sigma}{n}}.$$

Similarly, in the second term of (3.8)

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x}) \mu_v(\mathbf{x})^{1-\sigma} d\omega(v) & \leq \frac{1}{2} \int_{\mathbb{S}^{n-1}} (\tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x})) (\tau_v(\mathbf{x}) + \tau_{-v}(\mathbf{x}))^{1-\sigma} d\omega(v) \\ & \leq 2^{1-\sigma} \left[\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right]^{\frac{2-\sigma}{n}} \end{aligned}$$

as before implying that

$$\int_{\mathbb{S}^{n-1}} \frac{d\omega(v)}{\rho_v(\mathbf{x}) \mu_v(\mathbf{x})^{1-\sigma}} \geq 2^{\sigma-1} \left(\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right)^{\frac{2-\sigma}{n}}.$$

For $0 < \sigma < 1$ we now have that

$$\int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^{\sigma-2} \left[1 + k(\sigma) \left(\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right)^{(1-\sigma)} \right]^2 d\omega(v) \geq \rho(\mathbf{x}; \sigma - 2) + 3 \left(\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right)^{\frac{2-\sigma}{n}}$$

since $k(\sigma) = 1$ in this case. Consequently,

$$\begin{aligned} & \int_{\Omega} \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^\sigma |\cos(v, \nabla u(\mathbf{x}))|^2 d\omega(v) |\nabla u(\mathbf{x})|^2 d\mathbf{x} \\ & \geq \left(\frac{1-\sigma}{2} \right)^2 \int_{\Omega} \left[\rho(\mathbf{x}; \sigma - 2) + 3 \left(\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right)^{\frac{2-\sigma}{n}} \right] |u(\mathbf{x})|^2 d\mathbf{x}. \end{aligned}$$

According to (3.12) it follows that

$$\int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^\sigma |\cos(v, \nabla u(\mathbf{x}))|^2 d\omega(v) \leq \frac{D(\Omega)^\sigma}{2^\sigma n}.$$

Therefore, (3.6) holds.

The inequalities in the statement of the theorem for the case of a convex domain Ω follow from (3.3) and the fact that $|\Omega_{\mathbf{x}}| = |\Omega|$ for all $\mathbf{x} \in \Omega$. □

4. An $L^p(\Omega)$ inequality

With the guidance of Tidblom’s analysis for the Hardy inequality in [11], L^p versions of the weighted Hardy theorem in the last section can be proved by similar techniques. When $\sigma = 0$, the next theorem reduces to Theorem 2.1 of [11].

THEOREM 2. *Let $u \in C_0^1(\Omega)$ and $p \in (1, \infty)$. If $\sigma \leq 0$, then for $B(n, p)$ defined in (3.2)*

$$\int_{\Omega} \delta(\mathbf{x})^\sigma |\nabla u(\mathbf{x})|^p d\mathbf{x} \geq \frac{[|p - \sigma - 1|/p]^p}{B(n, p)} \int_{\Omega} \left\{ \rho(\mathbf{x}; \sigma - p) + (p - 1) \left[\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right]^{\frac{p-\sigma}{n}} \right\} |u(\mathbf{x})|^p d\mathbf{x} \tag{4.1}$$

and if $\sigma \in [0, p - 1]$, then

$$\int_{\Omega} |\nabla u(\mathbf{x})|^p d\mathbf{x} \geq \frac{2^\sigma [|p - \sigma - 1|/p]^p}{B(n, p) D(\Omega)^\sigma} \int_{\Omega} \left\{ \rho(\mathbf{x}; \sigma - p) + (p - 1) \left[\frac{s_{n-1}}{n|\Omega_{\mathbf{x}}|} \right]^{\frac{p-\sigma}{n}} \right\} |u(\mathbf{x})|^p d\mathbf{x}. \tag{4.2}$$

If Ω is convex, $\rho(\mathbf{x}, \sigma - p)$ can be replaced in (4.1) and (4.2) by the term $B(n, p - \sigma)/\delta(\mathbf{x})^{p-\sigma}$ (in view of (3.3)) and $|\Omega_{\mathbf{x}}|$ by $|\Omega|$.

Proof. From Lemma 3 we have that for $\sigma \leq p - 1$, any $v \in \mathbb{S}^{n-1}$, and $u \in C_0^1(\Omega)$

$$\int_{\Omega} \rho_v(\mathbf{x})^\sigma |\partial_v u(\mathbf{x})|^p d\mathbf{x} \geq \left[\frac{|p - \sigma - 1|}{p} \right]^p \int_{\Omega} \left\{ \rho_v(\mathbf{x})^{\sigma-p} + \frac{(p - 1)2^{p-\sigma}}{D_v(\mathbf{x})^{p-\sigma}} \right\} |u(\mathbf{x})|^p d\mathbf{x}. \tag{4.3}$$

If $\sigma \leq 0$ we bound $\rho_v(\mathbf{x})^\sigma$ for any $v \in \mathbb{S}^{n-1}$ by $\delta(\mathbf{x})^\sigma$ in the first integral above. If $\sigma > 0$, we bound it by $D(\Omega)^\sigma/2^\sigma$. As in [11] we may use the fact that

$$\int_{\mathbb{S}^{n-1}} |\partial_v u(\mathbf{x})|^p d\omega(v) = B(n, p) |\nabla u(\mathbf{x})|^p. \tag{4.4}$$

After bounding $\rho_v(\mathbf{x})^\sigma$ as described above, integrate in (4.3) over \mathbb{S}^{n-1} with respect to $d\omega(v)$. In order to evaluate the integral of $(2/D_v(\mathbf{x}))^{p-\sigma}$, we proceed as in [11]. Since $\sigma \leq p - 1$, then $f(t) = t^{\sigma-p}$ is convex for $t > 0$ and we have that

$$\int_{\mathbb{S}^{n-1}} \left(\frac{2}{D_v(\mathbf{x})}\right)^{p-\sigma} d\omega(v) \geq \left(\int_{\mathbb{S}^{n-1}} \frac{D_v(\mathbf{x})}{2} d\omega(v)\right)^{\sigma-p} \geq \left(\frac{n|\Omega_{\mathbf{x}}|}{s_{n-1}}\right)^{-\frac{p-\sigma}{n}} \tag{4.5}$$

by Jensen’s inequality and Lemma 2.1 of [11]. The conclusion follows. \square

5. Rellich’s inequality

The methods described above with Proposition 1 below can be used to prove a weighted Rellich inequality which, for $n \geq 4$ and without weights, improves the constant given in a Rellich inequality proved recently by Barbatis ([1], Theorem 1.2). A comparison is made below. The methods used by Barbatis depends upon the identity (5.2) first proved by M.P. Owen ([10], see the proof of Theorem 2.3). In order to incorporate weights, our proof requires the point-wise identity (5.1) which does not follow from the proof of Owen.

PROPOSITION 1. *Let Ω be a domain in \mathbb{R}^n . Then, for all $u \in C^2(\mathbb{R}^n)$*

$$\int_{\mathbb{S}^{n-1}} |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) = \frac{1}{n(n+2)} \left[|\Delta u(\mathbf{x})|^2 + 2 \sum_{i,j=1}^n \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 \right], \tag{5.1}$$

and for all $u \in C_0^2(\Omega)$

$$\int_{\Omega} \int_{\mathbb{S}^{n-1}} |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) d\mathbf{x} = \frac{3}{n(n+2)} \int_{\Omega} |\Delta u(\mathbf{x})|^2 d\mathbf{x}. \tag{5.2}$$

Proof.

For $v = (v_1, \dots, v_n)$ we have

$$\begin{aligned} \partial_v^2 u &= (v \cdot \nabla)^2 u = \sum_{\ell,m=1}^n v_\ell v_m u_{\ell m} \\ &= \sum_{\ell=1}^n v_\ell^2 u_{\ell\ell} + 2 \sum_{1 \leq \ell < m \leq n} v_\ell v_m u_{\ell m} \end{aligned}$$

in which $u_{pq}(\mathbf{x}) := \frac{\partial^2 u(\mathbf{x})}{\partial x_p \partial x_q}$. Consequently,

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} |\partial_v^2 u|^2 d\omega(v) &= \sum_{\ell,m=1}^n u_{\ell\ell} \overline{u_{mm}} \int_{\mathbb{S}^{n-1}} (v_\ell)^2 (v_m)^2 d\omega(v) \\ &+ 4 \sum_{m=1}^n \sum_{1 \leq p < q \leq n} \Re(u_{mm} \overline{u_{pq}}) \int_{\mathbb{S}^{n-1}} (v_m)^2 v_p v_q d\omega(v) \\ &+ 4 \sum_{1 \leq j < k \leq n} \sum_{1 \leq p < q \leq n} \Re(u_{pq} \overline{u_{jk}}) \int_{\mathbb{S}^{n-1}} v_p v_q v_j v_k d\omega(v). \end{aligned} \tag{5.3}$$

Let $\theta_j \in [0, \pi]$ for $j = 1, \dots, n - 2$, and $\theta_{n-1} \in [0, 2\pi]$. Using the convention that $\prod_{j=q}^p = 1$ for $p < q$ and $\theta_n = 0$, we have

$$v_j = \prod_{k=1}^{j-1} \sin \theta_k \cos \theta_j, \quad j = 1, \dots, n,$$

$$d\omega(v) := \frac{(n-2)!!}{\gamma_n} \prod_{k=1}^{n-2} (\sin \theta_k)^{n-1-k} d\theta_k d\theta_{n-1}, \quad (5.4)$$

for $n!! := n \cdot (n - 2) \cdot (n - 4) \cdots 1$ and

$$\gamma_n = \begin{cases} 2(2\pi)^{(n-1)/2} & \text{for } n \text{ odd,} \\ (2\pi)^{n/2} & \text{for } n \text{ even.} \end{cases}$$

Calculations show that

$$\int_{\mathbb{S}^{n-1}} (v_m)^2 v_p v_q d\omega(v) = 0, \quad m = 1, \dots, n, \quad 1 \leq p < q \leq n$$

implying that the second term on the right-hand side of (5.3) vanishes.

A similar consideration for the third term on the right-hand side of (5.3) shows that

$$\int_{\mathbb{S}^{n-1}} v_p v_q v_j v_k d\omega(v) \neq 0, \quad 1 \leq p < q \leq n, \quad 1 \leq j < k \leq n,$$

only if $j = p$ and $k = q$. Therefore, (5.3) reduces to

$$\int_{\mathbb{S}^{n-1}} |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) = \sum_{\ell, m=1}^n u_{\ell\ell} \overline{u_{mm}} \int_{\mathbb{S}^{n-1}} (v_\ell)^2 (v_m)^2 d\omega(v) + 4 \sum_{1 \leq p < q \leq n} |u_{pq}|^2 \int_{\mathbb{S}^{n-1}} (v_p)^2 (v_q)^2 d\omega(v). \quad (5.5)$$

However, further calculations show that

$$\int_{\mathbb{S}^{n-1}} v_p^2 v_q^2 d\omega(v) = \begin{cases} \frac{1}{n(n+2)} & 1 \leq p < q \leq n, \\ \frac{3}{n(n+2)} & p = q = 1, \dots, n \end{cases} \quad (5.6)$$

implying that

$$\int_{\mathbb{S}^{n-1}} |\partial_v^2 u|^2 d\omega(v) = \frac{3}{n(n+2)} \sum_{m=1}^n |u_{mm}|^2 + \frac{1}{n(n+2)} \sum_{1 \leq p < q \leq n} [4|u_{pq}|^2 + 2\Re(u_{pp} \overline{u_{qq}})] = \frac{1}{n(n+2)} \left[|\Delta u(\mathbf{x})|^2 + 2 \sum_{i,j=1}^n \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 \right] \quad (5.7)$$

which is (5.1). Equality (5.2) now follows since

$$\sum_{i,j=1}^n \int_{\Omega} \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 d\mathbf{x} = \int_{\Omega} |\Delta u(\mathbf{x})|^2 d\mathbf{x}.$$

□

Define

$$d(\mathbf{x}; \sigma) := \begin{cases} \delta(\mathbf{x})^\sigma, & \sigma < 0, \\ \left(\frac{\rho(\Omega)}{2}\right)^\sigma, & \sigma \in [0, 1]; \end{cases}$$

$$\beta(n, \sigma) := \frac{(1 - \sigma)^2(3 - \sigma)^2 n(n + 2)}{16};$$

and

$$C_R(n, \sigma) := 2^{4-\sigma} k(\sigma - 2) \left[\frac{s_{n-1}}{n} \right]^{\frac{4-\sigma}{n}} (1 + 2^{2-\sigma} k(\sigma - 2)) \quad (5.8)$$

for $\sigma \leq 1$ and $k(\sigma)$ defined in Lemma 2.

THEOREM 3. For $\sigma \leq 1$ and $u \in C_0^2(\Omega)$,

$$\begin{aligned} \int_{\Omega} d(\mathbf{x}; \sigma) \left[|\Delta u(\mathbf{x})|^2 + 2 \sum_{i,j=1}^n \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 \right] d\mathbf{x} \\ \geq \beta(n, \sigma) \left\{ \int_{\Omega} \rho(\mathbf{x}; \sigma - 4) |u(\mathbf{x})|^2 d\mathbf{x} \right. \\ \left. + 2^{4-\sigma} k(\sigma - 2) \left[\frac{s_{n-1}}{n} \right]^{\frac{4-\sigma}{n}} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4-\sigma}{n}}} d\mathbf{x} \right\} \end{aligned} \quad (5.9)$$

holds when $n \geq 4 - \sigma$ and

$$\begin{aligned} \int_{\Omega} d(\mathbf{x}; \sigma) \left[|\Delta u(\mathbf{x})|^2 + 2 \sum_{i,j=1}^n \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 \right] d\mathbf{x} \\ \geq \beta(n, \sigma) \left\{ \int_{\Omega} \rho(\mathbf{x}; \sigma - 4) |u(\mathbf{x})|^2 d\mathbf{x} \right. \\ + 2^{4-\sigma} k(\sigma - 2) \left[\frac{s_{n-1}}{n} \right]^{\frac{4-\sigma}{n}} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4-\sigma}{n}}} d\mathbf{x} \\ \left. + 2^{2(3-\sigma)} k(\sigma - 2)^2 \left[\frac{s_{n-1}}{n} \right]^{\frac{4+t-\sigma}{n}} \int_{\Omega} \frac{\delta(\mathbf{x})^t |u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4+t-\sigma}{n}}} d\mathbf{x} \right\} \end{aligned} \quad (5.10)$$

holds when $n \geq 4 + t - \sigma$ and $t \geq 2 - \sigma$.

Proof. For $\sigma \leq 1$, it follows that

$$\begin{aligned} \int_0^{2b} \rho(t)^\sigma |u''(t)|^2 dt &\geq \int_0^{2b} \rho(t)^\sigma \left[1 - \left(\frac{\rho(t)}{\mu(t)} \right)^{1-\sigma} \right]^2 |u''(t)|^2 dt \\ &\geq \left(\frac{1-\sigma}{2} \right)^2 \int_0^{2b} \rho(t)^{\sigma-2} |u'(t)|^2 dt \end{aligned}$$

by (2.4). Therefore, for $\sigma \leq 1$ and $u \in C_0^2(0, 2b)$,

$$\begin{aligned} & \int_0^{2b} \rho(t)^\sigma |u''(t)|^2 dt \\ & \geq \left(\frac{(1-\sigma)(3-\sigma)}{4} \right)^2 \int_0^{2b} \rho(t)^{\sigma-4} \left[1 + k(\sigma-2) \left(\frac{2\rho(t)}{\mu(t)} \right)^{3-\sigma} \right]^2 |u(t)|^2 dt \end{aligned} \tag{5.11}$$

by (2.3).

From (5.11) we have for $u \in C_0^2(\Omega)$

$$\begin{aligned} & \int_\Omega \rho_v(\mathbf{x})^\sigma |\partial_v^2 u(\mathbf{x})|^2 d\mathbf{x} \\ & \geq \left(\frac{(1-\sigma)(3-\sigma)}{4} \right)^2 \int_\Omega \rho_v(\mathbf{x})^{\sigma-4} \left\{ 1 + k(\sigma-2) \left(\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right)^{3-\sigma} \right\}^2 |u(\mathbf{x})|^2 d\mathbf{x} \end{aligned} \tag{5.12}$$

for $\sigma \leq 1$. As in (3.8) we write

$$\begin{aligned} & \rho_v(\mathbf{x})^{\sigma-4} \left\{ 1 + k(\sigma-2) \left(\frac{2\rho_v(\mathbf{x})}{\mu_v(\mathbf{x})} \right)^{3-\sigma} \right\}^2 \\ & = \rho_v(\mathbf{x})^{\sigma-4} + 2^{4-\sigma} k(\sigma-2) \frac{\rho_v^{-1}(\mathbf{x})}{\mu_v(\mathbf{x})^{3-\sigma}} + 2^{2(3-\sigma)} k(\sigma-2)^2 \frac{\rho_v^{-\sigma+2}(\mathbf{x})}{\mu_v(\mathbf{x})^{2(3-\sigma)}}. \end{aligned} \tag{5.13}$$

Since $\rho_v(\mathbf{x})\mu_v(\mathbf{x}) = \tau_v(\mathbf{x})\tau_{-v}(\mathbf{x})$, in the second term on the right-hand side of (5.13) we may write

$$\frac{\rho_v^{-1}(\mathbf{x})}{\mu_v(\mathbf{x})^{3-\sigma}} = \frac{1}{[\tau_v(\mathbf{x})\tau_{-v}(\mathbf{x})]\mu_v(\mathbf{x})^{2-\sigma}} =: I(v; \mathbf{x}).$$

Thus

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} I(v; \mathbf{x}) d\omega(v) &= \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{\sigma-3}(\mathbf{x}) \tau_{-v}(\mathbf{x})^{-1} d\omega(v) \\ & \quad + \int_{\tau_v(\mathbf{x}) \leq \tau_{-v}(\mathbf{x})} \tau_{-v}(\mathbf{x})^{\sigma-3}(\mathbf{x}) \tau_v(\mathbf{x})^{-1} d\omega(v) \\ & \geq \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{\sigma-4}(\mathbf{x}) d\omega(v) \\ & \quad + \int_{\tau_v(\mathbf{x}) \leq \tau_{-v}(\mathbf{x})} \tau_{-v}(\mathbf{x})^{\sigma-4}(\mathbf{x}) d\omega(v) \end{aligned}$$

and

$$\begin{aligned} \left\{ \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{\sigma-4} d\omega(v) \right\}^{-1} &\leq \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{-\sigma+4} d\omega(v) \\ &\leq \left\{ \int_{\mathbb{S}^{n-1}} \tau_v^n(\mathbf{x}) d\omega(v) \right\}^{(4-\sigma)/n} \\ &= \left(\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right)^{(4-\sigma)/n} \end{aligned}$$

for $n \geq 4 - \sigma$. Therefore for the second term on the right-hand side of (5.13), for $\sigma \leq 1$ and $n \geq 4 - \sigma$, it follows that

$$\int_{\Omega} \int_{\mathbb{S}^{n-1}} \frac{\rho_v^{-1}(\mathbf{x}) d\omega(v)}{\mu_v(\mathbf{x})^{3-\sigma}} |u(\mathbf{x})|^2 d\mathbf{x} \geq \left(\frac{s_{n-1}}{n} \right)^{\frac{4-\sigma}{n}} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4-\sigma}{n}}} d\mathbf{x}. \tag{5.14}$$

For any $t \in (-\infty, \infty)$, we may write the third term in (5.13) as

$$\frac{\rho_v^{-\sigma+2}(\mathbf{x})}{\mu_v(\mathbf{x})^{2(3-\sigma)}} = \rho_v(\mathbf{x})^t (\tau_v(\mathbf{x}) \tau_{-v}(\mathbf{x}))^{2-\sigma-t} \mu(\mathbf{x})^{-8+3\sigma+t} =: \rho_v(\mathbf{x})^t J(v, \mathbf{x}).$$

If $t \geq 2 - \sigma$

$$\int_{\mathbb{S}^{n-1}} J(v; \mathbf{x}) d\omega(v) \geq \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{-4+\sigma-t} d\omega(v) + \int_{\tau_v(\mathbf{x}) \leq \tau_{-v}(\mathbf{x})} \tau_{-v}(\mathbf{x})^{-4+\sigma-t} d\omega(v).$$

As before

$$\begin{aligned} \left\{ \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{-4+\sigma-t} d\omega(v) \right\}^{-1} &\leq \int_{\tau_v(\mathbf{x}) \geq \tau_{-v}(\mathbf{x})} \tau_v(\mathbf{x})^{4-\sigma+t} d\omega(v) \\ &\leq \left\{ \int_{\mathbb{S}^{n-1}} \tau_v^n(\mathbf{x}) d\omega(v) \right\}^{(4-\sigma+t)/n} \\ &= \left(\frac{n}{s_{n-1}} |\Omega_{\mathbf{x}}| \right)^{(4-\sigma+t)/n} \end{aligned}$$

if $n \geq 4 - \sigma + t$. Associated with the third term on the right-hand side of (5.13), we have for $\sigma \leq 1$, $t \geq 2 - \sigma > 0$, and $n \geq 4 - \sigma + t$

$$\int_{\Omega} \int_{\mathbb{S}^{n-1}} \frac{\rho_v^{-\sigma+2}(\mathbf{x}) d\omega(v)}{\mu_v(\mathbf{x})^{2(3-\sigma)}} |u(\mathbf{x})|^2 d\mathbf{x} \geq \left(\frac{s_{n-1}}{n} \right)^{\frac{4+t-\sigma}{n}} \int_{\Omega} \frac{\delta(\mathbf{x})^t |u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4+t-\sigma}{n}}} d\mathbf{x}. \tag{5.15}$$

From (5.12) – (5.15) we obtain

$$\begin{aligned} \int_{\Omega} \int_{\mathbb{S}^{n-1}} \rho_v(\mathbf{x})^\sigma |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) d\mathbf{x} \geq \\ \frac{(1-\sigma)^2(3-\sigma)^2}{16} \left\{ \int_{\Omega} \rho(\mathbf{x}; \sigma-4) |u(\mathbf{x})|^2 d\mathbf{x} + 2^{4-\sigma} k(\sigma-2) \left[\frac{s_{n-1}}{n} \right]^{\frac{4-\sigma}{n}} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4-\sigma}{n}}} d\mathbf{x} \right. \\ \left. + 2^{2(3-\sigma)} k(\sigma-2)^2 \left[\frac{s_{n-1}}{n} \right]^{\frac{4+t-\sigma}{n}} \int_{\Omega} \frac{\delta(\mathbf{x})^t |u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4+t-\sigma}{n}}} d\mathbf{x} \right\} \end{aligned}$$

provided $\sigma \leq 1$, $t \geq 2 - \sigma$, and $n \geq 4 + t - \sigma$.

Note, that we may simply choose zero as a lower bound for the third term on the right-hand side of (5.13) and conclude that

$$\int_{\Omega} \int_{S^{n-1}} \rho_v(\mathbf{x})^\sigma |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) d\mathbf{x} \geq \frac{(1-\sigma)^2(3-\sigma)^2}{16} \left\{ \int_{\Omega} \rho(\mathbf{x}; \sigma-4) |u(\mathbf{x})|^2 d\mathbf{x} + 2^{4-\sigma} k(\sigma-2) \left[\frac{s_{n-1}}{n} \right]^{\frac{4-\sigma}{n}} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{|\Omega_{\mathbf{x}}|^{\frac{4-\sigma}{n}}} d\mathbf{x} \right\}$$

for $\sigma \leq 1$ and $n \geq 4 - \sigma$.

Now, it follows from Proposition 1 that

$$\int_{\Omega} \int_{S^{n-1}} \rho_v(\mathbf{x})^\sigma |\partial_v^2 u(\mathbf{x})|^2 d\omega(v) d\mathbf{x} \leq \frac{1}{n(n+2)} \int_{\Omega} d(\mathbf{x}; \sigma) \left[|\Delta u(\mathbf{x})|^2 + 2 \sum_{i,j=1}^n \left| \frac{\partial^2 u(\mathbf{x})}{\partial x_i \partial x_j} \right|^2 \right] d\mathbf{x}.$$

Thus, (5.9) and (5.10) are proved. □

It follows from Theorem 1.2 of Barbatis [1] that for a convex bounded domain Ω and all $u \in C_0^\infty(\Omega)$

$$\int_{\Omega} |\Delta u(\mathbf{x})|^2 d\mathbf{x} \geq \frac{9}{16} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{\delta(\mathbf{x})^4} d\mathbf{x} + \frac{11}{48} n(n+2) \left[\frac{s_{n-1}}{n|\Omega|} \right]^{4/n} \int_{\Omega} |u(\mathbf{x})|^2 d\mathbf{x}. \tag{5.16}$$

As in Theorem 2, for a convex domain $\Omega \subset \mathbb{R}^n$, we may replace $\rho(\mathbf{x}, \sigma - 4)$ in Theorem 3 by $B(n, 4 - \sigma)/\delta(\mathbf{x})^{4-\sigma}$ and $|\Omega_{\mathbf{x}}|$ by $|\Omega|$ to conclude from (5.9) that for $n \geq 4$

$$\int_{\Omega} |\Delta u(\mathbf{x})|^2 d\mathbf{x} \geq \frac{9}{16} \int_{\Omega} \frac{|u(\mathbf{x})|^2}{\delta(\mathbf{x})^4} d\mathbf{x} + c_4 n(n+2) \left[\frac{s_{n-1}}{n|\Omega|} \right]^{4/n} \int_{\Omega} |u(\mathbf{x})|^2 d\mathbf{x} \tag{5.17}$$

for all $u \in C_0^\infty(\Omega)$ in which $c_4 = 3k(-2) \approx 1.25$. Therefore (5.17) improves the bound given by (5.16) for all $n \geq 4$.

REFERENCES

[1] G. BARBATIS, *Improved Rellich inequalities for the polyharmonic operator*, Indiana University Mathematics Journal **55**(4) (2006), 1401-1422.
 [2] H. BREZIS AND M. MARCUS, *Hardy's inequalities revisited*, Dedicated to Ennio De Giorgi, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1997, 1998), 217-237.
 [3] E.B. DAVIES, *A Review of Hardy Inequalities*. The Maz'ya Anniversary Collection, Vol. 2, Oper. Theory Adv. Appl., Vol. 110, pp. 55-67, Birkäuser, Basel, 1999.
 [4] E.B. DAVIES, *Spectral Theory and Differential Operators*, Cambridge Studies in Advanced Mathematics, Vol. 42, Cambridge Univ. Press, Cambridge, 1995.
 [5] S. FILIPPAS, V. MAZ'YA, AND A. TERTIKAS, *On a question of Brezis and Marcus*, Calc. Var. Partial Differential Equations **25**(4) (2006), 491-501.
 [6] M. HOFFMANN-OSTENHOF, T. HOFFMANN-OSTENHOF AND A. LAPTEV, *A geometrical version of Hardy's inequality*, J. Funct. Anal., **189** (2002), 539-548.
 [7] E.H. LIEB AND M. LOSS, *Analysis*, Graduate Studies in Mathematics, vol. 14, 2nd edition, American Mathematical Society, Providence, R.I., 2001.

- [8] M. MARCUS, V.J. MIZEL, AND Y. PINCHOVER, *On the best constant for Hardy's inequality in \mathbb{R}^n* , Trans. Amer. Math. Soc. **350** (1998), 3237–3255.
- [9] T. MATSKEWICH AND P.E. SOBOLEVSKII, *The best possible constant in a generalized Hardy's inequality for convex domains in \mathbb{R}^n* , Nonlinear Analysis TMA, **28** (1997), 1601–1610.
- [10] M.P. OWEN, *The Hardy-Rellich inequality for polyharmonic operators*, Proc. Royal Society of Edinburgh, A **129** (1999), 825–839.
- [11] J. TIDBLÖM, *A geometrical version of Hardy's inequality for $W_0^{1,p}(\Omega)$* , Proc. A.M.S., **132**(8) (2004), 2265–2271.

(Received March 28, 2007)

W.D.Evans
School of Mathematics
Cardiff University
Senghennydd Road
Cardiff CF24 4AG
UK

e-mail: EvansWD@cardiff.ac.uk

R.T.Lewis
Department of Mathematics
University of Alabama at Birmingham
Birmingham
AL 35294-1170
USA

e-mail: lewis@math.uab.edu