

THE GOLDEN–THOMPSON–SEGAL TYPE INEQUALITIES RELATED TO THE WEIGHTED GEOMETRIC MEAN DUE TO LAWSON–LIM

JUN ICHI FUJII, MASATOSHI FUJII AND YUKI SEO

Abstract. In this paper, by using the weighted geometric mean $G[n,t]$ and the weighted arithmetic one $A[n,t]$ due to Lawson-Lim for each $t \in [0, 1]$, we investigate n -variable versions of a complement of the Golden-Thompson-Segal type inequality due to Ando-Hiai: Let H_1, H_2, \dots, H_n be selfadjoint operators such that $m \leq H_i \leq M$ for $i = 1, 2, \dots, n$ and some scalars $m \leq M$. Then

$$\begin{aligned} S(e^{p(M-m)})^{-\frac{2}{p}} & \| G[n,t](e^{pH_1}, \dots, e^{pH_n})^{\frac{1}{p}} \| \\ & \leq \| e^{A[n,t](H_1, \dots, H_n)} \| \leq S(e^{p(M-m)})^{\frac{2}{p}} \| G[n,t](e^{pH_1}, \dots, e^{pH_n})^{\frac{1}{p}} \| \end{aligned}$$

for all $p > 0$ and the both-hand sides of the inequality above converge to the middle-hand side as $p \downarrow 0$, where $S(\cdot)$ is the Specht ratio and $\|\cdot\|$ stands for the operator norm.

Mathematics subject classification (2000): 47A30, 47A63 and 47A64.

Keywords and phrases: Positive operator, Specht ratio, Golden-Thompson-Segal inequality, geometric mean, Lie-Trotter formula.

REFERENCES

- [1] M. ALIĆ, P. S. BULLEN, J. E. PEČARIĆ AND V. VOLNEC, *On the geometric-arithmetic mean inequality for matrices*, Math. Communications, **2** (1997), 125–128.
- [2] T. ANDO AND F. HIAI, *Log-majorization and complementary Golden-Thompson type inequalities*, Linear Alg. Appl., **197/198** (1994), 113–131.
- [3] T. ANDO, C.-K. LI AND R. MATHIAS, *Geometric means*, Linear Alg. Appl., **385** (2004), 305–334.
- [4] J. I. FUJII, M. FUJII, M. NAKAMURA, J. E. PEČARIĆ AND Y. SEO, *A reverse of the weighted geometric mean due to Lawson-Lim*, Linear Alg. Appl., **427** (2007), 272–284.
- [5] M. FUJII AND R. NAKAMOTO, *A geometric mean in the Furuta inequality*, Sci. Math. Japon., **55** (2002), 615–621.
- [6] M. FUJII, Y. SEO AND M. TOMINAGA, *Golden-Thompson type inequalities related to a geometric mean via Specht's ratio*, Math. Inequal. Appl., **5** (2002), 573–582.
- [7] T. FURUTA, J. MIĆIĆ, J. E. PEČARIĆ AND Y. SEO, *Mond-Pečarić Method in Operator Inequalities*, Monographs in Inequalities 1, Element, Zagreb, 2005.
- [8] S. GOLDEN, *Lower bounds for Helmholtz function*, Phys. Rev., **137** (1965), B1127–B1128.
- [9] F. HIAI AND D. PETZ, *The Golden-Thompson trace inequality is complemented*, Linear Algebra Appl., **181** (1993), 153–185.
- [10] F. KUBO AND T. ANDO, *Means of positive linear operators*, Math. Ann., **246** (1980), 205–224.
- [11] J. LAWSON AND Y. LIM, *A general framework for extending means to higher orders*, preprint. (http://arxiv.org/PS_cache/math/pdf/0612/0612293v1.pdf).
- [12] J. LAWSON AND Y. LIM, *Higher order weighted matrix means and related matrix inequalities*, preprint.
- [13] I. SEGAL, *Notes toward the construction of nonlinear relativistic quantum fields, III*, Bull. Amer. Math. Soc., **75** (1969), 1390–1395.
- [14] Y. SEO, *Reverses of the Golden-Thompson type inequalities due to Ando-Hiai-Petz*, Banach J. Math. Anal., **2** (2008), 140–149.

- [15] W. SPECHT, *Zur Theorie der elementaren Mittel*, Math. Z., **74** (1960), 91–98.
- [16] K. SYMANZIK, *Proof and refinements of an inequality of Feynman*, J.Math. Phys., **6** (1965), 1155–1156.
- [17] C. J. THOMPSON, *Inequality with applications in statistical mechanics*, J. Math. Phys., **6** (1965), 469–480.
- [18] M. TOMINAGA, *Specht's ratio in the Young inequality*, Sci. Math. Japon., **55** (2002), 585–588.
- [19] A. M. TURING, *Rounding off-errors in matrix processes*, Quart. J. Mech. Appl. Math., **1** (1948), 287–308.
- [20] T. YAMAZAKI AND M. YANAGIDA, *Characterizations of chaotic order associated with Kantorovich inequality*, Sci. Math., **2** (1999), 37–50.