

INEQUALITIES OF GRÜSS TYPE INVOLVING THE p -HH-NORMS IN THE CARTESIAN PRODUCT SPACE

EDER KIKIANTY, S. S. DRAGOMIR, AND P. CERONE

Abstract. Inequalities in estimating a type of Čebyšev functional involving the p -HH-norms are obtained by applying the known results by Grüss, Ostrowski, Čebyšev, and Lupaş. Some of these inequalities are proven to be sharp. In 1998, Dragomir and Fedotov considered a generalised Čebyšev functional, in order to approximate the Riemann-Stieltjes integral. In this paper, some sharp bounds for the generalised Čebyšev functional with convex integrand and monotonically increasing integrator are established as well. An application for the Čebyšev functional involving the p -HH-norms is also considered; and the bounds are proven to be sharp.

1. Introduction

Let $(\mathbf{X}, \|\cdot\|)$ be a normed space and consider the Cartesian product space $\mathbf{X}^2 = \{(x, y) : x, y \in \mathbf{X}\}$, where the addition and scalar multiplication are defined in the usual way. This space is a normed space together with any of the following p -norms (cf. Clarkson [2, p. 397–398], Li and Tsing [11, p. 36], and Milne [13, p. 142]):

$$\|(x, y)\|_p := \begin{cases} (\|x\|^p + \|y\|^p)^{\frac{1}{p}}, & 1 \leq p < \infty; \\ \max\{\|x\|, \|y\|\}, & p = \infty, \end{cases}$$

for any $(x, y) \in \mathbf{X}^2$. Kikianty and Dragomir [8] introduced another type of norm on \mathbf{X}^2 which is called the p -HH-norm, and is defined as follows:

$$\|(x, y)\|_{p-HH} := \left(\int_0^1 \|(1-t)x + ty\|^p dt \right)^{\frac{1}{p}}, \quad (1.1)$$

for any $1 \leq p < \infty$ and $(x, y) \in \mathbf{X}^2$. For fundamental properties of this norm, we refer to the paper by Kikianty and Dragomir [8]. We note that the p -norms and the p -HH-norms are all equivalent in \mathbf{X}^2 .

Some inequalities of Ostrowski type, which involve the p -HH-norms and the p -norms, have been considered by Kikianty, Dragomir, and Cerone [9, 10]. Continuing these works, we are interested in obtaining some new inequalities involving the p -HH-norms.

In this paper, we consider bounds in estimating the difference of $\|(\cdot, \cdot)\|_{p+q-HH}^{p+q}$ and the product $\|(\cdot, \cdot)\|_{p-HH}^p \|(\cdot, \cdot)\|_{q-HH}^q$, for any $p, q \geq 1$. This difference, however,

Mathematics subject classification (2000): 26D15, 46B20, 46C50.

Keywords and phrases: Grüss inequality, Čebyšev functional, Cartesian product, semi-inner product.

is a particular type of Čebyšev functional. In the following, we recall some classical facts concerning this functional.

For two Lebesgue integrable functions $f, g : [a, b] \rightarrow \mathbb{R}$, the Čebyšev functional is defined by

$$T(f, g) := \frac{1}{b-a} \int_a^b f(t)g(t)dt - \frac{1}{b-a} \int_a^b f(t)dt \cdot \frac{1}{b-a} \int_a^b g(t)dt.$$

In 1935, Grüss proved the following inequality which bounds the Čebyšev functional [14, p. 295–296]:

$$|T(f, g)| \leq \frac{1}{4}(\Phi - \phi)(\Gamma - \gamma), \quad (1.2)$$

provided that f and g satisfy the condition $\phi \leq f(t) \leq \Phi$ and $\gamma \leq g(t) \leq \Gamma$ for all $t \in [a, b]$. The constant $\frac{1}{4}$ is best possible and is achieved for $f(t) = g(t) = \text{sgn}(t - \frac{a+b}{2})$. Some related results regarding the sharp upper bounds for this functional can be summarised as follows:

1. Čebyšev (1882): If f, g are continuously differentiable functions on $[a, b]$, then

$$|T(f, g)| \leq \frac{1}{12} \|f'\|_{L^\infty} \|g'\|_{L^\infty} (b-a)^2, \quad (1.3)$$

where $\|f'\|_{L^\infty} := \sup_{t \in [a, b]} |f'(t)|$. Equality holds iff f' and g' are constants [14, p. 297]. Inequality (1.3) is also valid for absolutely continuous functions f, g where $f', g' \in L^\infty[a, b]$.

2. Ostrowski (1970): If f is Lebesgue integrable on $[a, b]$, $m, M \in \mathbb{R}$ such that $-\infty \leq m \leq f \leq M \leq \infty$, g is absolutely continuous and $g' \in L^\infty[a, b]$, then

$$|T(f, g)| \leq \frac{1}{8} (b-a)(M-m) \|g'\|_{L^\infty}, \quad (1.4)$$

and the constant $\frac{1}{8}$ is the best possible [14, p. 300].

3. Lupaş (1973): If f, g are absolutely continuous, $f', g' \in L^2[a, b]$, then

$$|T(f, g)| \leq \frac{1}{\pi^2} (b-a) \|f'\|_{L^2} \|g'\|_{L^2}, \quad (1.5)$$

where $\|f'\|_{L^2} = \int_a^b |f'(t)|^2 dt$. Equality holds iff

$$f(x) = A + B \sin \left[\frac{\pi}{b-a} \left(x - \frac{a+b}{2} \right) \right] \text{ and } g(x) = C + D \sin \left[\frac{\pi}{b-a} \left(x - \frac{a+b}{2} \right) \right],$$

where $A, B, C,$ and D are constants [14, p. 301].

In Section 3, we apply these results to obtain upper bounds in estimating the difference of $\|(x, y)\|_{p+q-HH}^{p+q}$ and $\|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q$ ($p, q \geq 1$). Some of these inequalities are proven to be sharp.

More results regarding the Čebyšev functional were pointed out by Dragomir and Fedotov [4]. In order to approximate the Riemann-Stieltjes integral, they considered a generalised Čebyšev functional

$$D(f, u) := \int_a^b f(t)du(t) - \frac{1}{b-a} [u(b) - u(a)] \int_a^b f(s)ds,$$

where f is Riemann integrable and Stieltjes integrable with respect to a function u . Some bounds for D , when u is monotonically nondecreasing, were obtained by Dragomir [3]; and we summarised the results as follows:

1. If $f : [a, b] \rightarrow \mathbb{R}$ is L -Lipschitzian on $[a, b]$, then

$$|D(f, u; a, b)| \leq \frac{1}{2}L(b-a) \left[u(b) - u(a) - \frac{4}{(b-a)^2} \int_a^b u(t) \left(t - \frac{a+b}{2} \right) dt \right] \leq \frac{1}{2}L(b-a)[u(b) - u(a)],$$

and the constant $\frac{1}{2}$ is best possible in both inequalities.

2. If $f : [a, b] \rightarrow \mathbb{R}$ is a function of bounded variation on $[a, b]$, and $\int_a^b f(t)du(t)$ exists, then

$$|D(f, u; a, b)| \leq \left[u(b) - u(a) - \frac{1}{b-a} \int_a^b \operatorname{sgn} \left(t - \frac{a+b}{2} \right) dt \right] \bigvee_a^b(f) \leq [u(b) - u(a)] \bigvee_a^b(f),$$

where $\bigvee_a^b(f)$ denotes the total variation of f on $[a, b]$; and the first inequality is sharp.

In Section 4, we establish some sharp bounds for the generalised Čebyšev functional D in order to approximate the Riemann-Stieltjes integral for differentiable convex integrand and monotonically increasing integrator. The result follows by utilising an Ostrowski type inequality. Then in Section 5, we apply this result for the Čebyšev functional T , and the obtained bounds are sharp. A similar result is established for general convex functions, and the obtained bounds are also sharp. By applying the result for the p -HH-norms, we also obtain some upper and lower bounds for the difference between $\|(x, y)\|_{p+q-HH}^{p+q}$ and $\|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q$ ($p, q \geq 1$). These bounds are proven to be sharp.

2. Definitions and notation

Throughout this paper, we assume that all vector spaces are over the field of real numbers and the measure that we consider is the Lebesgue measure.

Let $x, y \in \mathbf{X}$, $x \neq y$ and define the segment $[x, y] := \{(1-t)x + ty, t \in [0, 1]\}$. Let $f : [x, y] \rightarrow \mathbb{R}$ and the associated function $h : [0, 1] \rightarrow \mathbb{R}$, defined by $h(t) := f[(1-t)x + ty]$, $t \in [0, 1]$. It is well known that the function h is convex on $[0, 1]$ if and only if f is convex on $[x, y]$.

In any normed space \mathbf{X} , the norm $\|\cdot\|$ is right-(left-)Gâteaux differentiable at $x \in \mathbf{X} \setminus \{0\}$, i.e. the following limits

$$(\nabla_{+(-)}\|\cdot\|(x))(y) := \lim_{t \rightarrow 0^{+(-)}} \frac{\|x+ty\| - \|x\|}{t}$$

exist for all $y \in \mathbf{X}$ (cf. Megginson [12, p. 483–485]). The norm $\|\cdot\|$ is Gâteaux differentiable at $x \in \mathbf{X} \setminus \{0\}$ if and only if $(\nabla_+ \|\cdot\|(x))(y) = (\nabla_- \|\cdot\|(x))(y)$, for all $y \in \mathbf{X}$. The function $f_0(x) = \frac{1}{2}\|x\|^2$ ($x \in \mathbf{X}$) is convex and the following

$$(x, y)_{s(i)} := (\nabla_{+(-)} f_0(y))(x) = \lim_{t \rightarrow 0^{+(-)}} \frac{\|y + tx\|^2 - \|y\|^2}{2t},$$

exist for any $x, y \in \mathbf{X}$. They are called the superior (inferior) semi-inner products (s.i.p.) associated with the norm $\|\cdot\|$. We refer to the work by Dragomir [6] for further properties of these semi-inner products.

The function $f_p : \mathbf{X} \rightarrow \mathbb{R}$ defined by $f_p(x) = \|x\|^p$ ($1 \leq p < \infty$) is also convex, and the following limit

$$(\nabla f_p[(1-t)x + ty])(y-x) = p\|(1-t)x + ty\|^{p-1}(\nabla \|\cdot\|[(1-t)x + ty])(y-x)$$

exists almost everywhere on $[0, 1]$. Note that for any $y \neq 0$,

$$(\nabla_+ \|\cdot\|(x))(y) = (y, x)_s \quad \text{and} \quad (\nabla_- \|\cdot\|(x))(y) = (y, x)_i.$$

Therefore,

$$(\nabla f_p[(1-t)x + ty])(y-x) = p\|(1-t)x + ty\|^{p-2}(y-x, (1-t)x + ty)_{s(i)} \tag{2.1}$$

exists almost everywhere on $[0, 1]$, for any $x, y \in \mathbf{X}$, whenever $p \geq 2$; otherwise it holds for any linearly independent $x, y \in \mathbf{X}$.

3. Grüss type inequality involving the p -HH-norms

In this section, we obtain some inequalities involving the p -HH-norms in the Cartesian product space \mathbf{X}^2 from the results due to Grüss, Čebyšev, Ostrowski, and Lupaş which have been stated in Section 1.

LEMMA 1. *Let $(\mathbf{X}, \|\cdot\|)$ be a normed space, $x, y \in \mathbf{X}$, and $p, q \geq 1$. Then,*

$$\|(x, y)\|_{p+q-HH}^{p+q} \geq \|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q. \tag{3.1}$$

Equality holds in (3.1) for $x = y$.

Proof. Define $f_p(t) := \|(1-t)x + ty\|^p$, where $t \in [0, 1]$. We claim that for any $p, q \geq 1$, f_p and f_q are synchronous (similarly ordered, cf. Hardy, Littlewood, and Polya [7, p. 43] on $[0, 1]$). The proof is as follows: let $t, s \in [0, 1]$ and assume that $f_1(t) \leq f_1(s)$ (as for the other case, the proof follows similarly). Since $f_1(t) \geq 0$ for any $t \in [0, 1]$, it implies that $f_p(t) \leq f_p(s)$, for any $p \geq 1$. Thus, for any $t, s \in [0, 1]$ and $p, q \geq 1$, we have

$$[f_p(t) - f_p(s)][f_q(t) - f_q(s)] \geq 0.$$

Since f and g are synchronous, the Čebyšev inequality holds (cf. Hardy, Littlewood, and Polya [7, p. 43]), i.e.,

$$\int_0^1 f_p(t)f_q(t)dt \geq \int_0^1 f_p(t)dt \int_0^1 f_q(t)dt,$$

or, equivalently,

$$\|(x, y)\|_{p+q-HH}^{p+q} \geq \|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q,$$

as desired. It is easily shown that equality holds for $x = y$. \square

THEOREM 1. *Let $(\mathbf{X}, \|\cdot\|)$ be a normed linear space, $x, y \in \mathbf{X}$, $p, q \geq 1$, and set*

$$T_{p,q}(x, y) := \|(x, y)\|_{p+q-HH}^{p+q} - \|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q \geq 0.$$

Then,

$$0 \leq T_{p,q}(x, y) \leq \frac{1}{12}pq\|y - x\|^2 \max\{\|x\|, \|y\|\}^{p+q-2} =: C_{p,q}(x, y). \tag{3.2}$$

The constant $\frac{1}{12}$ in (3.2) is sharp.

Proof. Let $x, y \in \mathbf{X}$, and define $f(t) = \|(1-t)x + ty\|^p$ and $g(t) = \|(1-t)x + ty\|^q$ ($t \in [0, 1]$). By (2.1),

$$f'(t) = \nabla_{\pm} \|\cdot\|^{p-1} [(1-t)x + ty](y-x) = p\|(1-t)x + ty\|^{p-2}(y-x, (1-t)x + ty)_{s(i)},$$

and by the Cauchy-Schwarz inequality,

$$\begin{aligned} \|f'\|_{L^\infty} &= \sup_{t \in [0,1]} p\|(1-t)x + ty\|^{p-2} |(y-x, (1-t)x + ty)_{s(i)}| \\ &\leq p\|y-x\| \sup_{t \in [0,1]} \|(1-t)x + ty\|^{p-1} = p\|y-x\| \max\{\|x\|, \|y\|\}^{p-1}. \end{aligned}$$

Similarly for g , we have $\|g'\|_{L^\infty} \leq q\|y-x\| \max\{\|x\|, \|y\|\}^{q-1}$. Due to Čebyšev's result (1.3), we have

$$T_{p,q}(x, y) \leq \frac{1}{12}\|f'\|_{L^\infty}\|g'\|_{L^\infty} \leq \frac{1}{12}pq\|y-x\|^2 \max\{\|x\|, \|y\|\}^{p+q-2}.$$

To prove the sharpness of the constant, we assume that the (3.2) holds for a constant $A > 0$ instead of $\frac{1}{12}$, i.e.

$$\|(x, y)\|_{p+q-HH}^{p+q} - \|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q \leq A pq\|y-x\|^2 \max\{\|x\|, \|y\|\}^{p+q-2}.$$

Choose $p = 1$, $q = 1$, $\mathbf{X} = \mathbb{R}$, and $0 < x < y$, to obtain

$$\frac{1}{3}(x^2 + xy + y^2) - \left(\frac{y+x}{2}\right)^2 = \frac{1}{12}(y-x)^2 \leq A(y-x)^2.$$

Since $x \neq y$, then $A \geq \frac{1}{12}$; and the proof is completed. \square

For any x and y in the normed space $(\mathbf{X}, \|\cdot\|)$, we set the following quantities for $p, q \geq 1$:

$$G_{p,q}(x,y) := \frac{1}{4} \max\{\|x\|, \|y\|\}^{p+q}, \quad O_{p,q}(x,y) := \frac{1}{8} q \|y-x\| \max\{\|x\|, \|y\|\}^{p+q-1},$$

$$\text{and } L_{p,q}(x,y) := \frac{1}{\pi^2} p q \|y-x\|^2 \|(x,y)\|_{(2p-2)-HH}^{p-1} \|(x,y)\|_{(2q-2)-HH}^{q-1}.$$

The following proposition is due to the results by Grüss, Ostrowski, and Lupaş. However, these upper bounds are not yet proven to be sharp.

PROPOSITION 1. *Under the assumptions of Theorem 1 and the above notation, we have*

$$0 \leq T_{p,q}(x,y) \leq G_{p,q}(x,y), \quad 0 \leq T_{p,q}(x,y) \leq O_{p,q}(x,y), \quad \text{and}$$

$$0 \leq T_{p,q}(x,y) \leq L_{p,q}(x,y),$$

for any $p, q \geq 1$ and $x, y \in \mathbf{X}$. \square

Proof. Let $x, y \in \mathbf{X}$, and define $f(t) = \|(1-t)x + ty\|^p$, and $g(t) = \|(1-t)x + ty\|^q$, for $t \in [0, 1]$. Since $p, q \geq 1$, we have $0 \leq f(t) \leq \max\{\|x\|, \|y\|\}^p$ and $0 \leq g(t) \leq \max\{\|x\|, \|y\|\}^q$. Then, due to Grüss' result (1.2), we have

$$T_{p,q}(x,y) \leq \frac{1}{4} \max\{\|x\|, \|y\|\}^{p+q} = G_{p,q}(x,y).$$

Similarly to the proof of Theorem 1, we have $\|g'\|_{L^\infty} \leq q \|y-x\| \max\{\|x\|, \|y\|\}^{q-1}$. By (1.4), we have

$$T_{p,q}(x,y) \leq \frac{1}{8} \max\{\|x\|, \|y\|\}^p \|g'\|_{L^\infty} \leq \frac{1}{8} q \|y-x\| \max\{\|x\|, \|y\|\}^{p+q-1} = O_{p,q}(x,y).$$

Note that for any $p \geq 1$, we have

$$\|f'\|_{L^2} = \left[\int_0^1 |p|(1-t)x + ty\|^{p-2} (y-x, (1-t)x + ty)_{s(t)}^2 dt \right]^{\frac{1}{2}}$$

$$\leq p \|y-x\| \left[\int_0^1 \|(1-t)x + ty\|^{2p-2} dt \right]^{\frac{1}{2}} = p \|y-x\| \|(x,y)\|_{(2p-2)-HH}^{p-1}$$

by the Cauchy-Schwarz inequality; and similarly for $q \geq 1$, we have $\|g'\|_{L^2} \leq q \|y-x\| \|(x,y)\|_{(2q-2)-HH}^{q-1}$. Therefore, by Lupaş' result (1.5), we obtain

$$T_{p,q}(x,y) \leq \frac{1}{\pi^2} \|f'\|_{L^2} \|g'\|_{L^2} \leq \frac{1}{\pi^2} pq \|y-x\|^2 \|(x,y)\|_{(2p-2)\text{-HH}}^{p-1} \|(x,y)\|_{(2q-2)\text{-HH}}^{q-1} = L_{p,q}(x,y). \quad \square$$

REMARK 1. We note that none of the upper bounds for $T_{p,q}(x,y)$ that we have obtained in Proposition 1 is better than the other ones, for each $x,y \in \mathbf{X}$. For example, choose $\mathbf{X} = \mathbb{R}$, $p = q = 1$, and $x = 1$. By utilising MAPLE, we obtain (see Figure 1(a))

$$\begin{aligned} G(1,y) &\geq O(1,y) \geq L(1,y), & y \in [0, 1], \\ G(1,y) &\geq L(1,y) \geq O(1,y), & y \in [-3, -2], \\ L(1,y) &\geq G(1,y) \geq O(1,y), & y \in [-\frac{3}{2}, -1]. \end{aligned}$$

Again, by utilising MAPLE, for $p = q = 2$, and $x = -1$, we have (see Figure 1(b))

$$\begin{aligned} O(-1,y) &\geq L(-1,y) \geq G(-1,y), & y \in [\frac{3}{5}, \frac{4}{5}], \\ O(-1,y) &\geq G(-1,y)(x,y) \geq L(-1,y), & y \in [0, \frac{2}{5}], \\ L(-1,y) &\geq O(-1,y)(x,y) \geq G(-1,y), & y \in [\frac{19}{20}, 1]. \end{aligned}$$




Figure 1.

PROBLEM 1. Are the constants $\frac{1}{4}$, $\frac{1}{8}$ and $\frac{1}{\pi^2}$ in Proposition 1 the best possible?

4. New bounds for the generalised Čebyšev functional D

The following result gives upper and lower bounds for the generalised Čebyšev functional $D(\cdot, \cdot)$ in order to approximate the Riemann-Stieltjes integral.

THEOREM 2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable convex function, and $u : [a, b] \rightarrow \mathbb{R}$ be a monotonically increasing function. Then,*

$$\begin{aligned} & \frac{b-a}{2} [f'(a)u(b) + f'(b)u(a)] - \int_a^b u(t) \left[\frac{t-a}{b-a} f'(a) + \frac{b-t}{b-a} f'(b) \right] dt \\ & \leq D(f, u) \leq \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) du(t). \end{aligned} \quad (4.1)$$

The constants $\frac{1}{2}$ and 1 in (4.1) are sharp.

Proof. Since f is a differentiable convex function on $[a, b]$, we have the following Ostrowski type inequality (cf. Dragomir [5])

$$\begin{aligned} \frac{1}{2} [(b-t)^2 - (t-a)^2] f'(t) & \leq \int_a^b f(s) ds - (b-a)f(t) \\ & \leq \frac{1}{2} [(b-t)^2 f'(b) - (t-a)^2 f'(a)], \end{aligned} \quad (4.2)$$

for any $t \in [a, b]$. Since u is a monotonically increasing function on $[a, b]$, we may integrate the inequality (4.2) (in the Riemann-Stieltjes sense) with respect to u , i.e.

$$\begin{aligned} \frac{1}{2} \int_a^b [(b-t)^2 - (t-a)^2] f'(t) du(t) & \leq \int_a^b \left[\int_a^b f(s) ds - (b-a)f(t) \right] du(t) \\ & \leq \frac{1}{2} \int_a^b [(b-t)^2 f'(b) - (t-a)^2 f'(a)] du(t). \end{aligned} \quad (4.3)$$

Note that

$$\frac{1}{2} \int_a^b [(b-t)^2 - (t-a)^2] f'(t) du(t) = (b-a) \int_a^b \left(\frac{b+a}{2} - t \right) f'(t) du(t),$$

and

$$\begin{aligned} \int_a^b \left[\int_a^b f(s) ds - (b-a)f(t) \right] du(t) & = \int_a^b f(s) ds \int_a^b du(t) - (b-a) \int_a^b f(t) du(t) \\ & = [u(b) - u(a)] \int_a^b f(s) ds - (b-a) \int_a^b f(t) du(t), \end{aligned}$$

and, using integration by parts

$$\begin{aligned} & \frac{1}{2} \int_a^b [(b-t)^2 f'(b) - (t-a)^2 f'(a)] du(t) \\ & = \frac{1}{2} (b-a)^2 [-f'(b)u(a) - f'(a)u(b)] + \int_a^b u(t) [(b-t)f'(b) + (t-a)f'(a)] dt. \end{aligned}$$

Therefore, by (4.3) we get

$$\begin{aligned}
 & (b-a) \int_a^b \left(\frac{b+a}{2} - t \right) f'(t) du(t) \\
 & \leq [u(b) - u(a)] \int_a^b f(s) ds - (b-a) \int_a^b f(t) du(t) \\
 & \leq \frac{1}{2}(b-a)^2 [-f'(b)u(a) - f'(a)u(b)] + \int_a^b u(t)[(b-t)f'(b) + (t-a)f'(a)] dt,
 \end{aligned} \tag{4.4}$$

and the proof follows by multiplying inequality (4.4) by $(-\frac{1}{b-a})$.

The sharpness of the constants follows by a particular case which will be stated in Corollary 3. \square

COROLLARY 1. *Under the assumptions of Theorem 2, if $f'(b) = -f'(a)$, then*

$$\begin{aligned}
 & f'(a) \left[\frac{b-a}{2}(u(b) - u(a)) - \frac{1}{b-a} \int_a^b u(t) (2t - (a+b)) dt \right] \\
 & \leq D(f, u) \leq \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) du(t).
 \end{aligned} \tag{4.5}$$

REMARK 2. A common example of such function is a function defined on interval $[a, b]$ which is symmetric with respect to the midpoint $\frac{a+b}{2}$, e.g. $f(t) = |t - \frac{a+b}{2}|^p$, where $p \geq 1$.

COROLLARY 2. *Under the assumptions of Theorem 2, if $f'(a) = -f'(b)$ and f'' exists, then*

$$\begin{aligned}
 & f'(a) \left[\frac{b-a}{2}(u(b) - u(a)) - \frac{1}{b-a} \int_a^b u(t) (2t - (a+b)) dt \right] \\
 & \leq D(f, u) \leq \left(\frac{b-a}{2} \right) f'(b)[u(b) - u(a)] - \int_a^b u(t) \left[f'(t) + \left(t - \frac{b+a}{2} \right) f''(t) \right] dt.
 \end{aligned} \tag{4.6}$$

Proof. This is a particular case of Corollary 1. Note that

$$\begin{aligned}
 & \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) du(t) \\
 & = \left(\frac{b-a}{2} \right) f'(b)[u(b) - u(a)] - \int_a^b u(t) \left[f'(t) + \left(t - \frac{b+a}{2} \right) f''(t) \right] dt,
 \end{aligned}$$

and the details are omitted. \square

PROBLEM 2. Are the inequalities in Corollaries 1 and 2 sharp?

5. Application for the Čebyšev functional

In this section, we apply the result of Section 4 to obtain bounds for the classical Čebyšev functional.

COROLLARY 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable convex function, and $g : [a, b] \rightarrow \mathbb{R}$ be a nonnegative Lebesgue integrable function. Then,*

$$\begin{aligned} & \frac{1}{2} \int_a^b \left[\left(\frac{t-a}{b-a} \right)^2 f'(a) - \left(\frac{b-t}{b-a} \right)^2 f'(b) \right] g(t) dt \\ & \leq T(f, g) \leq \frac{1}{b-a} \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) g(t) dt. \end{aligned} \quad (5.1)$$

The constants $\frac{1}{2}$ and 1 in (5.1) are sharp.

Proof. Recall that Theorem 2 gives us

$$\begin{aligned} & \frac{1}{2(b-a)} \int_a^b [(t-a)^2 f'(a) - (b-t)^2 f'(b)] du(t) \\ & \leq D(f, u) \leq \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) du(t). \end{aligned} \quad (5.2)$$

Since g is nonnegative on $[a, b]$, $u(t) = \int_a^t g(s) ds$ is monotonically increasing on $[a, b]$. Thus, inequality (5.1) follows by applying (5.2) to u and multiply the obtained inequality by $\frac{1}{b-a}$. The sharpness of the constants in (5.1) is demonstrated by choosing $f(t) = g(t) = t$ on $[a, b]$, and the details are omitted. \square

EXAMPLE 1. Let $f(t) = g(t) = \frac{1}{t}$ defined on $[x, y]$, where $x, y > 0$. Then by Corollary 3, we obtain

$$0 \leq \left(\frac{1}{G(x, y)} \right)^2 - \left(\frac{1}{L(x, y)} \right)^2 \leq \left(\frac{y-x}{2xy} \right)^2, \quad (5.3)$$

where $G(x, y)$ and $L(x, y)$ are the geometric mean and logarithmic mean of x and y , respectively (note that $G(x, y) = \sqrt{xy}$ and $L(x, y) = \frac{x-y}{\log x - \log y}$). We do not consider the lower bound in this case, as it is not always positive.

5.1. Čebyšev functional for convex functions

In Corollary 3, we assume that f is a differentiable convex function. However, we can ‘drop’ the assumption of differentiability, and get a similar result for general convex functions, where the derivative exists almost everywhere.

PROPOSITION 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function, and $g : [a, b] \rightarrow \mathbb{R}$ be a nonnegative Lebesgue integrable function. Then,

$$\begin{aligned} & \frac{1}{2} \int_a^b \left[\left(\frac{t-a}{b-a} \right)^2 f'(a) - \left(\frac{b-t}{b-a} \right)^2 f'(b) \right] g(t) dt \\ & \leq T(f, g) \leq \frac{1}{b-a} \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) g(t) dt. \end{aligned} \quad (5.4)$$

The constants 1 and $\frac{1}{2}$ in (5.4) are sharp.

Proof. Since f is a convex function on $[a, b]$, we have the following Ostrowski type inequality for any $t \in [a, b]$ (cf. Dragomir [5])

$$\begin{aligned} & \frac{1}{2} [(b-t)^2 f'_+(t) - (t-a)^2 f'_-(t)] \\ & \leq \int_a^b f(s) ds - (b-a)f(t) \leq \frac{1}{2} [(b-t)^2 f'_-(b) - (t-a)^2 f'_+(a)] \end{aligned} \quad (5.5)$$

We multiply the (5.5) by $g(t)$, take the integral over $[a, b]$ and multiply it by $-\frac{1}{(b-a)^2}$ to obtain

$$\begin{aligned} & \frac{1}{2} \int_a^b \left[\left(\frac{t-a}{b-a} \right)^2 f'_+(a) - \left(\frac{b-t}{b-a} \right)^2 f'_-(b) \right] g(t) dt \\ & \leq T(f, g) \leq \frac{1}{2} \int_a^b \left[\left(\frac{t-a}{b-a} \right)^2 f'_-(t) - \left(\frac{b-t}{b-a} \right)^2 f'_+(t) \right] g(t) dt \end{aligned}$$

Since f is convex, then f' exists almost everywhere and we may write $f'(t) = f'_\pm(t)$, for almost every $t \in [a, b]$, and the details are omitted. The sharpness of the constants follows by Remark 3. \square

In a similar way, we have the generalised version of Theorem 2 as follows, and the proof is omitted.

PROPOSITION 3. Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function, and $u : [a, b] \rightarrow \mathbb{R}$ be a monotonically increasing function. Then,

$$\begin{aligned} & \frac{b-a}{2} [f'(a)u(b) + f'(b)u(a)] - \int_a^b u(t) \left[\frac{t-a}{b-a} f'(a) + \frac{b-t}{b-a} f'(b) \right] dt \\ & \leq D(f, u) \leq \int_a^b \left(t - \frac{b+a}{2} \right) f'(t) du(t). \end{aligned} \quad (5.6)$$

The constants $\frac{1}{2}$ and 1 in (5.6) are sharp.

The following result is a consequence of Proposition 2, for convex functions on linear spaces.

COROLLARY 4. *Let \mathbf{X} be a linear space and x, y be two distinct vectors in \mathbf{X} . Let g be a nonnegative functional on $[x, y]$ such that $\int_0^1 g[(1-t)x + ty]dt < \infty$. Then, for any convex function f defined on the segment $[x, y]$ and $t \in (0, 1)$, we have*

$$\begin{aligned} & \frac{1}{2} \int_0^1 [t^2(\nabla f(x))(y-x) - (1-t)^2(\nabla f(y))(y-x)] g[(1-t)x + ty]dt \\ & \leq \int_0^1 f[(1-t)x + ty]g[(1-t)x + ty]dt \\ & \quad - \int_0^1 f[(1-t)x + ty]dt \int_0^1 g[(1-t)x + ty]dt \\ & \leq \int_0^1 \left(t - \frac{1}{2}\right) (\nabla f[(1-t)x + ty])(y-x)g[(1-t)x + ty]dt. \end{aligned} \tag{5.7}$$

The constants $\frac{1}{2}$ and 1 in (5.7) are sharp.

Proof. Consider the functions h, k defined on $[0, 1]$ by $h(t) = f[(1-t)x + ty]$ and $k(t) = g[(1-t)x + ty]$. Since f is convex on the segment $[x, y]$, then h is also convex on $[0, 1]$. Thus we may apply Proposition 2 to h and k . Note that $h'_\pm(t) = (\nabla_\pm f[(1-t)x + ty])(y-x)$, by the chain rule; and since h is convex,

$$h'(t) := h'_\pm(t) = (\nabla_\pm f[(1-t)x + ty])(y-x) =: (\nabla f[(1-t)x + ty])(y-x)$$

exists almost everywhere on $[0, 1]$ (we get a similar identity for k). The proof for the sharpness follows by the particular case given later in Corollary 5. \square

5.2. Application to the p -HH-norms

Let $(\mathbf{X}, \|\cdot\|)$ be a normed space. Recall from Lemma 1 that

$$T_{p,q}(x, y) := \|(x, y)\|_{p+q-HH}^{p+q} - \|(x, y)\|_{p-HH}^p \|(x, y)\|_{q-HH}^q \geq 0,$$

for any $x, y \in \mathbf{X}$ and $p, q \geq 1$.

COROLLARY 5. *Under the above notation and assumptions, we have*

$$\begin{aligned} & \frac{1}{2} p \int_0^1 [t^2 \|x\|^{p-2}(y-x, x) - (1-t)^2 \|y\|^{p-2}(y-x, y)] \|(1-t)x + ty\|^q dt \\ & \leq T_{p,q}(x, y) \leq p \int_0^1 \left(t - \frac{1}{2}\right) \|(1-t)x + ty\|^{p+q-2}(y-x, (1-t)x + ty) dt, \end{aligned} \tag{5.8}$$

for any $x, y \in \mathbf{X}$ whenever $p \geq 2$. If $1 \leq p < 2$, then the inequality (5.8) holds for any nonzero $x, y \in \mathbf{X}$.

The constants $\frac{1}{2}$ and 1 are sharp in (5.8).

Proof. Define $f[(1-t)x+ty] = \|(1-t)x+ty\|^p$ and $g[(1-t)x+ty] = \|(1-t)x+ty\|^q$ for $t \in [0, 1]$. By (2.1), the following exists almost everywhere on $[0, 1]$,

$$(\nabla \|\cdot\|^p[(1-t)x+ty])(y-x) = p\|(1-t)x+ty\|^{p-2}(y-x, (1-t)x+ty)_{s(i)},$$

provided that $p \geq 2$; otherwise, it holds for any linearly independent x and y . By denoting $(\cdot, \cdot) := (\cdot, \cdot)_{s(i)}$, we have

$$(\nabla \|\cdot\|^p[(1-t)x+ty])(y-x) = p\|(1-t)x+ty\|^{p-2}(y-x, (1-t)x+ty),$$

and we obtain the similar identity for g . Therefore, by Corollary 4,

$$\begin{aligned} & \frac{1}{2}p \int_0^1 [t^2\|x\|^{p-2}(y-x, x) - (1-t)^2\|y\|^{p-2}(y-x, y)] \|(1-t)x+ty\|^q dt \\ & \leq T_{p,q}(x, y) \leq p \int_0^1 \left(t - \frac{1}{2}\right) \|(1-t)x+ty\|^{p+q-2}(y-x, (1-t)x+ty) dt, \end{aligned}$$

for any $x, y \in \mathbf{X}$ whenever $p \geq 2$; otherwise, it holds for any nonzero $x, y \in \mathbf{X}$. The proof for the sharpness of the constants follows by a particular case which will be stated in Remark 3. \square

REMARK 3. (Case of inner product space) Let $(\mathbf{X}, \langle \cdot, \cdot \rangle)$ be an inner product space and x, y be two distinct vectors in \mathbf{X} . Then, for any $p, q \geq 1$, we have

$$\begin{aligned} & \frac{1}{2}p \int_0^1 \langle y-x, t^2\|x\|^{p-2}x - (1-t)^2\|y\|^{p-2}y \rangle \|(1-t)x+ty\|^q dt \tag{5.9} \\ & \leq T_{p,q}(x, y) \leq p \int_0^1 \left(t - \frac{1}{2}\right) \|(1-t)x+ty\|^{p+q-2} \langle y-x, (1-t)x+ty \rangle dt. \end{aligned}$$

If $p = q = 1$, then

$$\begin{aligned} & \frac{1}{2} \int_0^1 \|(1-t)x+ty\| \left\langle y-x, \frac{t^2}{\|x\|}x - \frac{(1-t)^2}{\|y\|}y \right\rangle dt \tag{5.10} \\ & \leq \|(x, y)\|_{2-HH}^2 - \|(x, y)\|_{1-HH}^2 \leq \frac{1}{12}\|y-x\|^2. \end{aligned}$$

Note that when $\mathbf{X} = \mathbb{R}$, and $x, y > 0$ (some details are omitted),

$$\begin{aligned} & \frac{1}{2} \int_0^1 ((1-t)x+ty)(y-x) (t^2 - (1-t)^2) dt \\ & = \frac{y-x}{2} \int_0^1 (t^2(1-t) - (1-t)^3)x + (t^3 - t(1-t)^2)y dt \\ & = \frac{y-x}{2} \left(\frac{y-x}{6}\right) = \frac{1}{12}(y-x)^2, \end{aligned}$$

and

$$\|(x, y)\|_{2-HH}^2 - \|(x, y)\|_{1-HH}^2 = \frac{y^3 - x^3}{3(y-x)} - \left(\frac{y+x}{2}\right)^2 = \frac{1}{12}(y-x)^2.$$

Therefore, we obtain equality in (5.10).

REMARK 4. Although the inequality that we obtain in Corollary 5 is sharp, the bounds are complicated to compute. We remark that the lower bound is not always positive, e.g., take $\mathbf{X} = \mathbb{R}$, $p = q = 1$, $x = -1, y = 1$, we have

$$\frac{1}{2}p \int_0^1 \left(t^2 |x|^{p-2} (y-x)x - (1-t)^2 |y|^{p-2} (y-x)y \right) (|(1-t)x + ty|)^q dt = -\frac{3}{8}.$$

In this case, the lower bound cannot be used to improve the Čebyšev inequality. We obtain coarser but simpler upper bounds for $T_{p,q}(x, y)$, as follows:

$$\begin{aligned} 0 \leq T_{p,q}(x, y) &\leq p \int_0^1 \left(t - \frac{1}{2} \right) \|(1-t)x + ty\|^{p+q-2} (y-x, (1-t)x + ty) dt, \\ &\leq p \|y-x\| \int_0^1 \left| t - \frac{1}{2} \right| \|(1-t)x + ty\|^{p+q-1} dt, \\ &\leq p \|y-x\| \begin{cases} \frac{1}{2} \|(x, y)\|_{(p+q-1)\text{-HH}}^{p+q-1} \\ \left(\frac{1}{2^{s'(s'+1)}} \right)^{\frac{1}{s'}} \|(x, y)\|_{(p+q-1)s}^{p+q-1}, s > 1, \frac{1}{s} + \frac{1}{s'} = 1; \\ \frac{1}{4} \max\{\|x\|, \|y\|\}^{p+q-1}. \end{cases} \quad (5.11) \end{aligned}$$

REMARK 5. Although, in general, these upper bounds are not always better than those obtained in Section 3, we remark that under certain conditions, they are better. For example, when $p \leq \frac{1}{2}q$, we have

$$\frac{1}{4}p \|y-x\| \max\{\|x\|, \|y\|\}^{p+q-1} \leq O_{p,q}(x, y)$$

(recall that $O_{p,q}(x, y) := \frac{1}{8}q \|y-x\| \max\{\|x\|, \|y\|\}^{p+q-1}$). Also, when $p \leq 1$ and $\|y-x\| \leq \max\{\|x\|, \|y\|\}$, we have

$$\frac{1}{4}p \|y-x\| \max\{\|x\|, \|y\|\}^{p+q-1} \leq G_{p,q}(x, y)$$

(recall that $G_{p,q}(x, y) := \frac{1}{4} \max\{\|x\|, \|y\|\}^{p+q}$).

PROBLEM 3. Are the constants $\frac{1}{2}$, $\left(\frac{1}{2^{s'(s'+1)}} \right)^{\frac{1}{s'}}$, and $\frac{1}{4}$ in (5.11) the best possible?

REFERENCES

- [1] P.S. BULLEN, *Handbook of Means and their Inequalities*, Mathematics and its Applications, vol. 560, Kluwer Academic Publishers Group, Dordrecht, 2003. Revised from the 1988 original [P. S. Bullen, D. S. Mitrinović and P. M. Vasić, *Means and their inequalities*, Reidel, Dordrecht].
- [2] J.A. CLARKSON, *Uniformly convex spaces*, Trans. Amer. Math. Soc., **40**, 3 (1936), 396–414.
- [3] S.S. DRAGOMIR, *Inequalities of Grüss type for the Stieltjes integral and applications*, Kragujevac J. Math., **26** (2004), 89–122.

- [4] S.S. DRAGOMIR AND I.A. FEDOTOV, *An inequality of Grüss' type for Riemann-Stieltjes integral and applications for special means*, Tamkang J. Math., **29**, 4 (1998), 287–292.
- [5] S.S. DRAGOMIR, *An Ostrowski like inequality for convex functions and applications*, Rev. Mat. Complut., **16**, 2 (2003), 373–382.
- [6] S.S. DRAGOMIR, *Semi-inner Products and Applications*, Nova Science Publishers, Inc., Hauppauge, NY, 2004.
- [7] G.H. HARDY, J.E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, Cambridge, at the University Press, 1952, 2nd ed.
- [8] E. KIKIANTY AND S.S. DRAGOMIR, *Hermite-Hadamard's inequality and the p -HH-norm on the Cartesian product of two copies of a normed space*, Math. Inequal. Appl., to appear.
- [9] E. KIKIANTY, S.S. DRAGOMIR, AND P. CERONE, *Ostrowski type inequality for absolutely continuous functions on segments of linear spaces*, Bull. Korean Math. Soc., **45**, 4 (2008), 763–780.
- [10] E. KIKIANTY, S.S. DRAGOMIR, AND P. CERONE, *Sharp inequalities of Ostrowski type for convex functions defined on linear spaces and applications*, Comput. Math. Appl., **56**, 9 (2008), 2235–2246.
- [11] C.-K. LI AND N.-K. TSING, *Norms on Cartesian product of linear spaces*, Tamkang J. Math., **21**, 1 (1990), 35–39.
- [12] R.E. MEGGINSON, *An Introduction to Banach Space Theory*, Graduate Texts in Mathematics, vol. 183, Springer-Verlag, New York, 1998.
- [13] R.D. MILNE, *Applied Functional Analysis*, An introductory treatment, Applicable Mathematics Series, Pitman Advanced Publishing Program, Boston, Mass., 1980.
- [14] D.S. MITRINOVIĆ, J.E. PEČARIĆ, AND A.M. FINK, *Classical and new inequalities in analysis*, Mathematics and its Applications (East European Series), vol. 61, Kluwer Academic Publishers Group, Dordrecht, 1993.

(Received October 24, 2008)

Eder Kikianty
School of Engineering and Science
Victoria University
PO Box 14428, Melbourne 8001
Victoria
Australia

e-mail: eder.kikianty@research.vu.edu.au

S. S. Dragomir
School of Engineering and Science
Victoria University
PO Box 14428, Melbourne 8001
Victoria
Australia

e-mail: sever.dragomir@vu.edu.au

P. Cerone
School of Engineering and Science
Victoria University
PO Box 14428, Melbourne 8001
Victoria
Australia

e-mail: pietro.cerone@vu.edu.au