

ON MONOTONICITY OF SOME OPERATOR FUNCTIONS RELATED TO ORDER PRESERVING OPERATOR INEQUALITIES

TATSUYA KOIZUMI AND KEIICHI WATANABE

(Communicated by J. Pečarić)

Abstract. In this paper, we shall show an equivalence relation between extensions of order preserving operator inequalities and monotonicity of related operator functions.

1. Introduction

A capital letter means a bounded linear operator on a complex Hilbert space H . An operator T is said to be positive semidefinite (denoted by $0 \leq T$) if $0 \leq (Tx, x)$ for all $x \in H$ and also an operator T is said to be positive definite (denoted by $0 < T$) if T is positive semidefinite and invertible.

THEOREM 1. [10], [13] *Let $0 \leq p \leq 1$. If $0 \leq B \leq A$, then $B^p \leq A^p$ holds.*

It is well known that $0 \leq B \leq A$ does not always ensure $B^p \leq A^p$ for $1 < p$ in general. The next result has been obtained from this point of view.

THEOREM 2. [1] *Let $0 \leq p$, $1 \leq q$ and $0 \leq r$ with $p+r \leq (1+r)q$. If $0 \leq B \leq A$, then we have the following inequality:*

$$\left(A^{\frac{r}{q}} B^p A^{\frac{r}{q}} \right)^{\frac{1}{q}} \leq A^{\frac{p+r}{q}}.$$

It is known that the next Theorem 3 is an equivalence between generalized Furuta inequalities and monotonicity of related operator functions.

THEOREM 3. [6] *The following statements hold and follow from each other:*

(1) *Let $1 \leq p$, $1 \leq s$, $0 \leq t \leq 1$ and $t \leq r$. If $0 \leq B \leq A$ with $0 < A$, then the following inequality holds:*

$$\left\{ A^{\frac{r}{q}} \left(A^{-\frac{1}{2}} B^p A^{-\frac{1}{2}} \right)^s A^{\frac{r}{q}} \right\}^{\frac{1-t+r}{(p-t)s+r}} \leq A^{1-t+r}.$$

Mathematics subject classification (2010): 47A63.

Keywords and phrases: Löwner-Heinz inequality, Furuta inequality, order preserving operator inequality.

Supported in part by Grants-in-Aid for Scientific Research, Japan Society of the Promotion of Science.

(2) Let $0 \leq t \leq q \leq 1$, $q \leq p$, $1 \leq s$ and $t \leq r$. If $0 \leq B \leq A$ with $0 < A$, then the following inequality holds:

$$\left\{ A^{\frac{t}{2}} \left(A^{-\frac{t}{2}} B^p A^{-\frac{t}{2}} \right)^s A^{\frac{r}{2}} \right\}^{\frac{q-t+r}{(p-t)s+r}} \leq A^{q-t+r}.$$

(3) Let $0 \leq t \leq 1$ and $1 \leq p$. If $0 \leq B \leq A$ with $0 < A$, then

$$F(A, B, r, s) = A^{-\frac{r}{2}} \left\{ A^{\frac{t}{2}} \left(A^{-\frac{t}{2}} B^p A^{-\frac{t}{2}} \right)^s A^{\frac{r}{2}} \right\}^{\frac{1-t+r}{(p-t)s+r}} A^{-\frac{r}{2}}$$

is decreasing for $t \leq r$ and $1 \leq s$.

(4) Let $0 \leq t \leq 1$, $0 \leq q$ and $t \leq p$. If $0 \leq B \leq A$ with $0 < A$, then

$$G(A, B, r, s) = A^{-\frac{r}{2}} \left\{ A^{\frac{t}{2}} \left(A^{-\frac{t}{2}} B^p A^{-\frac{t}{2}} \right)^s A^{\frac{r}{2}} \right\}^{\frac{q-t+r}{(p-t)s+r}} A^{-\frac{r}{2}}$$

is decreasing for $t \leq r$ and $1 \leq s$ such that $q-t \leq (p-t)s$.

(1) and (3) in Theorem 3 have been proved as a theorem in [3]. (2) in Theorem 3 was shown in [9] and (4) in Theorem 3 was shown in [8] as an extension of [7]. Moreover, (1), (2), (3) and (4) have been proved to be equivalent each other in [6] (see also [11]).

LEMMA 4. [2] Let A, B be positive definite and let λ be a real number. Then

$$(ABA)^\lambda = AB^{\frac{1}{2}} \left(B^{\frac{1}{2}} A^2 B^{\frac{1}{2}} \right)^{\lambda-1} B^{\frac{1}{2}} A.$$

DEFINITION 1. [12] Let n be a natural number. We set

$$\alpha(2n) = 1 - t_1 + t_2 - \dots - t_{2n-1} + t_{2n}$$

$$\psi(2n) = \{ \dots (((p_1 - t_1)p_2 + t_2)p_3 - t_3)p_4 + t_4)p_5 - \dots - t_{2n-1} \} p_{2n} + t_{2n}.$$

Furuta [4] obtained an extension of (1) in Theorem 3 stated before. The next Theorem 5 is Corollary 11 in [12], which is an extension of Theorem 3.3 in [4].

THEOREM 5. [12] Let n be a natural number. Let $1 \leq p_j$ ($j = 1, \dots, 2n$), $0 \leq t_{2k-1} \leq 1$ and $t_{2k-1} \leq t_{2k}$ ($k = 1, \dots, n$). If $0 \leq B \leq A$ with $0 < A$, then the following inequality holds:

$$\left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} \leq A^{\alpha(2n)}.$$

2. Results

For the convenience we introduce the following notations.

DEFINITION 2. Let n be a natural number. We set

$$\begin{aligned} \tilde{\alpha}(2n) &= q - r_1 + r_2 - \dots - r_{2n-1} + r_{2n} \\ \tilde{\psi}(2n) &= \{\dots(((s_1 - r_1)s_2 + r_2)s_3 - r_3)s_4 + r_4)s_5 - \dots - r_{2n-1}\}s_{2n} + r_{2n} \\ \tilde{\gamma}(2n - 1) &= \{\dots(((s_1 - r_1)s_2 + r_2)s_3 - r_3)s_4 + r_4)s_5 - \dots + r_{2n-2}\}s_{2n-1} - r_{2n-1}. \end{aligned}$$

DEFINITION 3. Let n be a natural number. We set

$$\begin{aligned} F_{2n}(A, B, t_{2n}, p_{2n}) &= \\ A^{-\frac{t_{2n}}{2}} \left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} A^{-\frac{t_{2n}}{2}} \\ G_{2n}(A, B, r_{2n}, s_{2n}) &= \\ A^{-\frac{r_{2n}}{2}} \left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\tilde{\psi}(2n)}} A^{-\frac{r_{2n}}{2}}. \end{aligned}$$

The purpose of this paper is to show the following Theorem 6, which is an extension of Theorem 3. The method of our proof of Theorem 6 is almost similar to that of Furuta, Hashimoto and Ito [6].

THEOREM 6. *The following statements hold and follow from each other:*

(1) *Let n be a natural number. Let $1 \leq p_j$ ($j = 1, \dots, 2n$), $0 \leq t_{2k-1} \leq 1$ and $t_{2k-1} \leq t_{2k}$ ($k = 1, \dots, n$). If $0 \leq B \leq A$ with $0 < A$, then the following inequality holds:*

$$\begin{aligned} &\left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} \\ &\leq A^{\alpha(2n)}. \end{aligned}$$

(2) *Let n be a natural number. Let $0 \leq r_{2k-1} \leq q \leq 1$ ($k = 1, 2, \dots, n$), $q \leq s_1$, $1 \leq s_j$ ($j = 2, 3, \dots, 2n$), $r_{2k-1} \leq r_{2k}$ ($k = 1, 2, \dots, n$). If $0 \leq B \leq A$ with $0 < A$, then the following inequality holds:*

$$\begin{aligned} &\left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\tilde{\psi}(2n)}} \\ &\leq A^{\tilde{\alpha}(2n)}. \end{aligned}$$

(3) *Let n be a natural number. Let $0 \leq t_{2k-1} \leq 1$ ($k = 1, 2, \dots, n$), $1 \leq p_j$ ($j = 1, \dots, 2n - 1$) and $t_{2i-1} \leq t_{2i}$ ($i = 1, \dots, n - 1$). If $0 \leq B \leq A$ with $0 < A$, then*

$$\begin{aligned} F_{2n}(A, B, t_{2n}, p_{2n}) &= \\ A^{-\frac{t_{2n}}{2}} \left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} A^{-\frac{t_{2n}}{2}} \end{aligned}$$

is decreasing for $t_{2n-1} \leq t_{2n}$ and $1 \leq p_{2n}$.

(4) Let n be a natural number. Let $0 \leq r_{2k-1} \leq 1$ ($k = 1, 2, \dots, n$), $0 \leq q$, $r_{2k-1} \leq s_1$ ($k = 1, 2, \dots, n$), $1 \leq s_j$ ($j = 2, 3, \dots, 2n-1$) and $r_{2i-1} \leq r_{2i}$ ($i = 1, \dots, n-1$). If $0 \leq B \leq A$ with $0 < A$, then

$$G_{2n}(A, B, r_{2n}, s_{2n}) = A^{-\frac{r_{2n}}{2}} \left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\tilde{\psi}(2n)}} A^{-\frac{r_{2n}}{2}}$$

is a decreasing function of $r_{2n-1} \leq r_{2n}$ and $1 \leq s_{2n}$ such that $\tilde{\alpha}(2n) \leq \tilde{\psi}(2n)$.

Proof. We may assume that B is invertible without loss of generality. (1) has been already proved in [12]. We have only to prove the equivalence among (1), (2), (3) and (4).

$$(1) \Rightarrow (2)$$

We may assume that $q \neq 0$, for if $q = 0$, then $r_{2k-1} = 0$ ($k = 1, 2, \dots, n$) and it is just the case of Theorem 3.1 in [5]. One can also obtain the result by using the technique in [14].

Since $q \in (0, 1]$, $B^q \leq A^q$.

Put $A_1 = A^q$, $B_1 = B^q$, $p_1 = \frac{s_1}{q}$, $p_j = s_j$ ($j = 2, 3, \dots, 2n$) and $t_i = \frac{r_i}{q}$ ($i = 1, 2, \dots, 2n$).

Then it is obvious that $0 \leq B_1 \leq A_1$, $1 \leq p_j$ ($j = 1, 2, \dots, 2n$), $0 \leq t_{2k-1} \leq 1$ and $t_{2k-1} \leq t_{2k}$ ($k = 1, 2, \dots, n$).

Therefore we have the following inequality by (1):

$$\left\{ A_1^{\frac{t_{2n}}{2}} \left(A_1^{-\frac{t_{2n-1}}{2}} \dots \left(A_1^{\frac{t_2}{2}} \left(A_1^{-\frac{t_1}{2}} B_1^{p_1} A_1^{-\frac{t_1}{2}} \right)^{p_2} A_1^{\frac{t_2}{2}} \right)^{p_3} \dots A_1^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A_1^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} \leq A_1^{\alpha(2n)}.$$

It is easy to see that

$$A_1^{\alpha(2n)} = A^{\tilde{\alpha}(2n)}, A_1^{\frac{t_{2j}}{2}} = A^{\frac{r_{2j}}{2}} \quad (j = 1, 2, \dots, n), B_1^{p_1} = B^{s_1}$$

and

$$\begin{aligned} \frac{\alpha(2n)}{\psi(2n)} &= \frac{1 - t_1 + t_2 - \dots - t_{2n-1} + t_{2n}}{\{ \dots ((p_1 - t_1)p_2 + t_2)p_3 - \dots - t_{2n-1} \} p_{2n} + t_{2n}} \\ &= \frac{1 - \frac{r_1}{q} + \frac{r_2}{q} - \dots - \frac{r_{2n-1}}{q} + \frac{r_{2n}}{q}}{\{ \dots \left(\left(\frac{s_1}{q} - \frac{r_1}{q} \right) s_2 + \frac{r_2}{q} \right) s_3 - \dots - \frac{r_{2n-1}}{q} \} s_{2n} + \frac{r_{2n}}{q}} \\ &= \frac{q - r_1 + r_2 - \dots - r_{2n-1} + r_{2n}}{\{ \dots ((s_1 - r_1)s_2 + r_2)s_3 - \dots - r_{2n-1} \} s_{2n} + r_{2n}} \\ &= \frac{\tilde{\alpha}(2n)}{\tilde{\psi}(2n)}, \end{aligned}$$

that is, we have the following inequality:

$$\left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\psi(2n)}} \leq A^{\tilde{\alpha}(2n)}.$$

(2) ⇒ (4)

Put $q_1 = \max \{r_{2k-1}; k = 1, 2, \dots, n\}$. Then we have the following inequality by (2):

$$\left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{q_1 - r_1 + r_2 - \dots - r_{2n-1} + r_{2n}}{\psi(2n)}} \leq A^{q_1 - r_1 + r_2 - \dots - r_{2n-1} + r_{2n}}.$$

Raise each side of this inequality to the power $\frac{r_{2n}}{q_1 - r_1 + r_2 - \dots - r_{2n-1} + r_{2n}} \in [0, 1]$, then we have the following inequality by Theorem 1:

$$\left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n-1}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n-1}}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{r_{2n}}{\psi(2n)}} \leq A^{r_{2n}}.$$

Put

$$D = A^{-\frac{r_{2n-1}}{2}} \left(A^{\frac{r_{2n-2}}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{\frac{r_{2n-2}}{2}} \right)^{s_{2n-1}} A^{-\frac{r_{2n-1}}{2}}.$$

Then we have the following inequalities:

$$\begin{aligned} A^{r_{2n}} &\geq \left\{ A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{r_{2n}}{\psi(2n)}} \dots (a) \\ &= A^{\frac{r_{2n}}{2}} D^{\frac{s_{2n}}{2}} \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{-\tilde{\psi}(2n) + r_{2n}}{\psi(2n)}} D^{\frac{s_{2n}}{2}} A^{\frac{r_{2n}}{2}} \quad (\text{by Lemma 4}) \\ D^{s_{2n}} &\leq \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{\tilde{\psi}(2n) - r_{2n}}{\psi(2n)}} \dots (b). \end{aligned}$$

Put $\gamma = \frac{w}{s_{2n}} \in [0, 1]$ for $0 < w \leq s_{2n}$.

Raise each side of (b) to the power $\gamma = \frac{w}{s_{2n}} \in [0, 1]$, then we have the following inequality by Theorem 1:

$$D^w \leq \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{(\tilde{\psi}(2n) - r_{2n})\gamma}{\psi(2n)}}.$$

(i) Proof of the result that $G_{2n}(A, B, r_{2n}, s_{2n})$ is a decreasing function of s_{2n} .

$$\begin{aligned}
 &g(s_{2n}) \\
 &= \left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n}-1}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n}-1}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\psi(2n)}} \\
 &= \left\{ A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\psi(2n)}} \\
 &= \left\{ \left(A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right)^{\frac{\tilde{\gamma}(2n-1)(s_{2n}+w)+r_{2n}}{\psi(2n)}} \right\}^{\frac{\tilde{\alpha}(2n)}{\tilde{\gamma}(2n-1)(s_{2n}+w)+r_{2n}}} \\
 &= \left\{ A^{\frac{r_{2n}}{2}} D^{\frac{s_{2n}}{2}} \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{(\tilde{\psi}(2n)-r_{2n})\gamma}{\psi(2n)}} D^{\frac{s_{2n}}{2}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\tilde{\gamma}(2n-1)(s_{2n}+w)+r_{2n}}} \quad (\text{by Lemma 4}) \\
 &\geq \left(A^{\frac{r_{2n}}{2}} D^{s_{2n}+w} A^{\frac{r_{2n}}{2}} \right)^{\frac{\tilde{\alpha}(2n)}{\tilde{\gamma}(2n-1)(s_{2n}+w)+r_{2n}}} \\
 &= g(s_{2n} + w)
 \end{aligned}$$

and the last inequality holds by Theorem 1 because $\frac{\tilde{\alpha}(2n)}{\tilde{\gamma}(2n-1)(s_{2n}+w)+r_{2n}} \in [0, 1]$.

(ii) Proof of the result that $G_{2n}(A, B, r_{2n}, s_{2n})$ is a decreasing function of r_{2n} . Raise each side of (a) to the power $\frac{u}{r_{2n}} \in [0, 1]$ for $0 < u \leq r_{2n}$, then we have the following inequality by Theorem 1:

$$A^u \geq \left\{ A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{u}{\psi(2n)}}.$$

$$\begin{aligned}
 &G(A, B, r_{2n}, s_{2n}) \\
 &= A^{-\frac{r_{2n}}{2}} \left\{ A^{\frac{r_{2n}}{2}} \left(A^{-\frac{r_{2n}-1}{2}} \dots \left(A^{\frac{r_2}{2}} \left(A^{-\frac{r_1}{2}} B^{s_1} A^{-\frac{r_1}{2}} \right)^{s_2} A^{\frac{r_2}{2}} \right)^{s_3} \dots A^{-\frac{r_{2n}-1}{2}} \right)^{s_{2n}} A^{\frac{r_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)}{\psi(2n)}} A^{-\frac{r_{2n}}{2}} \\
 &= A^{-\frac{r_{2n}}{2}} \left(A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right)^{\frac{\tilde{\alpha}(2n)}{\psi(2n)}} A^{-\frac{r_{2n}}{2}} \\
 &= D^{\frac{s_{2n}}{2}} \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{\tilde{\alpha}(2n)-\tilde{\psi}(2n)}{\psi(2n)}} D^{\frac{s_{2n}}{2}} \quad (\text{by Lemma 4}) \\
 &= D^{\frac{s_{2n}}{2}} \left\{ \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}} D^{\frac{s_{2n}}{2}} \right)^{\frac{\tilde{\psi}(2n)+u}{\psi(2n)}} \right\}^{\frac{\tilde{\alpha}(2n)-\tilde{\psi}(2n)}{\psi(2n)+u}} D^{\frac{s_{2n}}{2}} \\
 &= D^{\frac{s_{2n}}{2}} \left\{ D^{\frac{s_{2n}}{2}} A^{\frac{r_{2n}}{2}} \left(A^{\frac{r_{2n}}{2}} D^{s_{2n}} A^{\frac{r_{2n}}{2}} \right)^{\frac{u}{\psi(2n)}} A^{\frac{r_{2n}}{2}} D^{\frac{s_{2n}}{2}} \right\}^{\frac{\tilde{\alpha}(2n)-\tilde{\psi}(2n)}{\psi(2n)+u}} D^{\frac{s_{2n}}{2}} \quad (\text{by Lemma 4}) \\
 &\geq D^{\frac{s_{2n}}{2}} \left(D^{\frac{s_{2n}}{2}} A^{r_{2n}+u} D^{\frac{s_{2n}}{2}} \right)^{\frac{\tilde{\alpha}(2n)-\tilde{\psi}(2n)}{\psi(2n)+u}} D^{\frac{s_{2n}}{2}} \\
 &= G(A, B, r_{2n} + u, s_{2n}).
 \end{aligned}$$

$\frac{\tilde{\alpha}(2n) - \tilde{\psi}(2n)}{\tilde{\psi}(2n) + u} \in [-1, 0]$ since $\tilde{\alpha}(2n) \leq \tilde{\psi}(2n)$. Therefore the last inequality holds by Theorem 1.

(4) \Rightarrow (3)

Put $q = 1$, $s_k = p_k$ and $r_k = t_k (k = 1, 2, \dots, 2n)$ in (4).

(3) \Rightarrow (1)

Since $F_{2k}(A, B, t_{2k}, p_{2k})$ is decreasing for t_{2k} and p_{2k} by (3), we have the following inequality: $F_{2k}(A, B, t_{2k}, p_{2k}) \leq F_{2k}(A, B, t_{2k}, 1) \leq F_{2k}(A, B, t_{2k-1}, 1)$ holds for any natural number k such that $1 \leq k \leq n$.

Therefore we have the following inequality:

$$\begin{aligned} & \left\{ A^{\frac{t_{2k}}{2}} \left(A^{-\frac{t_{2k-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2k-1}}{2}} \right)^{p_{2k}} A^{\frac{t_{2k}}{2}} \right\}^{\frac{\alpha(2k)}{\psi(2k)}} \\ & \leq A^{\frac{t_{2k} - t_{2k-1}}{2}} \left\{ A^{\frac{t_{2k-2}}{2}} \left(A^{-\frac{t_{2k-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2k-3}}{2}} \right)^{p_{2k-2}} \right. \\ & \quad \left. \times A^{\frac{t_{2k-2}}{2}} \right\}^{\frac{\alpha(2k-2)}{\psi(2k-2)}} A^{\frac{-t_{2k-1} + t_{2k}}{2}} \end{aligned}$$

holds for every natural number k such that $1 \leq k \leq n$.

So we have the following inequality:

$$\begin{aligned} & \left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} \\ & \leq A^{\frac{t_{2n} - t_{2n-1}}{2}} \left\{ A^{\frac{t_{2n-2}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-2}}{2}} \right\}^{\frac{\alpha(2n-2)}{\psi(2n-2)}} A^{\frac{t_{2n} - t_{2n-1}}{2}} \\ & \leq A^{\frac{t_{2n} - t_{2n-1} + t_{2n-2} - t_{2n-3}}{2}} \left\{ A^{\frac{t_{2n-4}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-4}}{2}} \right\}^{\frac{\alpha(2n-4)}{\psi(2n-4)}} \\ & \quad \times A^{\frac{t_{2n} - t_{2n-1} + t_{2n-2} - t_{2n-3}}{2}} \\ & \quad \vdots \\ & \leq A^{\frac{\alpha(2n) - t_2 + t_1 - 1}{2}} \left\{ A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right\}^{\frac{1 - t_1 + t_2}{(p_1 - t_1)p_2 + t_2}} A^{\frac{\alpha(2n) - t_2 + t_1 - 1}{2}} \\ & \leq A^{\frac{\alpha(2n) - 1}{2}} B A^{\frac{\alpha(2n) - 1}{2}} \\ & \leq A^{\alpha(2n)}. \quad \square \end{aligned}$$

Although we can show the following Proposition 7 by (3) of Theorem 6, we will give an alternative proof.

PROPOSITION 7. *Let n be a natural number. Let $1 \leq p_j (j = 1, 2, \dots, 2n - 1, 2n)$, $t_{2k-1} \in [0, 1]$ and $t_{2k-1} \leq t_{2k} (k = 1, 2, \dots, n)$. If $0 \leq B \leq A$ with $0 < A$, then the fol-*

lowing inequality holds:

$$\begin{aligned}
 & A^{-\frac{t_{2n}}{2}} \left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} A^{-\frac{t_{2n}}{2}} \\
 & \leq A^{-\frac{t_{2n-1}}{2}} \left\{ A^{\frac{t_{2n-2}}{2}} \left(A^{-\frac{t_{2n-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-3}}{2}} \right)^{p_{2n-2}} \right. \\
 & \quad \left. \times A^{\frac{t_{2n-2}}{2}} \right\}^{\frac{\alpha(2n-2)}{\psi(2n-2)}} A^{-\frac{t_{2n-1}}{2}}.
 \end{aligned}$$

Proof. Let $1 \leq p_j$ ($j = 1, 2, \dots, 2n - 3, 2n - 2$), $t_{2k-1} \in [0, 1]$ and $t_{2k-1} \leq t_{2k}$ ($k = 1, 2, \dots, n - 1$).

If $0 \leq B \leq A$ with $0 < A$, then we have the following inequality by Theorem 5:

$$\begin{aligned}
 & \left\{ A^{\frac{t_{2n-2}}{2}} \left(A^{-\frac{t_{2n-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-3}}{2}} \right)^{p_{2n-2}} A^{\frac{t_{2n-2}}{2}} \right\}^{\frac{\alpha(2n-2)}{\psi(2n-2)}} \\
 & \leq A^{\alpha(2n-2)}.
 \end{aligned}$$

Put

$$A_1 = A^{\alpha(2n-2)} \quad \text{and}$$

$$B_1 = \left\{ A^{\frac{t_{2n-2}}{2}} \left(A^{-\frac{t_{2n-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-3}}{2}} \right)^{p_{2n-2}} A^{\frac{t_{2n-2}}{2}} \right\}^{\frac{\alpha(2n-2)}{\psi(2n-2)}}.$$

Then we have $0 \leq B_1 \leq A_1$. Therefore we have the following inequality:

$$A_1^{-\frac{t}{2}} \left\{ A_1^{\frac{r}{2}} \left(A_1^{-\frac{s}{2}} B_1^{p_1} A_1^{-\frac{s}{2}} \right)^s A_1^{\frac{r}{2}} \right\}^{\frac{1-t+r}{(p-t)s+r}} A_1^{-\frac{t}{2}} \leq A_1^{-\frac{t}{2}} B_1 A_1^{-\frac{t}{2}}$$

holds for $t \in [0, 1]$, $1 \leq p, s$ and $t \leq r$ by (3) of Theorem 3.

Put $s = p_{2n}$, $p = \frac{\psi(2n-2)}{\alpha(2n-2)} p_{2n-1}$, $t = \frac{t_{2n-1}}{\alpha(2n-2)}$ and $r = \frac{t_{2n}}{\alpha(2n-2)}$.

Then it is obvious that

$$A_1^{\frac{t}{2}} = A^{\frac{t_{2n-1}}{2}}, \quad A_1^{\frac{r}{2}} = A^{\frac{t_{2n}}{2}},$$

$$B_1^p = \left\{ A^{\frac{t_{2n-2}}{2}} \left(A^{-\frac{t_{2n-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-3}}{2}} \right)^{p_{2n-2}} A^{\frac{t_{2n-2}}{2}} \right\}^{p_{2n-1}}$$

and

$$\begin{aligned}
 \frac{1-t+r}{(p-t)s+r} &= \frac{1 - \frac{t_{2n-1}}{\alpha(2n-2)} + \frac{t_{2n}}{\alpha(2n-2)}}{\left(\frac{\psi(2n-2)}{\alpha(2n-2)} p_{2n-1} - \frac{t_{2n-1}}{\alpha(2n-2)} \right) p_{2n} + \frac{t_{2n}}{\alpha(2n-2)}} \\
 &= \frac{\alpha(2n-2) - t_{2n-1} + t_{2n}}{(\psi(2n-2)p_{2n-1} - t_{2n-1})p_{2n} + t_{2n}} \\
 &= \frac{\alpha(2n)}{\psi(2n)}.
 \end{aligned}$$

Therefore we have the following inequality:

$$\begin{aligned}
 & A^{-\frac{t_{2n}}{2}} \left\{ A^{\frac{t_{2n}}{2}} \left(A^{-\frac{t_{2n-1}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-1}}{2}} \right)^{p_{2n}} A^{\frac{t_{2n}}{2}} \right\}^{\frac{\alpha(2n)}{\psi(2n)}} A^{-\frac{t_{2n}}{2}} \\
 \leq & A^{-\frac{t_{2n-1}}{2}} \left\{ A^{\frac{t_{2n-2}}{2}} \left(A^{-\frac{t_{2n-3}}{2}} \dots \left(A^{\frac{t_2}{2}} \left(A^{-\frac{t_1}{2}} B^{p_1} A^{-\frac{t_1}{2}} \right)^{p_2} A^{\frac{t_2}{2}} \right)^{p_3} \dots A^{-\frac{t_{2n-3}}{2}} \right)^{p_{2n-2}} \right. \\
 & \left. \times A^{\frac{t_{2n-2}}{2}} \right\}^{\frac{\alpha(2n-2)}{\psi(2n-2)}} A^{-\frac{t_{2n-1}}{2}}. \quad \square
 \end{aligned}$$

REFERENCES

- [1] T. FURUTA, $A \geq B \geq 0$ assures $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0, p \geq 0, q \geq 1$ with $(1+2r)q \geq p+2r$, Proc. Amer. Math. Soc. **101**, 1 (1987), 85–88.
- [2] T. FURUTA, Two operator functions with monotone property, Proc. Amer. Math. Soc. **111**, 2 (1991), 511–516.
- [3] T. FURUTA, Extension of the Furuta inequality and Ando-Hiai log-majorization, Linear Algebra Appl. **219** (1995), 139–155.
- [4] T. FURUTA, Further extension of an order preserving operator inequality, J. Math. Inequal. **2** (2008), 465–472.
- [5] T. FURUTA, Operator functions on chaotic order involving order preserving operator inequalities, to appear in J. Math. Inequal.
- [6] T. FURUTA, M. HASHIMOTO AND M. ITO, Equivalence relation between generalized Furuta inequality and related operator functions, Sci. Math. **1**, 2 (1998), 257–259.
- [7] T. FURUTA AND D. WANG, A decreasing operator function associated with the Furuta inequality, Proc. Amer. Math. Soc. **126**, 8 (1998), 2427–2432.
- [8] T. FURUTA, T. YAMAZAKI AND M. YANAGIDA, Operator functions implying generalized Furuta inequality, Math. Inequal. Appl. **1** (1998), 123–130.
- [9] T. FURUTA, T. YAMAZAKI AND M. YANAGIDA, Order preserving operator inequalities via Furuta inequality, Math. Japon. **48** (1998), 471–476.
- [10] E. HEINZ, Beiträge zur Störungstheorie der Spektralzerlegung, Math. Ann. **124** (1951), 415–438.
- [11] M. ITO, M. HASHIMOTO AND T. FURUTA, Equivalence relation between an order preserving operator inequality and related operator functions, Surikaiseikikenkyusho Kokyuroku No. **1080** (1999), 34–40.
- [12] T. KOIZUMI AND K. WATANABE, A remark on extension of order preserving operator inequality, to appear in J. Math. Inequal.
- [13] K. LÖWNER, Über monotone Matrixfunktionen, Math. Z. **38** (1934), 177–216.
- [14] M. UCHIYAMA, Some exponential operator inequalities, Math. Inequal. Appl. **2** (1999), 469–471.

(Received December 20, 2011)

Tatsuya Koizumi
 Graduate School of Science & Technology
 Niigata University
 950-2181, Japan
 e-mail: f10a061h@mail.cc.niigata-u.ac.jp

 Keiichi Watanabe
 Department of Mathematics, Faculty of Science
 Niigata University
 950-2181, Japan
 e-mail: wtnbk@math.sc.niigata-u.ac.jp