

OPTIMAL BOUNDS FOR TOADER MEAN IN TERMS OF ARITHMETIC AND CONTRAHARMONIC MEANS

YING-QING SONG, WEI-DONG JIANG, YU-MING CHU AND DAN-DAN YAN

Abstract. We find the greatest value α_1 and α_2 , and the least values β_1 and β_2 , such that the double inequalities $\alpha_1 C(a, b) + (1 - \alpha_1) A(a, b) < T(a, b) < \beta_1 C(a, b) + (1 - \beta_1) A(a, b)$ and $\alpha_2/A(a, b) + (1 - \alpha_2)/C(a, b) < 1/T(a, b) < \beta_2/A(a, b) + (1 - \beta_2)/C(a, b)$ hold for all $a, b > 0$ with $a \neq b$. As applications, we get new bounds for the complete elliptic integral of the second kind. Here, $C(a, b) = (a^2 + b^2)/(a + b)$, $A(a, b) = (a + b)/2$, and

$$T(a, b) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} d\theta$$

denote the contraharmonic, arithmetic, and Toader means of two positive numbers a and b , respectively.

Mathematics subject classification (2010): 26E60, 33E05.

Keywords and phrases: Contraharmonic mean, arithmetic mean, Toader mean, complete elliptic integrals.

REFERENCES

- [1] G.H. TOADER, *Some mean values related to the arithmetic-geometric mean*, J. Math. Anal. Appl., 1998, **218** (2): 358–368.
- [2] E. NEUMAN, *Bounds for symmetric elliptic integrals*, J. Approx. Theory, 2003, **122** (2): 249–259.
- [3] H. KAZI AND E. NEUMAN, *Inequalities and bounds for elliptic integrals*, J. Approx. Theory, 2007, **146** (2): 212–226.
- [4] H. KAZI AND E. NEUMAN, *Inequalities and bounds for elliptic integrals II*, Special functions and orthogonal Polynomials, 127–138, Contemp. Math., 471, Amer. Math. Soc., Providence, RI, 2008.
- [5] M.-K. WANG, Y.-M. CHU, S.-L. QIU AND Y.-P. JIANG, *Bounds for the perimeter of an ellipse*, J. Approx. Theory, 2012, **164** (7): 928–937.
- [6] Y.-M. CHU AND M.-K. WANG, *Optimal Lehmer mean bounds for the Toader mean*, Results Math., 2012, **61** (3–4): 223–229.
- [7] Y.-M. CHU, M.-K. WANG AND S.-L. QIU, *Optimal combinations bounds of root-square and arithmetic means for Toader mean*, Proc. Indian Acad. Sci. Math. Sci., 2012, **122** (1): 41–51.
- [8] Y.-M. CHU AND M.-K. WANG, *Inequalities between arithmetic-geometric, Gini, and Toader means*, Abstr. Appl. Anal., 2012, Article ID 830585, 11 pages.
- [9] Y.-M. CHU, M.-K. WANG, S.-L. QIU AND Y.-F. QIU, *Sharp generalized Seiffert mean bounds for Toader mean*, Abstr. Appl. Anal., 2011, Article ID 605259, 8 pages.
- [10] M. VUORINEN, *Hypergeometric functions in geometric function theory*, Special functions and differential equations (Madras, 1997), 119–126, Allied Publ., New Delhi, 1998.
- [11] S.-L. QIU AND J.-M. SHEN, *On two problems concerning means*, Journal of Hangzhou Institute of Electronic Engineering, 1997, **17** (3): 1–7 (in Chinese).
- [12] R. W. BARNARD, K. PEARCE AND K. C. RICHARDS, *An inequality involving the generalized hypergeometric function and the arc length of an ellipse*, SIAM J. Math. Anal., 2000, **31** (3): 693–699.
- [13] H. ALZER AND S.-L. QIU, *Monotonicity theorems and inequalities for the complete elliptic integrals*, J. Comput. Appl. Math., 2004, **172** (2): 289–312.

- [14] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.