

OPTIMAL BOUNDS FOR NEUMAN-SÁNDOR MEAN IN TERMS OF THE CONVEX COMBINATION OF LOGARITHMIC AND QUADRATIC OR CONTRA-HARMONIC MEANS

YUMING CHU, TIEHONG ZHAO AND BAORYU LIU

Abstract. In this article, we present the least values α_1 , α_2 , and the greatest values β_1 , β_2 such that the double inequalities

$$\alpha_1 L(a, b) + (1 - \alpha_1) Q(a, b) < M(a, b) < \beta_1 L(a, b) + (1 - \beta_1) Q(a, b)$$

$$\alpha_2 L(a, b) + (1 - \alpha_2) C(a, b) < M(a, b) < \beta_2 L(a, b) + (1 - \beta_2) C(a, b)$$

hold for all $a, b > 0$ with $a \neq b$, where $L(a, b)$, $M(a, b)$, $Q(a, b)$ and $C(a, b)$ are respectively the logarithmic, Neuman-Sándor, quadratic and contra-harmonic means of a and b .

Mathematics subject classification (2010): 26E60.

Keywords and phrases: Neuman-Sándor mean, logarithmic mean, quadratic mean, contra-harmonic mean.

REFERENCES

- [1] E. NEUMAN AND J. SÁNDOR, *On the Schwab-Borchardt mean*, Math. Pannon. **14**, 2(2003), 253–266.
- [2] E. NEUMAN AND J. SÁNDOR, *On the Schwab-Borchardt mean II*, Math. Pannon. **17**, 1 (2006), 49–59.
- [3] Y.-M. LI, B.-Y. LONG AND Y.-M. CHU, *Sharp bounds for the Neuman-Sándor mean in terms of generalized logarithmic mean*, J. Math. Inequal. **6**, 4 (2012), 567–577.
- [4] E. NEUMAN, *A note on a certain bivariate mean*, J. Math. Inequal. **6**, 4 (2012), 637–643.
- [5] E. NEUMAN AND J. SÁNDOR, *Bounds for the quotients of differences of certain bivariate means*, Adv. Stud. Contemp. Math. **23**, 1 (2013), 61–67.