

## SOME GENERALIZATIONS OF OPERATOR INEQUALITIES

XIAOHUI FU

*Abstract.* In this paper, we generalize some operator inequalities as follows: Let  $A, A_i$  ( $i = 1, \dots, n$ ) be positive operators on a Hilbert space with  $0 < m \leq A, A_i \leq M$  ( $i = 1, \dots, n$ ). Then for  $1 \leq p < \infty$  and every positive unital linear map  $\Phi$ ,

$$\Phi^p(A^{-1})\Phi^p(A) + \Phi^p(A)\Phi^p(A^{-1}) \leq \frac{(M+m)^{2p}}{2M^p m^p},$$

and

$$\left(\frac{A_1 + \dots + A_n}{n}\right)^{2p} \leq \left(\frac{(M+m)^{2p}}{4M^p m^p}\right)^2 G^{2p}(A_1, \dots, A_n),$$

where  $G(A_1, \dots, A_n)$  is Ando-Li-Mathias geometric mean [1].

*Mathematics subject classification* (2010): 47A63.

*Keywords and phrases:* Operator inequalities, positive linear maps, arithmetic mean, geometric mean.

### REFERENCES

- [1] T. ANDO, C.-K. LI AND R. MATHIAS, *Geometric means*, Linear Algebra Appl. **385** (2004), 305–334.
- [2] R. BHATIA, *Positive definite matrices*, Princeton University Press, 2007.
- [3] R. BHATIA, F. KITTANEH, *Notes on matrix arithmetic-geometric mean inequalities*, Linear Algebra Appl. **308** (2000), 203–211.
- [4] J. I. FUJII ET AL., *A reverse of the weighted geometric mean due to Lawson-Lim*, Linear Algebra Appl. **427** (2007), 272–284.
- [5] M. LIN, *On an operator Kantorovich inequality for positive linear maps*, J. Math. Anal. Appl. **402** (2013), 127–132.
- [6] M. LIN, *Squaring a reverse AM-GM inequality*, Studia Math. **215** (2013), 187–194.
- [7] X. ZHAN, *Matrix Theory*, Beijing: Higher Education Press, 2008. (In Chinese)